

# Pacific Journal of Mathematics

## **SPHERE TRANSITIVE STRUCTURES AND THE TRIALITY AUTOMORPHISM**

ALFRED GRAY AND PAUL STEPHEN GREEN

## SPHERE TRANSITIVE STRUCTURES AND THE TRIALITY AUTOMORPHISM

ALFRED GRAY AND PAUL GREEN

Let  $G$  be a compact connected Lie group which acts transitively and effectively on a sphere  $S^{n-1}$ . A manifold  $M$  is said to have a sphere transitive structure if the structure group of the tangent bundle of  $M$  can be reduced from  $O(n)$  to  $G$ . The study of the existence of such structures is a generalization of the well-known problem of the existence of almost complex structures. We completely solve the question of existence of sphere transitive structures on spheres.

For our study of sphere transitive structures we need to know some facts about the triality automorphism  $\lambda$  of  $\text{Spin}(8)$ . We completely determine the cohomology homomorphism induced by  $\lambda$  on the cohomology of the classifying space of  $\text{Spin}(8)$ .

Berger [1] has classified the holonomy groups of manifolds having an affine connection with zero torsion. Either from this classification or directly from Simons [11], it follows that the holonomy group of an irreducible Riemannian manifold which is not a symmetric space acts transitively on a sphere.

On the other hand we have the following elementary fact: if the holonomy group of a Riemannian manifold  $M$  is  $G$ , then the structure group of the tangent bundle of  $M$  can be reduced to  $G$ . Therefore a more fundamental question than whether or not a Riemannian manifold  $M$  has a given Lie group  $G$  as its holonomy group is the question of the reduction of the structure group of the tangent bundle of  $M$  to  $G$ . In this paper we consider the latter question and give some necessary conditions and some sufficient conditions in terms of characteristic classes. From the remarks above it suffices to consider the case when  $G$  is a connected Lie group which acts transitively and effectively on a sphere.

We introduce the following notions.

DEFINITIONS. Let  $\xi = (E, M, p, F)$  be a vector bundle where  $M$  is a  $CW$ -complex and  $\dim F = n$ . Then a *sphere transitive reduction* is a reduction of the structure group  $O(n)$  of  $\xi$  to a connected Lie subgroup  $G$  of  $O(n)$  which acts transitively and effectively on the sphere  $S^{n-1}$ . In the special case when  $\xi$  is the tangent bundle of  $M$  we call the reduction a *sphere transitive structure* on  $M$ .

According to [10] the connected Lie groups  $G$  which act effectively and transitively on spheres are the following:  $SO(n)$ ,  $U(n)$ ,

$SU(n)$ ,  $Sp(n)$ ,  $Sp(n) \cdot SO(2)$ ,  $Sp(n) \cdot Sp(1)$ ,  $G_2$ ,  $Spin(7)$ , and  $Spin(9)$ . We have

$$\begin{aligned} SO(n)/SO(n-1) &= S^{n-1}, \quad U(n)/U(n-1) = SU(n)/SU(n-1) = S^{2n-1}, \\ Sp(n)/Sp(n-1) &= Sp(n) \cdot SO(2)/Sp(n-1) \cdot SO(2) \\ &= Sp(n) \cdot Sp(1)/Sp(n-1) \cdot Sp(1) = S^{4n-1}, \\ G_2/SU(3) &= S^6, \quad Spin(7)/G_2 = S^7, \quad Spin(9)/Spin(7) = S^{15}. \end{aligned}$$


In § 2 we discuss the triality automorphism  $\lambda$  of  $Spin(8)$  and the cohomology of the self homeomorphism of the classifying space induced by  $\lambda$ . The results of § 2 are then used in § 3 to determine the cohomology of the classifying space  $B Spin(n)$  ( $n = 7, 8, 9$ ) and a good deal of the cohomology of  $BG_2$ . Then we determine some necessary conditions for sphere transitive reductions for the cases  $G = G_2, Spin(7), Spin(9)$ . In § 4 we discuss the existence of sphere transitive structures on certain homogeneous spaces. In particular we completely solve the problem of the existence of sphere transitive structures on spheres.

2. The cohomology of the triality automorphism.  $Spin(8)$  is the simply connected compact Lie group whose Lie algebra is of type  $D_4$ . Now  $D_4$  is the unique simple Lie algebra with an outer automorphism of order 3. In fact, if  $Aut(D_4)$  (resp.  $Inn(D_4)$ ) denotes the group of all (resp. inner) automorphisms of  $D_4$ , then the factor group  $Aut(D_4)/Inn(D_4)$  is isomorphic to the symmetric group on 3 letters. Let  $\kappa, \lambda \in Aut(D_4)$  be such that their images in  $Aut(D_4)/Inn(D_4)$  generate this group and satisfy the relations  $\lambda^3 = 1, \kappa^2 = 1, \kappa\lambda\kappa = \lambda^2$ .

According to [7] it is possible to choose  $\kappa$  and  $\lambda$  so that the *principle of triality* holds. This means the following. Let  $V$  be the 8-dimensional algebra of Cayley numbers and denote the product of  $x, y \in V$  by  $xy$ . Then for  $A \in D_4, x, y \in V$  we have

$$(Ax)y + x(\lambda(A)y) = ((\lambda\kappa)(A))(xy).$$

The Dynkin diagram of  $D_4$  is



where  $\{\gamma_1, \gamma_2, \gamma_3, \gamma_4\}$  is a simple system of roots of  $D_4$ . Since  $\kappa$  and  $\lambda$  are outer, they give rise to symmetries of the Dynkin diagram of  $D_4$ . It may be checked that  $\{\gamma_1, \gamma_2, \gamma_3, \gamma_4\}$  may be chosen so that

$\kappa(\gamma_1) = \gamma_1$ ,  $\kappa(\gamma_2) = \gamma_2$ ,  $\kappa(\gamma_3) = \gamma_4$ ,  $\kappa(\gamma_4) = \gamma_3$ ,  $\lambda(\gamma_1) = \gamma_3$ ,  $\lambda(\gamma_2) = \gamma_2$ ,  $\lambda(\gamma_3) = \gamma_4$ ,  $\lambda(\gamma_4) = \gamma_1$ . Henceforth we assume that the principal of triality holds and that the above choice of simple roots has been made.

Since  $\text{Spin}(8)$  is simply connected,  $\lambda$  and  $\kappa$  induce outer automorphisms of  $\text{Spin}(8)$ ; these in turn induce homeomorphisms of  $B\text{Spin}(8)$ , which we continue to denote by  $\lambda$  and  $\kappa$ . In order to determine the cohomology of  $\lambda$  and  $\kappa$ , it will be convenient to use some cohomology classes introduced by Thomas [12]. Let  $p: B\text{Spin}(n) \rightarrow BSO(n)$  be the map defined by the covering homomorphism of  $\text{Spin}(n)$  over  $SO(n)$ . Denote by  $w_i$  the universal Stiefel-Whitney classes, by  $P_i$  the universal Pontryagin classes, and by  $X$  the Euler class of  $BSO(8)$ . Then  $H^*(BSO(8), \mathbf{Z}) = \mathbf{Z}[P_1, P_2, P_3, X] + 2\text{-torsion}$  and  $H^*(BSO(8), \mathbf{Z}_2) = \mathbf{Z}_2[w_2, \dots, w_8]$ . According to Thomas [12] there exist cohomology classes  $Q_i \in H^*(B\text{Spin}, \mathbf{Z})$  ( $i = 1, 2, 3, 4$ ) and  $w_i^* \in H^*(B\text{Spin}, \mathbf{Z}_2)$  ( $i = 4, 6, 7, 8$ ) (where  $\text{Spin}$  denotes the stable Spin group) such that

$$\begin{aligned}
 p^*(P_1) &= 2Q_1 & p^*(w_i) &= w_i^* \quad (i = 4, 6, 7, 8) \\
 p^*(P_2) &= 2Q_2 + Q_1^2 & p^*(w_i) &= 0 \quad (i = 2, 3, 5) \\
 p^*(P_3) &= Q_3 & \rho_2(Q_1) &= w_4^*, \quad \rho_2(Q_2) = w_8^* \\
 p^*(P_4) &= 2Q_4 + Q_2^2 & \rho_2(Q_3) &= w_6^{*2}, \quad \rho_4(Q_4) = w_{16}^* .
 \end{aligned}$$

The cohomology classes  $Q_1, Q_2, Q_3, w_4^*, w_6^*, w_7^*, w_8^*$  give rise to the cohomology classes in  $H^*(B\text{Spin}(8), \mathbf{Z})$  and  $H^*(B\text{Spin}(8), \mathbf{Z}_2)$  which we denote by the same letters.

**THEOREM 2.1.** (i) *There exist*

$$Y \in H^8(B\text{Spin}(8), \mathbf{Z}) \quad \text{and} \quad \omega \in H^8(B\text{Spin}(8), \mathbf{Z}_2)$$

such that

$$\begin{aligned}
 H^*(B\text{Spin}(8), \mathbf{Z}) &= \mathbf{Z}[Q_1, Q_2, Q_3, Y] + 2\text{-torsion} \\
 H^*(B\text{Spin}(8), \mathbf{Z}_2) &= \mathbf{Z}_2[w_4^*, w_6^*, w_7^*, w_8^*, \omega] .
 \end{aligned}$$

Furthermore  $Y$  and  $\omega$  can be chosen so that  $p^*(X) = 2Y - Q_2$  and  $\rho_2(Y) = \omega$ .

(ii) *The cohomology homomorphisms  $\lambda^*$  and  $\kappa^*$  are given as follows:*

$$\begin{aligned}
 \lambda^*(Q_1) &= Q_1, & \lambda^*(w_i^*) &= w_i^* \quad (i = 4, 6, 7), \\
 \lambda^*(Q_2) &= 3Y - 2Q_2, & \lambda^*(w_8^*) &= \omega, \\
 \lambda^*(Y) &= Y - Q_2, & \lambda^*(\omega) &= w_8^* + \omega, \\
 \lambda^*(Q_3) &= Q_3 + 2Q_1Y - 2Q_1Q_2, & \kappa^*(w_i^*) &= w_i^* \quad (i = 4, 6, 7, 8), \\
 \kappa^*(Q_i) &= Q_i \quad (i = 1, 2, 3), & \kappa^*(\omega) &= w_8^* + \omega, \\
 \kappa^*(Y) &= -Y + Q_2.
 \end{aligned}$$

Before proving this theorem we state without proof a lemma which we shall need.

LEMMA 2.2. *Let  $s: K \rightarrow L$  be a  $p^n$ -fold covering of a compact connected Lie group where  $p$  is a prime, and denote by*

$$s^*: H^*(BL, \mathbf{Z}) \longrightarrow H^*(BK, \mathbf{Z})$$

*the corresponding cohomology homomorphism of classifying spaces. Let  $S$  be a subset of  $H^*(BK, \mathbf{Z})$  such that  $S$  generates  $s^*(H^*(BL, \mathbf{Z}))$  as a group (ring) and  $\rho_p(S)$  generates  $\rho_p(H^*(BK), \mathbf{Z}) \cong H^*(BK, \mathbf{Z}_p)$  as a group (ring). ( $\rho_p$  denotes reduction mod  $p$ .) Then  $S$  generates  $H^*(BK, \mathbf{Z})$  as a group (ring).*

*Proof of Theorem 2.1.* Using a result of Borel [2] it is not hard to see that  $w_4^*$ ,  $w_6^*$ ,  $w_7^*$ , and  $w_8^*$  are generators of

$$H^*(B \text{Spin}(8), \mathbf{Z}_2).$$

Furthermore if  $\rho_0: \mathbf{Z} \rightarrow \mathbf{R}_0$  denotes the inclusion, where  $\mathbf{R}_0$  is the rationals, then it is obvious that

$$H^*(B \text{Spin}(8), \mathbf{R}_0) = \mathbf{R}_0[\rho_0(Q_1), \rho_0(Q_2), \rho_0(Q_3), \rho_0(p^*(X))].$$

We first establish part of (ii). The automorphism  $\kappa$  of  $\text{Spin}(8)$  gives rise to an outer automorphism  $\tilde{\kappa}$  of  $SO(8)$ ; this is the ordinary orientation reversing automorphism of  $SO(8)$ . The induced homomorphism  $\tilde{\kappa}^*$  is the identity on  $H^*(BSO(8), \mathbf{Z}_2)$  and satisfies  $\tilde{\kappa}^*(P_i) = P_i$  ( $i = 1, 2, 3$ ),  $\tilde{\kappa}^*(X) = -X$ . Hence  $\kappa^*(w_i^*) = w_i^*$  ( $i = 4, 6, 7, 8$ ), and  $\kappa^*(Q_i) = Q_i$  ( $i = 1, 2, 3$ ). It is also easy to see that  $\lambda^*(Q_1) = Q_1$  and  $\lambda^*(w_i^*) = w_i^*$  for  $i = 4, 6, 7$ .

We may write

$$\begin{aligned} \lambda^*(P^*(\rho_0(X))) &= a\rho_0(X) + b\rho_0(Q_2) + c\rho_0(Q_2^2), \\ \lambda^*(\rho_0(Q_2)) &= d\rho_0(X) + e\rho_0(Q_2) + f\rho_0(Q_2^2), \end{aligned}$$

where  $a, b, c, d, e, f$  are rational numbers. Using the facts that  $\lambda^*(Q_1^2) = Q_1^2$ ,  $\lambda^3 = 1$ ,  $\kappa\lambda\kappa = \lambda^2$ , and the knowledge of  $\kappa^*$ , we calculate that  $c = f = 0$ ,  $a = e = -1/2$ , and  $bd = -3/4$ .

To compute  $b, d$ , and  $\lambda^*(\rho_0(Q_3))$  we must resort to some calculations with roots. Let  $\tilde{Q}_1, \tilde{Q}_2, \tilde{Q}_3$ , and  $\tilde{X}$  denote the real cohomology classes corresponding to  $Q_1, Q_2, Q_3$ , and  $p^*(X)$ . Then we may regard  $\tilde{Q}_1, \tilde{Q}_2, \tilde{Q}_3$  and  $\tilde{X}$  as polynomials on the Lie algebra of a maximal torus of  $\text{Spin}(8)$ , i.e., polynomials in the roots of  $\text{Spin}(8)$ . A calculation shows in fact that (if we write  $\gamma_0 = -\gamma_1 - 2\gamma_2 - \gamma_3 - \gamma_4$ ),

$$\begin{aligned}
 \tilde{Q}_1 &= -2\varepsilon(\gamma_0^2 + \gamma_1^2 + \gamma_3^2 + \gamma_4^2), \\
 \tilde{Q}_2 &= \varepsilon^2(-\gamma_0^2\gamma_1^2 - 2\gamma_3^2\gamma_4^2 + \gamma_0^2\gamma_3^2 + \gamma_0^2\gamma_4^2 + \gamma_1^2\gamma_3^2 + \gamma_1^2\gamma_4^2), \\
 \tilde{X} &= \varepsilon^2(-\gamma_0^2\gamma_3^2 - \gamma_1^2\gamma_4^2 + \gamma_0^2\gamma_4^2 + \gamma_1^2\gamma_3^2), \\
 \tilde{Q}_3 &= -2\varepsilon^3(\gamma_0^4\gamma_3^2 + \gamma_0^4\gamma_4^2 + \gamma_1^4\gamma_3^2 + \gamma_1^4\gamma_4^2 + \gamma_3^4\gamma_0^2 + \gamma_3^4\gamma_1^2 + \gamma_4^4\gamma_0^2 + \gamma_4^4\gamma_1^2 \\
 &\quad - 2\gamma_0^2\gamma_1^2\gamma_3^2 - 2\gamma_0^2\gamma_1^2\gamma_4^2 - 2\gamma_0^2\gamma_3^2\gamma_4^2 - 2\gamma_1^2\gamma_3^2\gamma_4^2).
 \end{aligned}$$

Thus we obtain

$$(*) \quad \begin{cases} \lambda^*(\rho_0(X)) = -\frac{1}{2}\rho_0(X) - \frac{1}{2}\rho_0(Q_2), \\ \lambda^*(\rho_0(Q_2)) = \frac{3}{2}\rho_0(X) - \frac{1}{2}\rho_0(Q_2), \\ \lambda^*(\rho_0(Q_3)) = \rho_0(Q_3) + \rho_0(Q_1X) - \rho_0(Q_1Q_2). \end{cases}$$

Define  $Y = -\lambda^*(p^*(X))$  and  $\omega = \rho_2(Y)$ . Then  $\lambda^*(w_8^*) = \omega$ . From this, equations (\*), and the fact that  $H^*(B\text{Spin}(8), \mathbf{Z})$  has only 2-torsion, we obtain the rest of (ii).

From (ii) and Borel [2] we see that  $\omega$  may be taken to be the remaining generator of  $H^*(B\text{Spin}(8), \mathbf{Z}_2)$ . This fact together with (ii) and Lemma 2.2 imply (i).

**3. The cohomology of  $B\text{Spin}(7)$ ,  $B\text{Spin}(9)$ , and  $BG_2$ .** We first compute the cohomology of  $B\text{Spin}(7)$  and its inclusion in  $BSO(8)$ . Actually there are two natural 8-dimensional representations of  $\text{Spin}(7)$  according to [8]. These are equivalent in  $O(8)$  but not in  $SO(8)$ . Denote these representations by  $j_+$  and  $j_-$ . In the terminology of [8]  $j_+$  and  $j_-$  give rise to the two distinct 3-fold vector cross products on  $R^8$ . Let  $i: \text{Spin}(7) \rightarrow \text{Spin}(8)$  be the natural inclusion. The following lemma [8], [13] will be necessary.

LEMMA 3.1. *We have the following commutative diagrams*

$$\begin{array}{ccc}
 \text{Spin}(8) & \xrightarrow{\lambda} & \text{Spin}(8) \\
 i \uparrow & & \downarrow p \\
 \text{Spin}(7) & \xrightarrow{j_+} & SO(8)
 \end{array}
 \qquad
 \begin{array}{ccc}
 \text{Spin}(8) & \xrightarrow{\lambda^2} & \text{Spin}(8) \\
 i \uparrow & & \downarrow p \\
 \text{Spin}(7) & \xrightarrow{j_-} & SO(8).
 \end{array}$$

Where it is convenient we write  $j_{\pm}$  to mean either  $j_+$  or  $j_-$ . Let  $i^*: H^*(B\text{Spin}(8)) \rightarrow H^*(B\text{Spin}(7))$  and

$$j_{\pm}^*: H^*(BSO(8)) \rightarrow H^*(B\text{Spin}(7))$$

be the induced cohomology homomorphisms of  $i$  and  $j_{\pm}$  on classifying spaces.

**THEOREM 3.2.** *Identify  $i^*(w_i^*)$  with  $w_i^*$  ( $i = 4, 6, 7$ ),  $i^*(\omega)$  with  $\omega$ ,  $i^*(Q_i)$  with  $Q_i$  ( $i = 1, 3$ ) and  $i^*(Y)$  with  $Y$ . Then we have*

$$(i) \quad H^*(B \operatorname{Spin}(7), \mathbf{Z}) = \mathbf{Z}[Q_1, Q_3, Y] + 2\text{-torsion},$$

$$H^*(B \operatorname{Spin}(7), \mathbf{Z}_2) = \mathbf{Z}_2[w_4^*, w_6^*, w_7^*, \omega];$$

$$(ii) \quad i^*(w_8^*) = 0 \text{ and } i^*(Q_2) = 2Y;$$

$$(iii) \quad j_{\pm}^*(P_1) = 2Q_1, \quad j_{\pm}^*(w_i) = w_i^* \quad (i = 4, 6, 7)$$

$$j_{\pm}^*(P_2) = -2Y + Q_1^2, \quad j_{\pm}^*(w_8^*) = \omega;$$

$$j_{\pm}^*(P_3) = Q_3 - 2Q_1Y,$$

$$j_{\pm}^*(X) = \mp Y,$$

(iv) *The kernel of  $j_{\pm}^*$  on integral cohomology is the ideal generated by  $4P_2 - P_1^2 \mp 8X$ .*

*Proof.* Since  $i: \operatorname{Spin}(7) \rightarrow \operatorname{Spin}(8)$  covers the ordinary inclusion of  $SO(7)$  in  $SO(8)$ , we have  $(i^* \circ p^*)(X) = 0$ . Thus  $i^*(Q_2) = 2Y$ . From this fact, Theorem 2.1 and Lemma 2.2 we obtain (i) and (ii). Furthermore (iii) follows from (i), (ii), and Lemma 3.1; finally (iv) is an easy calculation from (iii).

Let  $M$  be a  $CW$ -complex and let  $\xi$  be an oriented vector bundle over  $M$  with fiber dimension 8. Denote by  $f: M \rightarrow BSO(8)$  the classifying map determined by  $\xi$ . We shall say that  $\xi$  admits a *nontransitive*  $\operatorname{Spin}(7)$  reduction if  $f = p \circ i \circ g$  for some  $g: M \rightarrow B \operatorname{Spin}(7)$ :

$$\begin{array}{ccc} B \operatorname{Spin}(7) & \xrightarrow{i} & B \operatorname{Spin}(8) \\ g \uparrow & & \downarrow p \\ M & \xrightarrow{f} & BSO(8). \end{array}$$

(Here  $i$  and  $p$  denote the maps induced by the maps  $\operatorname{Spin}(7) \rightarrow \operatorname{Spin}(8)$  and  $\operatorname{Spin}(8) \rightarrow SO(8)$  which we also designate by  $i$  and  $p$ .) On the other hand by Lemma 3.1,  $M$  admits a sphere transitive  $\operatorname{Spin}(7)$  reduction in the sense of this paper if and only if for some  $g: M \rightarrow B \operatorname{Spin}(7)$  we have  $f = p \circ \lambda \circ i \circ g$  or  $f = p \circ \lambda^2 \circ i \circ g$ . Therefore we have the following lemma.

**LEMMA 3.3.** *Assume  $w_2(\xi) = 0$ . Then  $\xi$  has a transitive  $\operatorname{Spin}(7)$  reduction (that is a reduction of  $SO(8)$  to  $j_{\pm}(\operatorname{Spin}(7))$ ) if and only if  $\lambda^{\mp 1}(\xi)$  has a nontransitive  $\operatorname{Spin}(7)$  reduction.*

Next we determine the primary and secondary obstructions to the existence of sphere transitive  $\operatorname{Spin}(7)$  structures.

**THEOREM 3.4.** *Let  $M$  be a  $CW$ -complex and let  $\xi$  be an oriented vector bundle over  $\xi$  with fiber dimension 8. Denote by  $c^2(\xi)$  and*

$c^s(\xi)$  the primary and secondary obstructions to the existence of a transitive  $j_{\pm}(\text{Spin}(7))$  structure. Then  $c^2(\xi) \in H^2(M, \mathbf{Z}_2)$ ,  $c^8(\xi) \in H^8(M, \mathbf{Z})$ , and we have

$$c^2(\xi) = w_2(\xi) ,$$

$$16c^8(\xi) = 4P_2(\xi) - P_1^2(\xi) \pm 8X(\xi) .$$

*Proof.* We first note that  $SO(8)/\text{Spin}(7)$  is diffeomorphic to real projective space  $P^7$ . Hence  $c^2(\xi) \in H^2(M, \pi_1(P^7)) = H^2(M, \mathbf{Z}_2)$  and  $c^8(\xi) \in H^8(M, \pi_7(P^7)) = H^8(M, \mathbf{Z})$ . A transgression argument given in [8] shows that  $w_2(\xi) = c^2(\xi)$ .

Assume that  $w_2(\xi) = 0$ . By Lemma 3.3,  $\xi$  has a sphere transitive  $j_{\pm}(\text{Spin}(7))$  structure if and only if  $\lambda^{\mp 1}(\xi)$  has a nontransitive  $\text{Spin}(7)$  structure. The first obstruction to the latter is  $X(\lambda^{\mp 1}(\xi))$ , as is well-known. On the other hand by Theorem 2.1 and 3.2 we have

$$16X^{\pm 1}(\lambda(\xi)) = 4P_2(\xi) - P_1^2(\xi) \mp X(\xi) .$$

Hence the theorem follows.

**COROLLARY 3.5.** *Let  $\xi$  be an oriented vector bundle with fiber dimension 8 over a CW-complex  $M$ . Assume that  $\dim M \leq 8$  and that  $H_8(M, \mathbf{Z})$  has no 2-torsion. Then  $\xi$  has a sphere transitive  $j_{\pm}(\text{Spin}(7))$  structure if and only if  $w_2(\xi) = 0$  and*

$$4P_2(\xi) - P_1^2(\xi) \pm X(\xi) = 0 .$$

Theorems 2.1 and 3.4 and Corollary 3.5 correct an error in [8]. We now turn to  $\text{Spin}(9)$ . First we need a lemma.

**LEMMA 3.6.** *We have the following commutative diagram:*

$$\begin{array}{ccccc}
 \text{Spin}(8) \times \text{Spin}(8) & \xrightarrow{\Delta} & \text{Spin}(16) & & \\
 \lambda \times \lambda^2 \nearrow & & \downarrow p & & \\
 \text{Spin}(8) & \xrightarrow{k} & \text{Spin}(9) & \xrightarrow{l} & SO(16)
 \end{array}$$

where  $\Delta$  is the standard map of  $\text{Spin}(8) \times \text{Spin}(8)$  into  $\text{Spin}(16)$ ,  $p$  is the covering projection,  $k$  is the standard inclusion of  $\text{Spin}(8)$  in  $\text{Spin}(9)$ , and  $l$  is the sphere transitive 16-dimensional representation of  $\text{Spin}(9)$ .

*Proof.* Let  $F_4$  denote the automorphism group of the exceptional Jordan algebra of  $3 \times 3$  Hermitian matrices of Cayley numbers. Let



$$E_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad E_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad E_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

The subgroup  $H_i$  of  $F_4$  which leaves  $E_i$  fixed is isomorphic to  $\text{Spin}(9)$  (see [7]). On the other hand  $\text{Spin}(8)$  is isomorphic to  $H_1 \cap H_2 \cap H_3$ . Let

$$V_1 = \text{matrices of the form } \begin{pmatrix} 0 & z & w \\ \bar{z} & 0 & 0 \\ \bar{w} & 0 & 0 \end{pmatrix},$$

$$V_2 = \text{matrices of the form } \begin{pmatrix} 0 & z & 0 \\ \bar{z} & 0 & w \\ 0 & \bar{w} & 0 \end{pmatrix},$$

$$V_3 = \text{matrices of the form } \begin{pmatrix} 0 & 0 & z \\ 0 & 0 & w \\ \bar{z} & \bar{w} & 0 \end{pmatrix}.$$

Then  $V_i$  is an irreducible representation space for  $H_i$ . Since there is only one irreducible 16-dimensional representation of  $\text{Spin}(9)$ , each representation of  $H_i$  on  $V_i$  is just  $1$ . Now the representation of  $\text{Spin}(8)$  on  $V_1$  is  $\lambda \times 1$ , on  $V_2$  is  $\lambda \times \lambda^2$ , and on  $V_3$  is  $1 \times \lambda^2$ . Hence we get the commutative diagram

$$\begin{array}{ccccc} & & \text{Spin}(8) \times \text{Spin}(8) & \longrightarrow & \text{Spin}(16) \\ & \nearrow^{\lambda^{i-1} \times \lambda^i} & & & \downarrow p \\ \text{Spin}(8) & \xrightarrow{k_i} & \text{Spin}(9) & \xrightarrow{l} & SO(16). \end{array}$$

We claim that  $k_2$  is the standard inclusion of  $\text{Spin}(8)$  in  $\text{Spin}(9)$  while  $k_1$  and  $k_3$  are not. This may be proved by showing that  $k_i^*(H^*(B\text{Spin}(9), \mathbf{R}_0))$  is  $\mathbf{R}_0[P_1, P_2, P_3, X^2] \subseteq H^*(B\text{Spin}(8), \mathbf{R}_0)$  for  $i = 2$ , but not for  $i = 1$  or  $3$ . (See the proof of the next theorem.) This completes the proof of the lemma.

**THEOREM 3.7.** (i) *There exist cohomology classes*

$$Z \in H^{16}(B\text{Spin}(9), \mathbf{Z})$$

and  $\phi \in H^{16}(B\text{Spin}(9), \mathbf{Z}_2)$  such that

$$H^*(B\text{Spin}(9), \mathbf{Z}) = \mathbf{Z}[Q_1, Q_2, Q_3, Z] + 2\text{-torsion}$$

$$H^*(B\text{Spin}(9), \mathbf{Z}_2) = \mathbf{Z}_2[w_4, w_6, w_7, w_8, \phi].$$

Here  $k_2^*(Q_i) = Q_i$  ( $i = 1, 2, 3$ ),  $k_2^*(4Z) = p^*(X^2 - P_2^2)$ ,  $k_2^*(w_i^*) = w_i^*$  ( $i = 4, 6, 7, 8$ ), and  $k_2^*(\phi) = \omega^2 + \omega w_8^*$ .

(ii) We have (modulo elements of order 2)

$$l^*(P_1) = 4Q_1$$

$$l^*(P_2) = -2Q_2 + 6Q_1^2$$

$$l^*(P_3) = 2Q_3 - 6Q_1Q_2 + 4Q_1^2$$

$$l^*(P_4) = -34Z - 7Q_2^2 + 4Q_1Q_3 - 6Q_1^2Q_2 + Q_1^4$$

$$l^*(P_5) = 28Q_1Z - 2Q_2Q_3 + 2Q_1^2Q_3 + 10Q_1Q_2^2 - 2Q_1^3Q_2$$

$$l^*(P_6) = 22Q_2Z - 2Q_1^2Z + Q_3^2 - 2Q_1Q_2Q_3 + 5Q_2^3 + Q_1^2Q_2^2$$

$$l^*(P_7) = 2Q_3Z - 10Q_1Q_2Z + Q_2^2Q_3 - 3Q_1Q_2^3$$

$$l^*(X) = Z.$$

and

$$l^*(w_i) = 0 \text{ for } i = 1, 2, 3, 4, 5, 6, 7, 9, 10, 11, 13$$

$$l^*(w_8) = w_4^{*2} + w_8^*$$

$$l^*(w_{12}) = w_6^{*2} + w_4^*w_8^*$$

$$l^*(w_{14}) = w_7^{*2} + w_6^*w_8^*$$

$$l^*(w_{15}) = w_7^*w_8^*$$

$$l^*(w_{16}) = \phi.$$

*Proof.* Let  $\xi$  be an 8-dimensional vector bundle with  $w_2(\xi) = 0$  and set  $\nu = \lambda(\xi) \oplus \lambda^2(\xi)$ . Then the Pontryagin, Euler, and Stiefel-classes of  $\nu$  may be computed by means of the Whitney sum formula together with Theorem 2.1. On the other hand any maximal torus (maximal 2-subgroup) of  $\text{Spin}(8)$  is also a maximal torus (maximal 2-subgroup) of  $\text{Spin}(9)$ . Therefore the formulas for above mentioned characteristic classes are the most general possible.

Set  $Z = l^*(X)$  and  $\phi = l^*(w_{16})$ . Then we obtain (ii). Finally (i) follows from (ii) and Lemma 2.2.

Theoretically the kernel of  $l^*$  can be determined from Theorem 3.7 (ii). This yields some necessary conditions that a 16-dimensional vector bundle have a transitive  $\text{Spin}(9)$  reduction. However, we omit the details. In the only example we consider in § 4, namely the Cayley plane, it is simpler to use Theorem 3.7 itself.

We conclude this section by noting a few facts about the cohomology of  $BG_2$  and its inclusion in  $B\text{Spin}(7)$ .

LEMMA 3.8. *Let  $g$  be the standard inclusion of  $G_2$  in  $SO(7)$ , and denote by  $h$  the lifting of  $g$  into  $\text{Spin}(7)$ :*

$$\begin{array}{ccc}
 & \text{Spin (7)} & \xrightarrow{i} \text{Spin (8)} \\
 & \nearrow h & \downarrow \\
 G_2 & \xrightarrow{g} & SO(7) .
 \end{array}$$

If  $i$  denotes the standard inclusion of Spin (7) in Spin (8), then we have

$$\lambda \circ i \circ h = i \circ h .$$

*Proof.* This follows from the fact that  $G_2$  is the fixed point set of  $\lambda$ .

**THEOREM 3.9.** (i) *We have*

$$\begin{aligned}
 H^*(BG_2, \mathbf{R}_0) &= \mathbf{R}_0[g^*(P_1), g^*(P_3)] \\
 &= \mathbf{R}_0[h^*(Q_1), h^*(Q_3)] ,
 \end{aligned}$$

where  $g^*$  and  $h^*$  are induced by  $g$  and  $h$  defined in the previous lemma and  $\mathbf{R}_0$  denotes the rationals.

(ii) *In integral cohomology, the kernel of  $g^*$  is the ideal generated by  $4P_2 - P_1^2$  and the kernel of  $h^*$  is the ideal generated by  $Y$ .*

*Proof.* The proof of (i) and the fact that  $g^*(4P_2 - P_1^2) = 0$  consists of identifying the Pontryagin classes with polynomials in the roots of  $SO(7)$ , computing the images of these polynomials under  $g^*$ , and using the fact that there are two generators of  $M^*(BG_2, \mathbf{R}_0)$ , one 4-dimensional, and the other 12-dimensional. We omit the details. From Lemma 3.8, Theorem 2.1 and Theorem 3.2, we have  $h^*(Y) = 0$  and  $h^*(\omega) = 0$ . An easy calculation shows that  $g^*(4P_2 - P_1^2) = 0$ . That  $Y$  and  $4P_2 - P_1^2$  generate the kernels of  $h^*$  and  $g^*$  follows from (i).

**4. Sphere transitive structures on spheres and other homogeneous spaces.** The study of the existence of almost complex structures on spheres is a well-known problem in algebraic topology; it was solved by Borel and Serre [4]. Thus the results of this section can be viewed as a generalization of this problem. Many of the results we present are not new. However, we give them in order that we may write down in an organized fashion the complete solution to the problem of the existence of sphere-transitive structures on spheres.

We shall need two preliminary results.

**LEMMA 4.1.** *Let  $G$  act transitively and linearly on  $S^{2n-1}$  with*

isotropy subgroup  $H$ . Then if the tangent bundle of  $S^{2n}$  can be reduced to  $G$ , the subgroup of elements of  $\pi_{2n-2}(H)$  which are inessential in  $G$  has order at most 2.

*Proof.* Consider the following commutative diagram:

$$\begin{array}{ccccccc}
 H & \xrightarrow{k} & G & \xrightarrow{h} & SO(2n) & \xrightarrow{j} & SO(2n + 1) \\
 & & p_1 \downarrow & \swarrow p_2 & & & \downarrow p_3 \\
 & & S^{2n-1} & & & & S^{2n} .
 \end{array}$$

Here the  $p_i$  are evaluation maps,  $j$  and  $k$  denote the inclusion of the respective isotropy subgroups, and  $h$  denotes the representation of  $G$  arising from the action on  $S^{2n-1}$ . Let  $\partial_1$  be the boundary operator in the homotopy sequence of the fibration  $H \xrightarrow{k} G \xrightarrow{p} S^{2n-1}$  and  $\partial_2$  the boundary operator in the homotopy sequence of fibration  $SO(2n) \rightarrow SO(2n + 1) \rightarrow S^{2n}$ . Let  $\iota_k \in \pi_k(S^k)$  denote the homotopy class of the identity map of  $S^k$ . A reduction of the structure group of the tangent bundle of  $S^{2n}$  to  $G$  is equivalent to the existence of an element  $\alpha \in \pi_{2n-1}(G)$  such that  $h_*(\alpha) = \partial_2(\iota_{2n})$ . Then  $p_{1*}(\alpha) = p_{2*}h_*(\alpha) = p_{2*}\partial_2(\iota_{2n}) = 2\iota_{2n-1}$  and so  $\partial_1(2\iota_{2n-1}) = \partial_1(p_{1*}\alpha) = 0$ . Hence  $\partial_1(\pi_{2n-1}(S^{2n-1})) \subseteq \pi_{2n-2}(H)$  has order at most 2. By the exactness of the homotopy sequence this subgroup is equal to  $\ker(k: \pi_{2n-2}(H) \rightarrow \pi_{2n-2}(G))$ .

**LEMMA 4.2.** *We have  $\pi_{4n-2}(\mathrm{Sp}(n)) = 0$  and  $(2n - 1)!$  divides the order of  $\pi_{4n-2}(\mathrm{Sp}(n - 1))$  for  $n \geq 2$ .*

*Proof.*  $\pi_{4n-2}(\mathrm{Sp}(n))$  is in the stable range and is 0 by Bott periodicity. To prove the other assertion we consider the homomorphism of homotopy sequences of fibrations induced by the commutative diagram

$$\begin{array}{ccccc}
 \mathrm{Sp}(n - 1) & \longrightarrow & \mathrm{Sp}(n) & \longrightarrow & S^{4n-1} \\
 \downarrow \iota & & \downarrow & & \downarrow \text{identity} \\
 U(2n - 1) & \longrightarrow & U(2n) & \longrightarrow & S^{4n-1}
 \end{array}$$

where the horizontal lines are fibrations. Let  $\partial_1$  and  $\partial_2$  be the boundary maps of the homotopy sequences of the upper and lower lines, respectively. Then  $\iota_* \circ \partial_1 = \partial_2$ ; hence the order of  $\mathrm{Im}(\partial_1)$  is a multiple of the order of  $\mathrm{Im}(\partial_2)$ . But  $Z_{(2n-1)!} = \pi_{4n-2}(U(2n - 1)) \subset \mathrm{Im} \partial_2$  see [5]. Hence  $(2n - 1)!$  divides the order of  $\pi_{4n-2}(\mathrm{Sp}(n - 1))$ .

**THEOREM 4.3.** *Let  $\tau(S^n)$  denote the tangent bundle of  $S^n$ . The following is a complete list of sphere transitive structures on*

spheres:

- (i)  $SO(n)$  on  $\tau(S^n)$ ,
- (ii)  $U(3)$  on  $\tau(S^6)$ ,
- (iii)  $SU(3)$  on  $\tau(S^6)$ ,
- (iv)  $G_2$  on  $\tau(S^7)$ .

*Proof.* We have (i) because  $S^n$  is orientable and (iv) because  $S^7$  is parallelizable. (ii) is a consequence of the fact that  $S^6$  has an almost complex structure. Actually, however, it turns out that structure group of the tangent bundle  $\tau(S^6)$  can be reduced to  $SU(3)$  (see [8]) so that (iii) holds.

Next we show that there are no other sphere transitive structures. We do this case by case.

$U(n)$ : Borel and Serre proved that for  $n \neq 1, 3$   $\tau(S^{2n})$  cannot have a  $U(n)$  structure.

$SU(n)$ : Since  $\tau(S^{2n})$  ( $n \neq 1, 3$ ) cannot have a  $U(n)$  structure, it cannot have an  $SU(n)$  structure because  $SU(n) \subseteq U(n)$ .

$Sp(n)$ : Since  $\tau(S^{4n})$  ( $n \neq 1$ ) cannot have a  $U(2n)$  structure and  $Sp(n) \subseteq U(2n)$ ,  $\tau(S^{4n})$  cannot have a  $Sp(n)$  structure.

$Sp(n) \cdot SO(2)$ : We have  $Sp(n) \cdot SO(2) \subseteq U(2n)$ . Thus the argument for  $Sp(n)$  applies in this case also.

$Sp(n) \cdot Sp(1)$ : For  $n \geq 1$ ,  $Sp(n) \cdot Sp(1)$  is covered by

$$Sp(n) \times Sp(1) = Sp(n) \times S^3.$$

We have  $\pi_k(Sp(n) \cdot Sp(1)) \cong \pi_k Sp(n) \oplus \pi_k(S^3)$  for  $k > 1$ . By the second part of Lemma 4.2, it follows that for  $n \geq 2$ ,  $\pi_{4n-2}(Sp(n) \cdot Sp(1)) = \pi_{4n-2}(S^3)$  and  $\pi_{4n-2}(Sp(n-1) \cdot Sp(1))$  is the direct sum of  $\pi_{4n-2}(S^3)$  with a group of order at least  $(2n-1)!$ . Since  $\pi_{4n-2}(S^3)$  is finite, it follows that the necessary condition for a  $Sp(n) \cdot Sp(1)$ -structure on  $S^{4n}$  given by Lemma 4.1 fails, for  $n > 1$ .

$Spin(7)$ : According to Theorem 3.2 (iv) a necessary condition that an 8-dimensional vector bundle  $\xi$  have a transitive  $Spin(7)$  reduction is that  $4P_2(\xi) - P_1^2(\xi) \mp 8X(\xi) = 0$ . The tangent bundle of  $S^8$  (or its negative) does not satisfy this condition.

$Spin(9)$ : Suppose the tangent bundle  $\tau = \tau(S^{16})$  had a transitive  $Spin(9)$  structure. We have  $P_i(\tau) = 0$  ( $i = 1, \dots, 7$ ),  $X(\tau) = 2$ . Hence by Theorem 3.7 (ii),  $Q_i(\tau) = 0$  ( $i = 1, \dots, 7$ ) and  $Z(\tau) = 0$  (at least with rational coefficients). This contradicts the fact that we must have  $X(\tau) = Z(\tau)$ . The same argument shows that  $-\tau$  cannot have a transitive  $Spin(9)$  reduction.

We conclude with some brief remarks about the existence of sphere transitive structures on various simply connected compact homogeneous spaces other than spheres. Denote by  $P^n(C)$  and  $P^n(Q)$

complex and quaternionic projective spaces of real dimension  $2n$  and  $4n$ , respectively. Also let  $\bar{Q}_n$  denote the space of all nonoriented 2-planes in  $\mathbb{R}^{n+2}$ .

**THEOREM 4.4.** *The homogeneous spaces  $S^6 \times S^2$ ,  $S^4 \times S^4$ ,  $S^4 \times S^2 \times S^2$ ,  $(S^2)^4$ ,  $P^4(\mathbb{C})$ ,  $P^3(\mathbb{C}) \times S^2$ ,  $P^2(\mathbb{C}) \times P^2(\mathbb{C})$ ,  $P^2(\mathbb{C}) \times S^4$ ,  $P^2(\mathbb{C}) \times S^2 \times S^2$ ,  $P^1(\mathbb{Q}) \times P^2(\mathbb{C})$ ,  $P^1(\mathbb{Q}) \times S^4$ ,  $P^1(\mathbb{Q}) \times S^2 \times S^2$ ,  $\bar{Q}_2 \times P^1(\mathbb{Q})$ ,  $\bar{Q}_2 \times P^2(\mathbb{C})$ ,  $\bar{Q}_2 \times S^4$ ,  $\bar{Q}_2 \times S^2 \times S^2$  do not possess sphere transitive Spin (7) structures.*

*Proof.* For each case one computes (see [3]) the Pontryagin and Euler classes and verifies that they do not satisfy  $P_2 - 4P_1^2 \pm 8X = 0$ .

In contrast to Theorem 4.4 we have the following result.

**THEOREM 4.5.** *Either orientation of the spaces  $P^2(\mathbb{Q})$ ,  $\bar{Q}_1$ , and  $G_2/SO(4)$  possesses a sphere transitive Spin (7) structure.*

*Proof.* According to [3] each of these spaces has integral cohomology  $\mathbb{Z}[u]/(u^4)$  where  $u$  is a 4-dimensional generator. Furthermore  $P_1 = 2u$ ,  $P_2 = 7u^2$ , and  $X = \pm 3u^2$  for each of these spaces (with the proper choice of  $u$ ). Theorem (4.5) now follows from Theorem 3.4.

It would be interesting to construct explicitly a sphere transitive Spin (7) structure (i.e., a 3-fold vector cross product) on  $P^2(\mathbb{Q})$ .

Finally we have the following theorem.

**THEOREM 4.6.** *Let  $C = F_4/\text{Spin}(9)$  denote the Cayley plane with the canonical orientation. Then  $C$  does not possess a sphere transitive Spin (9) structure, but  $-C$  does.*

*Proof.* We have  $H^*(C, \mathbb{Z}) = \mathbb{Z}[u]/(u^4)$  where  $u$  is an 8-dimensional generator. With the proper choice of  $u$  we have by [3] that for the Cayley plane,  $P_2 = 6u$ ,  $P_4 = 39u^2$ ,  $P_1 = P_3 = 0$ , and  $X = \pm 3u^2$ . It is well known that at least one orientation of  $C$  possesses a sphere transitive Spin (9) structure. It is not hard to verify that  $-C$  satisfies the conclusions of Theorem 3.7 while  $C$  does not. Hence we get Theorem 4.6.

## REFERENCES

1. M. Berger, *Sur les groupes d'holonomie homogène des variétés à connexion affine et des variétés riemanniens*, Bull. Soc. Math. France **83** (1955), 279-330.
2. A. Borel, *Sur l'homologie et la cohomologie des groupes de Lie compacts connexes*, Amer. J. Math, **76** (1954), 273-342.

3. A. Borel and F. Hirzebruch, *Characteristic classes and homogeneous spaces I*, Amer. J. Math. **80** (1958), 458-538.
4. A. Borel and J. Serre, *Groupes de Lie et puissances réduits de Steenrod*, Amer. J. Math. **75** (1953), 409-448.
5. R. Bott, *A report on the unitary group*, Proc. of Symposia Pure Math. vol. 3; *Differential Geometry*, 1-6.
6. R. Brown and A. Gray, *Vector cross products*, Comment. Math. Helv. **42** (1967), 222-236.
7. H. Freudenthal, *Octaven Ausnahmen gruppen, und Oktavengeometrie*, Utrecht, 1951.
8. A. Gray, *Vector cross products on manifolds*, Trans. Amer. Math. Soc. **141** (1969), 465-504.
9. ———, *A note on Riemannian manifolds with holonomy groups  $Sp(n) \cdot Sp(1)$* , Michigan Math J. **16** (1969), 125-128.
10. D. Montgomery and H. Samelson, *Transformation groups of spheres*, Ann. of Math. **44** (1943), 454-470.
11. J. Simons, *On transitivity of holonomy systems*, Ann. of Math. **76** (1962), 213-234.
12. E. Thomas, *On the cohomology groups of the classifying space for the stable spinor group*, Bol. Soc. Math. Mexicana, **7** (1962), 57-69.
13. H. Toda, Y. Saito and I. Yokota, *Note on the generator of  $\pi_7(SO(n))$* , Memoirs of the College of Science, University of Kyoto (A) **30** (1957), 227-230.

Received October 27, 1969. The work of the first author was partially supported by a University of Maryland Faculty Fellowship.

UNIVERSITY OF MARYLAND  
COLLEGE PARK, MARYLAND

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD PIERCE  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER



Johan Aarnes, Edward George Effros and Ole A. Nielsen, <i>Locally compact spaces and two classes of <math>C^*</math>-algebras</i> .....	1
Allan C. Cochran, R. Keown and C. R. Williams, <i>On a class of topological algebras</i> .....	17
John Dauns, <i>Integral domains that are not embeddable in division rings</i> ....	27
Robert Jay Daverman, <i>On the number of nonpiercing points in certain crumpled cubes</i> .....	33
Bryce L. Elkins, <i>Characterization of separable ideals</i> .....	45
Zbigniew Fiedorowicz, <i>A comparison of two naturally arising uniformities on a class of pseudo-PM spaces</i> .....	51
Henry Charles Finlayson, <i>Approximation of Wiener integrals of functionals continuous in the uniform topology</i> .....	61
Theodore William Gamelin, <i>Localization of the corona problem</i> .....	73
Alfred Gray and Paul Stephen Green, <i>Sphere transitive structures and the triality automorphism</i> .....	83
Charles Lemuel Hagopian, <i>On generalized forms of aposynthesis</i> .....	97
J. Jakubík, <i>On subgroups of a pseudo lattice ordered group</i> .....	109
Cornelius W. Onneweer, <i>On uniform convergence for Walsh-Fourier series</i> .....	117
Stanley Joel Osher, <i>On certain Toeplitz operators in two variables</i> .....	123
Washek (Vaclav) Frantisek Pfeffer and John Benson Wilbur, <i>On the measurability of Perron integrable functions</i> .....	131
Frank J. Polansky, <i>On the conformal mapping of variable regions</i> .....	145
Kouei Sekigawa and Shûkichi Tanno, <i>Sufficient conditions for a Riemannian manifold to be locally symmetric</i> .....	157
James Wilson Stepp, <i>Locally compact Clifford semigroups</i> .....	163
Ernest Lester Stitzinger, <i>Frattini subalgebras of a class of solvable Lie algebras</i> .....	177
George Szeto, <i>The group character and split group algebras</i> .....	183
Mark Lawrence Teply, <i>Homological dimension and splitting torsion theories</i> .....	193
David Bertram Wales, <i>Finite linear groups of degree seven. II</i> .....	207
Robert Breckenridge Warfield, Jr., <i>An isomorphic refinement theorem for Abelian groups</i> .....	237
James Edward West, <i>The ambient homeomorphy of an incomplete subspace of infinite-dimensional Hilbert spaces</i> .....	257
Peter Wilker, <i>Adjoint product and hom functors in general topology</i> .....	269
Daniel Eliot Wulbert, <i>A note on the characterization of conditional expectation operators</i> .....	285