

# Lecture Notes in Physics

## Editorial Board

R. Beig, Wien, Austria  
W. Beiglböck, Heidelberg, Germany  
W. Domcke, Garching, Germany  
B.-G. Englert, Singapore  
U. Frisch, Nice, France  
P. Hänggi, Augsburg, Germany  
G. Hasinger, Garching, Germany  
K. Hepp, Zürich, Switzerland  
W. Hillebrandt, Garching, Germany  
D. Imboden, Zürich, Switzerland  
R. L. Jaffe, Cambridge, MA, USA  
R. Lipowsky, Golm, Germany  
H. v. Löhneysen, Karlsruhe, Germany  
I. Ojima, Kyoto, Japan  
D. Sornette, Nice, France, and Zürich, Switzerland  
S. Theisen, Golm, Germany  
W. Weise, Garching, Germany  
J. Wess, München, Germany  
J. Zittartz, Köln, Germany

## The Lecture Notes in Physics

The series Lecture Notes in Physics (LNP), founded in 1969, reports new developments in physics research and teaching – quickly and informally, but with a high quality and the explicit aim to summarize and communicate current knowledge in an accessible way. Books published in this series are conceived as bridging material between advanced graduate textbooks and the forefront of research to serve the following purposes:

- to be a compact and modern up-to-date source of reference on a well-defined topic;
- to serve as an accessible introduction to the field to postgraduate students and nonspecialist researchers from related areas;
- to be a source of advanced teaching material for specialized seminars, courses and schools.

Both monographs and multi-author volumes will be considered for publication. Edited volumes should, however, consist of a very limited number of contributions only. Proceedings will not be considered for LNP.

Volumes published in LNP are disseminated both in print and in electronic formats, the electronic archive is available at [springerlink.com](http://springerlink.com). The series content is indexed, abstracted and referenced by many abstracting and information services, bibliographic networks, subscription agencies, library networks, and consortia.

Proposals should be sent to a member of the Editorial Board, or directly to the managing editor at Springer:

Dr. Christian Caron  
Springer Heidelberg  
Physics Editorial Department I  
Tiergartenstrasse 17  
69121 Heidelberg/Germany  
[christian.caron@springer.com](mailto:christian.caron@springer.com)

Francesco Iachello

# Lie Algebras and Applications

 Springer

Author

Francesco Iachello  
Department of Physics  
Yale University  
P.O. Box 208120  
New Haven, CT 06520-8120  
USA  
E-mail: francesco.iachello@yale.edu

---

F. Iachello, *Lie Algebras and Applications*, Lect. Notes Phys. 708  
(Springer, Berlin Heidelberg 2006), DOI 10.1007/b11785361

---

Library of Congress Control Number: 2006929595

ISSN 0075-8450

ISBN-10 3-540-36236-3 Springer Berlin Heidelberg New York

ISBN-13 978-3-540-36236-4 Springer Berlin Heidelberg New York

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer. Violations are liable for prosecution under the German Copyright Law.

Springer is a part of Springer Science+Business Media  
springer.com  
© Springer-Verlag Berlin Heidelberg 2006

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

Typesetting: by the author and techbooks using a Springer L<sup>A</sup>T<sub>E</sub>X macro package  
Cover design: *design & production* GmbH, Heidelberg

Printed on acid-free paper      SPIN: 11785361      54/techbooks      5 4 3 2 1 0

# Preface

In the second part of the 20th century, algebraic methods have emerged as a powerful tool to study theories of physical phenomena, especially those of quantal systems. The framework of Lie algebras, initially introduced by Sophus Lie in the last part of the 19th century, has been considerably expanded to include graded Lie algebras, infinite-dimensional Lie algebras, and other algebraic constructions. Algebras that were originally introduced to describe certain properties of a physical system, in particular behavior under rotations and translations, have now taken center stage in the construction of physical theories.

This book contains a set of notes from lectures given at Yale University and other universities and laboratories in the last 20 years. The notes are intended to provide an introduction to Lie algebras at the level of a one-semester graduate course in physics. Lie algebras have been particularly useful in spectroscopy, where they were introduced by Eugene Wigner and Giulio Racah. Racah's lectures were given at Princeton University in 1951 (Group Theory and Spectroscopy) and they provided the impetus for the initial applications in atomic and nuclear physics. In the intervening years, many other applications have been made. This book contains a brief account of some of these applications to the fields of molecular, atomic, nuclear, and particle physics. The application of Lie algebraic methods in Physics is so wide that often students are overwhelmed by the sheer amount of material to absorb. This book is intended to give a basic introduction to the method and how it is applied in practice, with emphasis on bosonic systems as encountered in molecules (vibron model), and in nuclei (interacting boson model), and to fermionic systems as encountered in atoms (atomic shell model), and nuclei (nuclear shell model), and hadrons (quark model). Exactly solvable problems in quantum mechanics are also discussed.

Associated with a Lie algebra there is a Lie group. Although the emphasis of these lecture notes is on Lie algebras, a chapter is devoted to Lie groups and to the relation between Lie algebras and Lie groups.

Many exhaustive books exist on the subject of Lie algebras and Lie groups. Reference to these books is made throughout, so that the interested student can study the subject in depth. A selected number of other books, not

explicitly mentioned in the text, are also included in the reference list, to serve as additional introductory material and for cross-reference.

In the early stages of preparing the notes, I benefited from many conversations with Morton Hamermesh, Brian Wybourne, Asim Barut, and Jin-Quan Chen, who wrote books on the subject, but are no longer with us. This book is dedicated to their memory. I also benefited from many conversations with Robert Gilmore, who has written a book on the subject, and with Phil Elliott, Igal Talmi, Akito Arima, Bruno Gruber, Arno Böhm, Yuval Ne'eman, Marcos Moshinsky, and Yuri Smirnov, David Rowe, who have made major contributions to this field.

I am very much indebted to Mark Caprio for a critical reading of the manuscript, and for his invaluable help in preparing the final version of these lecture notes.

New Haven  
May 2006

*Francesco Iachello*

# Contents

<b>1</b>	<b>Basic Concepts</b> .....	1
1.1	Definitions .....	1
1.2	Lie Algebras .....	1
1.3	Change of Basis .....	3
1.4	Complex Extensions .....	4
1.5	Lie Subalgebras .....	4
1.6	Abelian Algebras .....	5
1.7	Direct Sum .....	5
1.8	Ideals (Invariant Subalgebras) .....	6
1.9	Semisimple Algebras .....	7
1.10	Semidirect Sum .....	7
1.11	Killing Form .....	8
1.12	Compact and Non-Compact Algebras .....	9
1.13	Derivations .....	9
1.14	Nilpotent Algebras .....	10
1.15	Invariant Casimir Operators .....	10
1.16	Invariant Operators for Non-Semisimple Algebras .....	12
1.17	Structure of Lie Algebras .....	12
	1.17.1 Algebras with One Element .....	12
	1.17.2 Algebras with Two Elements .....	12
	1.17.3 Algebras with Three Elements .....	13
<b>2</b>	<b>Semisimple Lie Algebras</b> .....	15
2.1	Cartan-Weyl Form of a (Complex) Semisimple Lie Algebra .....	15
2.2	Graphical Representation of Root Vectors .....	15
2.3	Explicit Construction of the Cartan-Weyl Form .....	17
2.4	Dynkin Diagrams .....	19
2.5	Classification of (Complex) Semisimple Lie Algebras .....	21
2.6	Real Forms of Complex Semisimple Lie Algebras .....	21
2.7	Isomorphisms of Complex Semisimple Lie Algebras .....	21
2.8	Isomorphisms of Real Lie Algebras .....	22
2.9	Enveloping Algebra .....	23
2.10	Realizations of Lie Algebras .....	23
2.11	Other Realizations of Lie Algebras .....	24

<b>3</b>	<b>Lie Groups</b> .....	27
3.1	Groups of Transformations .....	27
3.2	Groups of Matrices .....	27
3.3	Properties of Matrices .....	28
3.4	Continuous Matrix Groups .....	29
3.5	Examples of Groups of Transformations .....	32
3.5.1	The Rotation Group in Two Dimensions, $SO(2)$ .....	32
3.5.2	The Lorentz Group in One Plus One Dimension, $SO(1,1)$ .....	33
3.5.3	The Rotation Group in Three Dimensions .....	34
3.5.4	The Special Unitary Group in Two Dimensions, $SU(2)$ .....	34
3.5.5	Relation Between $SO(3)$ and $SU(2)$ .....	35
3.6	Lie Algebras and Lie Groups .....	37
3.6.1	The Exponential Map .....	37
3.6.2	Definition of Exp .....	37
3.6.3	Matrix Exponentials .....	38
<b>4</b>	<b>Irreducible Bases (Representations)</b> .....	39
4.1	Definitions .....	39
4.2	Abstract Characterization .....	39
4.3	Irreducible Tensors .....	40
4.3.1	Irreducible Tensors with Respect to $GL(n)$ .....	40
4.3.2	Irreducible Tensors with Respect to $SU(n)$ .....	41
4.3.3	Irreducible Tensors with Respect to $O(n)$ . Contractions .....	41
4.4	Tensor Representations of Classical Compact Algebras .....	42
4.4.1	Unitary Algebras $u(n)$ .....	42
4.4.2	Special Unitary Algebras $su(n)$ .....	42
4.4.3	Orthogonal Algebras $so(n)$ , $n = \text{Odd}$ .....	43
4.4.4	Orthogonal Algebras $so(n)$ , $n = \text{Even}$ .....	43
4.4.5	Symplectic Algebras $sp(n)$ , $n = \text{Even}$ .....	43
4.5	Spinor Representations .....	44
4.5.1	Orthogonal Algebras $so(n)$ , $n = \text{Odd}$ .....	44
4.5.2	Orthogonal Algebras $so(n)$ , $n = \text{Even}$ .....	44
4.6	Fundamental Representations .....	45
4.6.1	Unitary Algebras .....	45
4.6.2	Special Unitary Algebras .....	45
4.6.3	Orthogonal Algebras, $n = \text{Odd}$ .....	45
4.6.4	Orthogonal Algebras, $n = \text{Even}$ .....	46
4.6.5	Symplectic Algebras .....	46
4.7	Chains of Algebras .....	46
4.8	Canonical Chains .....	46
4.8.1	Unitary Algebras .....	47
4.8.2	Orthogonal Algebras .....	48



4.9	Isomorphisms of Spinor Algebras . . . . .	49
4.10	Nomenclature for $u(n)$ . . . . .	50
4.11	Dimensions of the Representations . . . . .	50
4.11.1	Dimensions of the Representations of $u(n)$ . . . . .	51
4.11.2	Dimensions of the Representations of $su(n)$ . . . . .	52
4.11.3	Dimensions of the Representations of $A_n \equiv su(n+1)$ . . . . .	52
4.11.4	Dimensions of the Representations of $B_n \equiv so(2n+1)$ . . . . .	52
4.11.5	Dimensions of the Representations of $C_n \equiv sp(2n)$ . . . . .	53
4.11.6	Dimensions of the Representations of $D_n \equiv so(2n)$ . . . . .	53
4.12	Action of the Elements of $g$ on the Basis $\mathcal{B}$ . . . . .	53
4.13	Tensor Products . . . . .	56
4.14	Non-Canonical Chains . . . . .	58
<b>5</b>	<b>Casimir Operators and Their Eigenvalues</b> . . . . .	<b>63</b>
5.1	Definitions . . . . .	63
5.2	Independent Casimir Operators . . . . .	63
5.2.1	Casimir Operators of $u(n)$ . . . . .	63
5.2.2	Casimir Operators of $su(n)$ . . . . .	64
5.2.3	Casimir Operators of $so(n)$ , $n = \text{Odd}$ . . . . .	64
5.2.4	Casimir Operators of $so(n)$ , $n = \text{Even}$ . . . . .	64
5.2.5	Casimir Operators of $sp(n)$ , $n = \text{Even}$ . . . . .	65
5.2.6	Casimir Operators of the Exceptional Algebras . . . . .	65
5.3	Complete Set of Commuting Operators . . . . .	65
5.3.1	The Unitary Algebra $u(n)$ . . . . .	66
5.3.2	The Orthogonal Algebra $so(n)$ , $n = \text{Odd}$ . . . . .	66
5.3.3	The Orthogonal Algebra $so(n)$ , $n = \text{Even}$ . . . . .	66
5.4	Eigenvalues of Casimir Operators . . . . .	66
5.4.1	The Algebras $u(n)$ and $su(n)$ . . . . .	67
5.4.2	The Orthogonal Algebra $so(2n+1)$ . . . . .	69
5.4.3	The Symplectic Algebra $sp(2n)$ . . . . .	71
5.4.4	The Orthogonal Algebra $so(2n)$ . . . . .	72
5.5	Eigenvalues of Casimir Operators of Order One and Two . . . . .	74
<b>6</b>	<b>Tensor Operators</b> . . . . .	<b>75</b>
6.1	Definitions . . . . .	75
6.2	Coupling Coefficients . . . . .	76
6.3	Wigner-Eckart Theorem . . . . .	77
6.4	Nested Algebras. Racah's Factorization Lemma . . . . .	79
6.5	Adjoint Operators . . . . .	81
6.6	Recoupling Coefficients . . . . .	83
6.7	Symmetry Properties of Coupling Coefficients . . . . .	84
6.8	How to Compute Coupling Coefficients . . . . .	85
6.9	How to Compute Recoupling Coefficients . . . . .	86
6.10	Properties of Recoupling Coefficients . . . . .	86

6.11	Double Recoupling Coefficients . . . . .	87
6.12	Coupled Tensor Operators . . . . .	88
6.13	Reduction Formula of the First Kind . . . . .	88
6.14	Reduction Formula of the Second Kind . . . . .	89
<b>7</b>	<b>Boson Realizations . . . . .</b>	<b>91</b>
7.1	Boson Operators . . . . .	91
7.2	The Unitary Algebra $u(1)$ . . . . .	92
7.3	The Algebras $u(2)$ and $su(2)$ . . . . .	93
7.3.1	Subalgebra Chains . . . . .	93
7.4	The Algebras $u(n), n \geq 3$ . . . . .	97
7.4.1	Racah Form . . . . .	97
7.4.2	Tensor Coupled Form of the Commutators . . . . .	98
7.4.3	Subalgebra Chains Containing $so(3)$ . . . . .	99
7.5	The Algebras $u(3)$ and $su(3)$ . . . . .	99
7.5.1	Subalgebra Chains . . . . .	100
7.5.2	Lattice of Algebras . . . . .	103
7.5.3	Boson Calculus of $u(3) \supset so(3)$ . . . . .	103
7.5.4	Matrix Elements of Operators in $u(3) \supset so(3)$ . . . . .	105
7.5.5	Tensor Calculus of $u(3) \supset so(3)$ . . . . .	106
7.5.6	Other Boson Constructions of $u(3)$ . . . . .	107
7.6	The Unitary Algebra $u(4)$ . . . . .	108
7.6.1	Subalgebra Chains not Containing $so(3)$ . . . . .	109
7.6.2	Subalgebra Chains Containing $so(3)$ . . . . .	109
7.7	The Unitary Algebra $u(6)$ . . . . .	115
7.7.1	Subalgebra Chains not Containing $so(3)$ . . . . .	115
7.7.2	Subalgebra Chains Containing $so(3)$ . . . . .	115
7.8	The Unitary Algebra $u(7)$ . . . . .	123
7.8.1	Subalgebra Chain Containing $g_2$ . . . . .	124
7.8.2	The Triplet Chains . . . . .	125
<b>8</b>	<b>Fermion Realizations . . . . .</b>	<b>131</b>
8.1	Fermion Operators . . . . .	131
8.2	Lie Algebras Constructed with Fermion Operators . . . . .	131
8.3	Racah Form . . . . .	132
8.4	The Algebras $u(2j + 1)$ . . . . .	133
8.4.1	Subalgebra Chain Containing $spin(3)$ . . . . .	134
8.4.2	The Algebras $u(2)$ and $su(2)$ . Spinors . . . . .	134
8.4.3	The Algebra $u(4)$ . . . . .	136
8.4.4	The Algebra $u(6)$ . . . . .	137
8.5	The Algebra $u(\sum_i (2j_i + 1))$ . . . . .	138
8.6	Internal Degrees of Freedom (Different Spaces) . . . . .	139
8.6.1	The Algebras $u(4)$ and $su(4)$ . . . . .	139
8.6.2	The Algebras $u(6)$ and $su(6)$ . . . . .	141
8.7	Internal Degrees of Freedom (Same Space) . . . . .	142

8.7.1	The Algebra $u((2l + 1)(2s + 1))$ : $L$ - $S$ Coupling . . . . .	142
8.7.2	The Algebra $u\left(\sum_j (2j + 1)\right)$ : $j$ - $j$ Coupling . . . . .	145
8.7.3	The Algebra $u\left(\left(\sum_l (2l + 1)\right)(2s + 1)\right)$ : Mixed $L$ - $S$ Configurations . . . . .	146
<b>9</b>	<b>Differential Realizations</b> . . . . .	147
9.1	Differential Operators . . . . .	147
9.2	Unitary Algebras $u(n)$ . . . . .	147
9.3	Orthogonal Algebras $so(n)$ . . . . .	148
9.3.1	Casimir Operators. Laplace-Beltrami Form . . . . .	150
9.3.2	Basis for the Representations . . . . .	151
9.4	Orthogonal Algebras $so(n, m)$ . . . . .	152
9.5	Symplectic Algebras $sp(2n)$ . . . . .	153
<b>10</b>	<b>Matrix Realizations</b> . . . . .	155
10.1	Matrices . . . . .	155
10.2	Unitary Algebras $u(n)$ . . . . .	155
10.3	Orthogonal Algebras $so(n)$ . . . . .	158
10.4	Symplectic Algebras $sp(2n)$ . . . . .	159
10.5	Basis for the Representation . . . . .	160
10.6	Casimir Operators . . . . .	161
<b>11</b>	<b>Spectrum Generating Algebras and Dynamic Symmetries</b> 163	
11.1	Spectrum Generating Algebras (SGA) . . . . .	163
11.2	Dynamic Symmetries (DS) . . . . .	163
11.3	Bosonic Systems . . . . .	164
11.3.1	Dynamic Symmetries of $u(4)$ . . . . .	165
11.3.2	Dynamic Symmetries of $u(6)$ . . . . .	167
11.4	Fermionic Systems . . . . .	170
11.4.1	Dynamic Symmetry of $u(4)$ . . . . .	170
11.4.2	Dynamic Symmetry of $u(6)$ . . . . .	171
<b>12</b>	<b>Degeneracy Algebras and Dynamical Algebras</b> . . . . .	173
12.1	Degeneracy Algebras . . . . .	173
12.2	Degeneracy Algebras in $\nu \geq 2$ Dimensions . . . . .	173
12.2.1	The Isotropic Harmonic Oscillator . . . . .	174
12.2.2	The Coulomb Problem . . . . .	177
12.3	Degeneracy Algebra in $\nu = 1$ Dimension . . . . .	181
12.4	Dynamical Algebras . . . . .	182
12.5	Dynamical Algebras in $\nu \geq 2$ Dimensions . . . . .	182
12.5.1	Harmonic Oscillator . . . . .	182
12.5.2	Coulomb Problem . . . . .	182
12.6	Dynamical Algebra in $\nu = 1$ Dimension . . . . .	183
12.6.1	Pöschl-Teller Potential . . . . .	183

XII Contents

12.6.2 Morse Potential . . . . .	185
12.6.3 Lattice of Algebras . . . . .	187
<b>References</b> . . . . .	189
<b>Index</b> . . . . .	193

# List of Symbols

$\mathcal{A}$	Abelian algebra
$[, ]_+$	anticommutator
$\{, \}$	anticommutator
$\mathcal{B}$	basis (bosons)
$\mathcal{F}$	basis (fermions)
$\in$	belongs to
$b$	boson annihilation operator
$b^\dagger$	boson creation operator
$\langle  $	bra
$C_i(g)$	Casimir operator of $g$
$[, ]$	commutator
$\mathbb{C}$	complex
$\supset$	contains
$\subset$	contained in
$\equiv$	defined as
Der	derivation
dim	dimension of the representation
$\oplus$	direct sum
$DS$	dynamic symmetry
$X_\alpha$	element of an algebra
$\square$	entry in Young tableau
$\doteq$	equal to
$\langle \rangle$	expectation value
$\downarrow \exp$	exponential map
$a$	fermion annihilation operator
$a^\dagger$	fermion creation operator
$\setminus$	Gel'fand pattern
$\approx$	homomorphic groups
$\cap$	intersect
$IRB$	irreducible basis
$\sim$	isomorphic algebras
$\langle   \rangle$	isoscalar factors
$  \rangle$	ket
$\downarrow$	labels of the representation

XIV List of Symbols

$\nabla^2$	Laplace operator
$\mathcal{G}$	Lie algebra
$g$	Lie algebra
$G$	Lie group
$\mathcal{V}$	linear vector space
$\circ$	long root
$\mathcal{F}$	number field
$O$	octonion
$\alpha^{(+)}$	positive root
$Q$	quaternion
$\langle     \rangle$	reduced matrix elements
$\cdot$	scalar product
$\oplus_s$	semidirect sum
$\bullet$	short root
$SGA$	spectrum generating algebra
$\otimes$	tensor product
$\oplus$	tensor sum
$ 0\rangle$	vacuum
$\mathcal{L}$	vector space

# 1 Basic Concepts

## 1.1 Definitions

The key notion in the definition of Lie algebras is that of the commutator (or bracket), denoted by  $[\cdot, \cdot]$ . The commutator of  $X$  and  $Y$  is defined as

$$[X, Y] = XY - YX . \quad (1.1)$$

It satisfies the relations

$$[X, X] = 0; \quad [X, Y] = -[Y, X] . \quad (1.2)$$

Another key notion is that of number field,  $\mathcal{F}$ . The number fields of interest are: Real,  $R$ , Complex,  $C$ , Quaternion,  $Q$ , and Octonion,  $O$ . Since these lecture notes are intended primarily for applications to quantal systems, where the basic commutation relations between coordinates and momenta are

$$\left[ x, \frac{1}{i} \frac{\partial}{\partial x} \right] = i , \quad (1.3)$$

only real and complex fields will be considered. Although formulations of quantum mechanics in terms of quaternions and octonions have been suggested, Lie algebras over the quaternion and octonion number fields will not be discussed here.

## 1.2 Lie Algebras

Lie algebras are named after the Norwegian mathematician Sophus Lie (1842–1899). Most of what we know about the original formulation comes from Lie's lecture notes in Leipzig, as collected by Scheffers. [S. Lie and G. Scheffers, *Vorlesungen über Kontinuierliche Gruppen*, Leipzig, 1893].

A set of elements  $X_\alpha (\alpha = 1, \dots, r)$  is said to form a Lie algebra  $\mathcal{G}$ , written as  $X_\alpha \in \mathcal{G}$ , if the following axioms are satisfied:

**Axiom 1** *The commutator of any two elements is a linear combination of the elements in the Lie algebra*

$$[X_\rho, X_\sigma] = \sum_\tau c_{\rho\sigma}^\tau X_\tau. \quad (1.4)$$

**Axiom 2.** *The double commutators of three elements satisfy the Jacobi identity*

$$[X_\rho, [X_\sigma, X_\tau]] + [X_\sigma, [X_\tau, X_\rho]] + [X_\tau, [X_\rho, X_\sigma]] = 0. \quad (1.5)$$

The coefficients  $c_{\rho\sigma}^\tau$  are called Lie structure constants. They define the Lie algebra. They satisfy

$$c_{\rho\sigma}^\tau = -c_{\sigma\rho}^\tau \quad (1.6)$$

and

$$c_{\rho\sigma}^\mu c_{\mu\tau}^\nu + c_{\sigma\tau}^\mu c_{\mu\rho}^\nu + c_{\tau\rho}^\mu c_{\mu\sigma}^\nu = 0. \quad (1.7)$$

A tensor notation with covariant,  $X_{\rho\dots}$ , and contravariant,  $X^{\rho\dots}$ , indices has been used in (1.4)–(1.7) and will be used in the remaining part of this chapter. In this notation, the structure constants  $c_{\rho\sigma}^\tau$  are rank-3 tensors with one contravariant and two covariant indices. The outer product of two tensors, for example two covariant vectors,  $X_\rho$  and  $Y_\sigma$ , is written  $X_\rho Y_\sigma$ . The inner product of two tensors, for example two vectors, is written  $X_\rho Y^\rho$ . Also, unless otherwise specified, a summation convention over repeated indices will be used

$$c_{\rho\sigma}^\tau X_\tau \equiv \sum_\tau c_{\rho\sigma}^\tau X_\tau. \quad (1.8)$$

Algebras are usually denoted by script (calligraphic) letters,  $\mathcal{G}$ , or by lowercase letters,  $g$ . The associated Lie groups, to be discussed in Chap. 3, are usually denoted by capital letters,  $G$ . However, often no distinction is made between groups and algebras and the same letter is used for both. Unless otherwise specified, the notation lowercase-capital will be used in these notes.

The  $r$  elements,  $X_\alpha$ , span a  $r$ -dimensional linear vector space  $\mathcal{L}$ . The definition given above in an ‘operational’ definition. A formal definition is:

**Definition 1.** *A vector space  $\mathcal{L}$  over a number field  $\mathcal{F}$ , with an operation  $\mathcal{L} \times \mathcal{L} \rightarrow \mathcal{L}$ , denoted  $[X, Y]$  and called the commutator of  $X$  and  $Y$ , is called a Lie algebra over  $\mathcal{F}$  if the following axioms are satisfied:*

- (1) *The operation is bilinear.*
- (2)  *$[X, X] = 0$  for all  $X$  in  $\mathcal{L}$ .*
- (3)  *$[X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = 0$  ( $X, Y, Z \in \mathcal{L}$ ).*

The properties of bilinearity, i.e.,  $[aX + bY, Z] = a[X, Z] + b[Y, Z]$  and  $[X, bY + cZ] = b[X, Y] + c[X, Z]$ , and of vanishing  $[X, X] = 0$  together guarantee the antisymmetry property (1.2) of the commutator. The closure of  $\mathcal{L}$  under the commutator ( $\mathcal{L} \times \mathcal{L} \rightarrow \mathcal{L}$ ) gives Axiom 1 above for the generators. Property (3) gives Axiom 2 above. [J.E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Graduate Texts in Mathematics, Springer-Verlag, 1972.]



A Lie algebra is called “real” if the field  $\mathcal{F}$  is  $R$ , it is called “complex” if  $\mathcal{F}$  is  $C$ . Real Lie algebras have real structure constants, while complex Lie algebras have structure constants which can be real or complex.

**Example 1.** *The algebra*

$$[X_1, X_2] = X_3, \quad [X_2, X_3] = X_1, \quad [X_3, X_1] = X_2 \quad (1.9)$$

is a real Lie algebra with three elements ( $r = 3$ ). This is the angular momentum algebra in three dimensions,  $so(3)$ .

**Example 2.** *The algebra*

$$[X_1, X_2] = X_3, \quad [X_2, X_3] = -X_1, \quad [X_3, X_1] = X_2 \quad (1.10)$$

is also a real Lie algebra with three elements ( $r = 3$ ). This is the Lorentz algebra in  $2 + 1$  dimensions,  $so(2, 1)$ .

Note the difference between the two (a sign in the commutation relations).

### 1.3 Change of Basis

It is possible to change the basis

$$X'_\sigma = a^\rho_\sigma X_\rho \quad (1.11)$$

where  $a^\rho_\sigma$  is non-singular. The new commutation relations of the algebra are

$$[X'_\rho, X'_\sigma] = c'^\tau_{\rho\sigma} X'_\tau. \quad (1.12)$$

From (1.4) and (1.11), the new structure constants  $c'^\tau_{\rho\sigma}$  are obtained from the old structure constants  $c^\kappa_{\nu\lambda}$  by solving the set of equations

$$c'^\tau_{\rho\sigma} a^\kappa_\tau = a^\nu_\rho a^\lambda_\sigma c^\kappa_{\nu\lambda}. \quad (1.13)$$

Particularly simple is the change of basis in which each element is multiplied by a real number (sometimes called a normalization transformation).

**Example 3.** *The transformation*

$$X'_1 = \sqrt{2}X_1, \quad X'_2 = \sqrt{2}X_2, \quad X'_3 = X_3 \quad (1.14)$$

changes the commutation relations of the Lie algebra  $so(3)$  into

$$[X'_1, X'_2] = 2X'_3; \quad [X'_2, X'_3] = X'_1; \quad [X'_3, X'_1] = X'_2. \quad (1.15)$$

Lie algebras that have the same commutation relations up to a change of basis are called isomorphic. Isomorphism of algebras will be denoted by the symbol  $\sim$ . Isomorphisms of Lie algebras will be discussed in Chap. 2. An example is:

**Example 4.** *The Lie algebras  $so(3)$  and  $su(2)$  are isomorphic*

$$so(3) \sim su(2). \quad (1.16)$$

## 1.4 Complex Extensions

The change of basis (1.11) can be complex. An example is multiplication by the imaginary unit on  $so(3)$ ,

$$J_1 = iX_1, \quad J_2 = iX_2, \quad J_3 = iX_3. \quad (1.17)$$

The commutation relations are now

$$[J_1, J_2] = iJ_3, \quad [J_2, J_3] = iJ_1, \quad [J_3, J_1] = iJ_2. \quad (1.18)$$

The algebra composed of elements  $J_1, J_2, J_3$  is the 'angular momentum algebra' often quoted in quantum mechanics books.

If one takes linear combinations of elements,  $A, B$ , of a real Lie algebra  $g$ , with complex coefficients, and defines  $[A + iB, C] = [A, C] + i[B, C]$  one obtains the complex extension of the real Lie algebra. Starting from a real algebra  $g$ , by making a complex change of basis, one can construct a complex extension of  $g$ . In some cases, the complex extension of different real Lie algebras is the same.

**Example 5.** *The real Lie algebras  $so(2, 1)$  and  $so(3)$  have the same complex extension.*

Consider the real Lie algebra  $so(2, 1)$  of Example 2. By making the change of basis

$$Y_1 = X_1, \quad Y_2 = -iX_2, \quad Y_3 = -iX_3, \quad (1.19)$$

one obtains

$$[Y_1, Y_2] = Y_3; \quad [Y_2, Y_3] = Y_1; \quad [Y_3, Y_1] = Y_2. \quad (1.20)$$

These are the same commutation relations of the real Lie algebra  $so(3)$  of Example 1.

## 1.5 Lie Subalgebras

A subset of elements,  $Y_\beta$ , closed with respect to commutation is called a subalgebra

$$X_\alpha \in g; \quad Y_\beta \in g'; \quad g \supset g' \quad (1.21)$$

The symbol  $\supset$  is used to indicate that  $g'$  is a subalgebra of  $g$ . The subset satisfies the commutation relations

$$[Y_\rho, Y_\sigma] = c_{\rho\sigma}^\tau Y_\tau \quad (1.22)$$

**Example 6.** *The single element  $X_3$  forms a Lie subalgebra of  $so(3) \ni X_1, X_2, X_3$ , since*

$$[X_3, X_3] = 0. \quad (1.23)$$

This is  $so(2)$ , the angular momentum algebra in two dimensions,  $so(3) \ni so(2)$ .

## 1.6 Abelian Algebras

These are a special type of algebras, named after the Norwegian mathematician Niels Abel (1802–1829). An Abelian algebra,  $\mathcal{A}$ , is an algebra for which all elements commute,

$$[X_\rho, X_\sigma] = 0 \quad \text{for any} \quad X_\rho \in \mathcal{A}, \quad X_\sigma \in \mathcal{A}. \quad (1.24)$$

**Example 7.** *The algebra  $so(2) \ni X_3$  is Abelian, since*

$$[X_3, X_3] = 0. \quad (1.25)$$

Any algebra contains a trivial Abelian subalgebra, composed of a single element  $X_\rho$ , since  $[X_\rho, X_\rho] = 0$ . Another non-trivial example is the translation algebra in two dimensions,  $t(2) \ni X_1, X_2$ .

**Example 8.** *The algebra  $t(2)$  with commutation relations*

$$[X_1, X_2] = 0, \quad [X_1, X_1] = 0, \quad [X_2, X_2] = 0, \quad (1.26)$$

*is Abelian.*

## 1.7 Direct Sum

Consider two commuting algebras  $g_1 \ni X_\alpha, g_2 \ni Y_\beta$ , satisfying

$$\begin{aligned} [X_\rho, X_\sigma] &= c_{\rho\sigma}^\tau X_\tau, \\ [Y_\rho, Y_\sigma] &= c'_{\rho\sigma}{}^\tau Y_\tau, \\ [X_\rho, Y_\sigma] &= 0. \end{aligned} \quad (1.27)$$

The commuting property is denoted by  $g_1 \cap g_2 = 0$ . The set of elements  $X_\alpha, Y_\beta$  forms an algebra  $g$ , called the direct sum,

$$g = g_1 \oplus g_2. \quad (1.28)$$

Sometimes, it is possible to rewrite a Lie algebra as a direct sum of other Lie algebras. Consider, the algebra  $so(4) \ni X_1, X_2, X_3, Y_1, Y_2, Y_3$ , satisfying commutation relations

$$\begin{aligned} [X_1, X_2] &= X_3 \quad ; \quad [X_2, X_3] = X_1 \quad ; \quad [X_3, X_1] = X_2 \\ [Y_1, Y_2] &= X_3 \quad ; \quad [Y_2, Y_3] = X_1 \quad ; \quad [Y_3, Y_1] = X_2 \\ [X_1, Y_1] &= 0 \quad ; \quad [X_2, Y_2] = 0 \quad ; \quad [X_3, Y_3] = 0 \\ [X_1, Y_2] &= Y_3 \quad ; \quad [X_1, Y_3] = -Y_2 \\ [X_2, Y_1] &= -Y_3 \quad ; \quad [X_2, Y_3] = Y_1 \\ [X_3, Y_1] &= Y_2 \quad ; \quad [X_3, Y_2] = -Y_1 \end{aligned} \quad (1.29)$$

By a change of basis

$$J_i = \frac{X_i + Y_i}{2} \quad , \quad K_i = \frac{X_i - Y_i}{2} \quad (i = 1, 2, 3) \quad (1.30)$$

the algebra can be brought to the form

$$\begin{aligned} [J_1, J_2] &= J_3 & [J_2, J_3] &= J_1 & [J_3, J_1] &= J_2 \\ [K_1, K_2] &= K_3 & [K_2, K_3] &= K_1 & [K_3, K_1] &= K_2 \\ [J_i, K_j] &= 0 & (i, j &= 1, 2, 3) . \end{aligned} \quad (1.31)$$

In the new form, one can see that the algebra  $so(4)$  is the direct sum of two  $so(3)$  algebras.

**Example 9.** *The algebra  $so(4)$  is isomorphic to the direct sums*

$$so(4) \sim so(3) \oplus so(3) \sim su(2) \oplus su(2) \sim sp(2) \oplus sp(2) . \quad (1.32)$$

The splitting is rarely possible. Consider for example, the algebra  $so(3, 1) \ni X_1, X_2, X_3, Y_1, Y_2, Y_3$ , satisfying commutation relations

$$\begin{aligned} [X_1, X_2] &= X_3 \quad ; \quad [X_2, X_3] = X_1 \quad ; \quad [X_3, X_1] = X_2 \\ [Y_1, Y_2] &= -X_3 \quad ; \quad [Y_2, Y_3] = -X_1 \quad ; \quad [Y_3, Y_1] = -X_2 \\ [X_1, Y_1] &= 0 \quad ; \quad [X_2, Y_2] = 0 \quad ; \quad [X_3, Y_3] = 0 \\ [X_1, Y_2] &= Y_3 \quad ; \quad [X_1, Y_3] = -Y_2 \\ [X_2, Y_1] &= -Y_3 \quad ; \quad [X_2, Y_3] = Y_1 \\ [X_3, Y_1] &= Y_2 \quad ; \quad [X_3, Y_2] = -Y_1 . \end{aligned} \quad (1.33)$$

This algebra cannot be split into a direct sum of real Lie algebras. However, sometimes, the splitting is possible by going to the complex extension of the algebra. For example by taking the combination

$$J_j = \frac{X_j + iY_j}{2} \quad , \quad K_j = \frac{X_j - iY_j}{2} \quad (j = 1, 2, 3) \quad (1.34)$$

one can show that the elements  $J_j, K_j$  satisfy (1.31). The algebras  $so(4)$  and  $so(3, 1)$  have the same complex extension, and can be split in the same fashion.

## 1.8 Ideals (Invariant Subalgebras)

Consider an algebra  $g$  and its subalgebra  $g'$ ,  $X_\alpha \in g$ ,  $Y_\beta \in g'$ ,  $g \supset g'$ . Since  $g'$  is a subalgebra, it satisfies

$$[Y_\rho, Y_\sigma] = c'_{\rho\sigma} Y_\tau . \quad (1.35)$$

If, in addition,

$$[Y_\rho, X_\sigma] = c_{\rho\sigma}^\tau Y_\tau, \tag{1.36}$$

then  $g'$  is called an invariant subalgebra. As an example, consider the Euclidean algebra  $e(2)$ , composed of three elements,  $X_1, X_2, X_3$ , satisfying

$$[X_1, X_2] = X_3 \quad [X_1, X_3] = -X_2 \quad [X_2, X_3] = 0. \tag{1.37}$$

**Example 10.** *The subalgebra  $g' \ni X_2, X_3$  is an (Abelian) ideal of  $g \equiv e(2) \ni X_1, X_2, X_3$ .*

### 1.9 Semisimple Algebras

An algebra which has no Abelian ideals is called semisimple.

**Example 11.** *The algebra  $so(3)$*

$$[X_1, X_2] = X_3 \quad [X_2, X_3] = X_1 \quad [X_3, X_1] = X_2 \tag{1.38}$$

*is semisimple.*

**Example 12.** *The algebra  $e(2)$*

$$[X_1, X_2] = X_3 \quad [X_1, X_3] = -X_2 \quad [X_2, X_3] = 0 \tag{1.39}$$

*is non-semisimple.*

### 1.10 Semidirect Sum

Consider two non-commuting algebras,  $X_\alpha \in g_1, Y_\beta \in g_2$ , satisfying

$$[X_\rho, X_\sigma] = c_{\rho\sigma}^\tau X_\tau, \quad [Y_\rho, Y_\sigma] = c_{\rho\sigma}'^\tau Y_\tau. \tag{1.40}$$

If  $g_2$  is an invariant subalgebra of  $g_1$ ,

$$[X_\rho, Y_\sigma] = c_{\rho\sigma}''^\tau Y_\tau, \tag{1.41}$$

then the algebra  $g$

$$g = g_1 \oplus_s g_2 \tag{1.42}$$

is called the semidirect sum of  $g_1$  and  $g_2$ .

**Example 13.** *The Euclidean algebra  $e(2)$ , composed of three elements,  $X_1, X_2, X_3$ , is the semidirect sum of the rotation algebra in two dimensions,  $so(2)$ , composed of the single element,  $X_1$ , and the translation algebra in two dimensions,  $t(2)$ , composed of two commuting elements,  $X_2, X_3$*

$$e(2) = so(2) \oplus_s t(2). \tag{1.43}$$

## 1.11 Killing Form

With the Lie structure constants one can form a tensor, called metric tensor, or Killing form

$$g_{\sigma\lambda} = g_{\lambda\sigma} = c_{\sigma\rho}^{\tau} c_{\lambda\tau}^{\rho} \quad (1.44)$$

named after Killing, who in a series of papers in the 1880's discussed its properties. [W. Killing, *Die Zusammensetzung der Stetigen Endlichen Transformations-gruppen, I-IV*, Math. Ann. 31, 252 (1888); 33, 1 (1889); 34, 57 (1889); 36, 161 (1890).] The Killing form was used by Cartan to identify semisimple Lie algebras. Cartan's criterion for deciding if a Lie algebra is semisimple is:

**Theorem 1.** *A Lie algebra  $g$  is semisimple if, and only if,*

$$\det |g_{\sigma\lambda}| \neq 0. \quad (1.45)$$

In other words, an inverse  $g^{\sigma\lambda}$  of the metric tensor  $g_{\sigma\lambda}$  exists

$$g^{\sigma\tau} g_{\tau\lambda} = \delta_{\lambda}^{\sigma}, \quad (1.46)$$

where

$$\delta_{\lambda}^{\sigma} = \begin{cases} 1 & \text{if } \sigma = \lambda \\ 0 & \text{if } \sigma \neq \lambda \end{cases}. \quad (1.47)$$

**Example 14.** *The algebra  $so(3)$  is semisimple.*

The metric tensor of  $so(3)$  is

$$g_{\sigma\lambda} = \begin{pmatrix} -2 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & -2 \end{pmatrix}, \quad (1.48)$$

also written in compact form  $g_{\sigma\lambda} = -2\delta_{\sigma\lambda}$ . The determinant of the metric tensor is

$$\det |g_{\sigma\lambda}| = -8 \quad (1.49)$$

and thus the algebra  $so(3)$  is semisimple.

**Example 15.** *The algebra  $so(2,1)$  is semisimple.*

The metric tensor of  $so(2,1)$  is

$$g_{\sigma\lambda} = \begin{pmatrix} -2 & 0 & 0 \\ 0 & +2 & 0 \\ 0 & 0 & +2 \end{pmatrix}. \quad (1.50)$$

The determinant of the metric tensor is again

$$\det |g_{\sigma\lambda}| = -8 \quad (1.51)$$

and thus the algebra  $so(2,1)$  is semisimple.

**Example 16.** *The algebra  $e(2)$  is non-semisimple.*

Finally, consider the algebra  $e(2)$  with metric tensor

$$g_{\sigma\lambda} = \begin{pmatrix} -2 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \tag{1.52}$$

In this case

$$\det |g_{\sigma\lambda}| = 0, \tag{1.53}$$

and thus the algebra  $e(2)$  is non-semisimple.

### 1.12 Compact and Non-Compact Algebras

A real semisimple Lie algebra is compact if its metric tensor is negative definite.

**Example 17.** *The algebra  $so(3)$  is compact.*

The metric tensor of  $so(3)$  is negative definite. In its diagonal form all elements are negative.

**Example 18.** *The algebra  $so(2, 1)$  is non-compact.*

The metric tensor of  $so(2, 1)$  is non-negative definite. In its diagonal form some elements are positive.

### 1.13 Derivations

Starting with a Lie algebra,  $g$ , with elements  $X_\rho$ , it is possible to construct other algebras, called derivations and denoted by  $\text{Der } g$ , by taking commutators

$$\begin{aligned} g^{(1)} &= [g, g] \\ g^{(2)} &= [g^{(1)}, g^{(1)}] \\ &\dots \end{aligned} \tag{1.54}$$

For example, starting with the Euclidean algebra,  $e(2)$ , with elements

$$g \doteq X_1, X_2, X_3 \tag{1.55}$$

satisfying the commutation relations (1.39), one has

$$\begin{aligned} \text{Der } g &\doteq X_2, X_3 \\ \text{Der}^2 g &\doteq 0. \end{aligned} \tag{1.56}$$

If, for some positive  $k$ ,

$$\text{Der}^k g \doteq 0 \quad (g^{(k)} = 0) \tag{1.57}$$

the algebra is called solvable.

**Example 19.** *The algebra  $e(2)$  is solvable.*

From (1.55),

$$\text{Der}^2 e(2) \doteq 0. \quad (1.58)$$

## 1.14 Nilpotent Algebras

Starting with a Lie algebra,  $g$ , with elements,  $X_\rho$ , it is possible to construct powers of  $g$  as

$$\begin{aligned} g^2 &= g^{(1)} = [g, g] \\ g^3 &= [g, g^2] \\ &\dots \end{aligned} \quad (1.59)$$

If, for some positive  $k$ ,

$$g^k = 0 \quad (1.60)$$

the algebra is called nilpotent.

**Example 20.** *The algebra  $e(2)$  is not nilpotent*

Starting with

$$g \doteq X_1, X_2, X_3, \quad (1.61)$$

satisfying (1.39), one has

$$\begin{aligned} g^2 &\doteq X_2, X_3 \\ g^3 &\doteq X_2, X_3 \\ &\dots \end{aligned} \quad (1.62)$$

## 1.15 Invariant Casimir Operators

These operators play a central role in applications. They are named after the Dutch physicist Casimir, who introduced them in 1931 for the angular momentum algebra  $so(3)$ . [H. Casimir, *Über die Konstruktion einer zu den irreduzibelen Darstellung halbeinfacher kontinuierlichen Gruppen gehörigen Differential-gleichung*, Proc. Roy. Akad. Amst. 34, 844 (1931).] An operator,  $C$ , that commutes with all the elements of a Lie algebra  $g$

$$[C, X_\tau] = 0 \quad \text{for all } X_\tau \in g \quad (1.63)$$

is called an invariant Casimir operator. Casimir operators can be linear, quadratic, cubic,  $\dots$  in the elements  $X_\tau$ . A Casimir operator is called of order  $p$  if it contains products of  $p$  elements  $X_\tau$ ,



$$C_p = \sum_{\alpha_1, \dots, \alpha_p} f^{\alpha_1 \alpha_2 \dots \alpha_p} X_{\alpha_1} X_{\alpha_2} \dots X_{\alpha_p} . \quad (1.64)$$

The summation is explicitly displayed in this formula. Also, if  $C$  commutes with  $g$ , so does  $aC$ ,  $C^2$ ,  $\dots$ . The number of independent Casimir operators of a Lie algebra will be discussed in Chap. 5. The quadratic ( $p = 2$ ) Casimir operator of a semisimple algebra can be simply constructed from the metric tensor

$$C_2 = g^{\rho\sigma} X_\rho X_\sigma = g_{\rho\sigma} X^\rho X^\sigma \equiv C . \quad (1.65)$$

*Proof.* Evaluate the commutator of  $C$  and  $X_\tau$

$$\begin{aligned} [C, X_\tau] &= g^{\rho\sigma} [X_\rho X_\sigma, X_\tau] = g^{\rho\sigma} X_\rho [X_\sigma, X_\tau] + g^{\rho\sigma} [X_\rho, X_\tau] X_\sigma \\ &= g^{\rho\sigma} X_\rho c_{\sigma\tau}^\lambda X_\lambda + g^{\rho\sigma} c_{\rho\tau}^\lambda X_\lambda X_\sigma \\ &= g^{\rho\sigma} c_{\sigma\tau}^\lambda X_\rho X_\lambda + g^{\sigma\rho} c_{\sigma\tau}^\lambda X_\lambda X_\rho = g^{\rho\sigma} c_{\sigma\tau}^\lambda (X_\rho X_\lambda + X_\lambda X_\rho) \\ &= g^{\rho\sigma} g^{\lambda\nu} c_{\nu\sigma\tau} (X_\rho X_\lambda + X_\lambda X_\rho) = 0 \end{aligned} \quad (1.66)$$

The last line follows from the fact that the product  $g^{\sigma\lambda} g^{\lambda\nu}$  is symmetric under  $\lambda \rightarrow \sigma, \nu \rightarrow \rho$ , the product  $(X_\rho X_\sigma + X_\sigma X_\rho)$  is symmetric under  $\lambda \rightarrow \rho$ , and the structure constant  $c_{\nu\rho\sigma}$  is antisymmetric under  $\nu \rightarrow \sigma$ . For a semisimple Lie algebra, indices can be raised and lowered using the metric tensor

$$c_{\sigma\tau}^\lambda = g^{\lambda\nu} c_{\nu\sigma\tau} . \quad (1.67)$$

Higher order Casimir operators can be constructed in a similar fashion

$$C_p = c_{\alpha_1\beta_1}^{\beta_2} c_{\alpha_2\beta_2}^{\beta_3} \dots c_{\alpha_p\beta_p}^{\beta_1} X^{\alpha_1} X^{\alpha_2} \dots X^{\alpha_p} . \quad (1.68)$$

For the algebra  $so(3)$ , the inverse of the metric tensor is

$$g^{\sigma\lambda} = \begin{pmatrix} -\frac{1}{2} & 0 & 0 \\ 0 & -\frac{1}{2} & 0 \\ 0 & 0 & -\frac{1}{2} \end{pmatrix} \quad (1.69)$$

giving

$$C = -\frac{1}{2} (X_1^2 + X_2^2 + X_3^2) . \quad (1.70)$$

For the algebra  $so(2,1)$

$$C = -\frac{1}{2} (X_1^2 - X_2^2 - X_3^2) . \quad (1.71)$$

Note the minus signs. By multiplying  $C$  by 2 and the elements by  $i$ , one obtains for  $so(3)$

$$C' = 2C = J_1^2 + J_2^2 + J_3^2 . \quad (1.72)$$

This is the usual form in which the Casimir operator of the angular momentum algebra  $so(3)$  appears in quantum mechanics textbooks.

## 1.16 Invariant Operators for Non-Semisimple Algebras

For non-semisimple Lie algebras, Casimir operators cannot be simply constructed. One introduces a related notion of invariant operators that commute with all elements.

**Example 21.** *The invariant operator of the Euclidean algebra  $e(2) \ni X_1, X_2, X_3$  is*

$$C = X_2^2 + X_3^2 \quad (1.73)$$

*Proof.* The commutators are

$$\begin{aligned} [X_2^2, X_1] &= X_2[X_2, X_1] + [X_2, X_1] X_2 = -2 X_2 X_3 \\ [X_3^2, X_1] &= X_3[X_3, X_1] + [X_3, X_1] X_3 = 2 X_2 X_3 . \end{aligned} \quad (1.74)$$

Hence

$$[C, X_\tau] = 0 \text{ for any } X_\tau \in g. \quad (1.75)$$

## 1.17 Structure of Lie Algebras

The structure of Lie algebras, semisimple or not, can be investigated by inspection. A detailed account is given by Kirillov. [A.A. Kirillov, *Elements of the Theory of Representation*, Nauka, Moscow, 1978.]

### 1.17.1 Algebras with One Element

We begin with the case  $r = 1$ . In this case there is only one element,  $X$ , and one possibility

$$(a) \quad [X, X] = 0 . \quad (1.76)$$

The algebra is Abelian.

**Example 22.** *The algebras  $so(2) \sim u(1)$  are examples of Kirillov's case 1a.*

### 1.17.2 Algebras with Two Elements

Next consider the case  $r = 2$ . In this case, there are two elements,  $X_1, X_2$ , and two possibilities

$$(a) \quad [X_1, X_2] = 0 , \quad (1.77)$$

and

$$(b) \quad [X_1, X_2] = X_1 . \quad (1.78)$$

In case (a), the algebra is Abelian. In case (b),  $X_1$  is an Abelian ideal.

**Example 23.** *The translation algebra  $t(2)$  is an example of Kirillov's case 2a.*

### 1.17.3 Algebras with Three Elements

For  $r = 3$ , there are three elements,  $X_1, X_2, X_3$  and four possibilities:

$$(a) \quad [X_1, X_2] = [X_2, X_3] = [X_3, X_1] = 0 \quad (1.79)$$

$$(b) \quad [X_1, X_2] = X_3; \quad [X_1, X_3] = [X_2, X_3] = 0 \quad \text{or} \\ [X_1, X_3] = X_2; \quad [X_1, X_2] = [X_2, X_3] = 0, \quad (1.80)$$

$$(c) \quad [X_1, X_2] = 0; \quad [X_3, X_1] = \alpha X_1 + \beta X_2; \quad [X_3, X_2] = \gamma X_1 + \delta X_2, \quad (1.81)$$

where the matrix  $\begin{vmatrix} \alpha & \beta \\ \gamma & \delta \end{vmatrix}$  is non-singular, and

$$(d) \quad [X_1, X_2] = X_3; \quad [X_2, X_3] = X_1; \quad [X_3, X_1] = X_2 \quad \text{or} \\ [X_1, X_2] = X_3; \quad [X_2, X_3] = -X_1; \quad [X_3, X_1] = X_2. \quad (1.82)$$

In case (a), the algebra is Abelian.

**Example 24.** *The translation algebra in three dimensions  $t(3)$  is an example of Kirillov's case 3a.*

**Example 25.** *The Euclidean algebra  $e(2)$  is Kirillov's case 3c, with  $\alpha = 0, \beta = 1, \gamma = -1, \delta = 0$ .*

**Example 26.** *The algebras  $so(3)$  and  $so(2, 1)$  are examples of Kirillov's case 3d.*

This procedure becomes very cumbersome as the number of elements in the algebra increases. However, it allows one to classify all Lie algebras, including some non-semisimple algebras of physical interest, such as the Euclidean algebras. It has explicitly been carried out to  $r = 8$ . A different method, due to Cartan, provides a classification of semisimple Lie algebras for any number of elements.

## 2 Semisimple Lie Algebras

### 2.1 Cartan-Weyl Form of a (Complex) Semisimple Lie Algebra

In 1895, the French mathematician Cartan provided a way to classify all semisimple Lie algebras. [Elie Cartan, *Sur la Structure des Groupes de Transformation Finis et Continus*, Thèse, Paris, 1894.] The subject was subsequently taken up by Weyl and van der Waerden. [Herman Weyl, *Theorie der Darstellung kontinuierlicher halbeinfacher Gruppen durch lineare Transformationen, I-IV*, Math. Zeitschr. 23, 271 (1925); 24, 328, 377 (1926); B.L. van der Waerden, *Die Gruppentheoretische Methode in der Quantenmechanik*, Math. Zeitschr. 37, 446 (1933).] We begin by rewriting the algebra as

$$X_{\sigma} \equiv (H_i, E_{\alpha}) ; \quad (i = 1, \dots, l) . \quad (2.1)$$

The elements  $H_i$  form the maximal Abelian subalgebra, often called the Cartan subalgebra,

$$[H_i, H_k] = 0 \quad (i, k = 1, \dots, l) . \quad (2.2)$$

The number of elements in the Cartan subalgebra,  $l$ , is called the rank of the algebra. The commutation relations of  $H_i$  with  $E_{\alpha}$  are

$$[H_i, E_{\alpha}] = \alpha_i E_{\alpha} \quad (2.3)$$

while those of the  $E$ 's among themselves are

$$[E_{\alpha}, E_{\beta}] = N_{\alpha\beta} E_{\alpha+\beta} \quad (\text{if } \alpha + \beta \neq 0) \quad (2.4)$$

$$[E_{\alpha}, E_{-\alpha}] = \alpha^i H_i . \quad (2.5)$$

The  $\alpha_i$ 's are called roots and  $N_{\alpha\beta}$  is a normalization. This form of the Lie algebra is called the Cartan-Weyl form.

### 2.2 Graphical Representation of Root Vectors

One considers the  $\alpha_i$ 's ( $i = 1, \dots, l$ ) as the components of a covariant vector lying in an  $l$ -dimensional weight space with scalar product

$$(\alpha, \beta) \equiv \alpha^i \beta_i = \alpha_i \beta^i . \quad (2.6)$$

Van der Waerden derived a set of rules for the algebra to be a semisimple Lie algebra.

**Rule 1** If  $\alpha$  is a root, so is  $-\alpha$ .

**Rule 2** If  $\alpha, \beta$  are roots,  $\frac{2(\alpha, \beta)}{(\alpha, \alpha)}$  is an integer.

**Rule 3** If  $\alpha, \beta$  are roots,  $\beta - 2\alpha \frac{(\alpha, \beta)}{(\alpha, \alpha)}$  is a root.

From these, it follows that the angle  $\varphi$  between roots

$$\cos \varphi = \frac{(\alpha, \alpha)}{\sqrt{(\alpha, \alpha)(\beta, \beta)}} \quad (2.7)$$

can take the values

$$\begin{aligned} \cos^2 \varphi &= 0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1 \\ \varphi &= 0^\circ, 30^\circ, 45^\circ, 60^\circ, 90^\circ \end{aligned} \quad (2.8)$$

Roots can be displayed graphically in a root diagram.

For rank  $l = 1$ , the root diagram is a line, and there is only one possibility (Fig. 2.1).

The algebra, called  $A_1$  by Cartan and  $so(3) \sim su(2)$  by physicists, has three elements,  $r = 3$ .

For rank  $l = 2$ , the root diagram is planar. There are several possibilities (Fig. 2.2):

(a)  $\varphi = 30^\circ$

This algebra called  $G_2$  by Cartan, has 14 elements,  $r = 14$ .

(b)  $\varphi = 45^\circ$

This algebra called  $B_2$  by Cartan and  $so(5)$  by physicists has 10 elements,  $r = 10$ .


(c)  $\varphi = 60^\circ$

This algebra, called  $A_2$  by Cartan and  $su(3)$  by physicists has 8 elements,  $r = 8$ .

(d)  $\varphi = 90^\circ$


This algebra, called  $D_2$  by Cartan and  $so(4)$  by physicists, has  $r = 6$  elements. It can be seen as the direct sum of two  $A_1$  algebras,  $D_2 \sim A_1 \oplus A_1$  or  $so(4) \sim so(3) \oplus so(3)$ .

The Cartan classification of rank-2 algebras contains also the algebra  $C_2$ , called  $sp(4)$  by physicists. The root diagram of  $C_2$  is identical to that of  $B_2$  rotated by  $45^\circ$  (Fig. 2.3).




$so(3) \sim su(2) \sim A_1$

**Fig. 2.1.** Root diagram of the rank  $l = 1$  algebra



**Fig. 2.2.** Root diagrams of rank  $l = 2$  algebras




**Fig. 2.3.** Root diagram of the algebra  $C_2$

The algebras  $C_2$  and  $B_2$  are isomorphic,  $sp(4) \sim so(5)$ .


For rank  $l = 3$  the root diagram is three-dimensional and it will not be drawn here.

### 2.3 Explicit Construction of the Cartan-Weyl Form

For applications, it is of interest to construct explicitly Lie algebras in Cartan-Weyl form. The rank  $l = 1$  algebra,  $A_1$  (Fig. 2.4) has three elements,



**Fig. 2.4.** Root diagram of the algebra  $A_1$  with roots explicitly displayed



**Fig. 2.5.** Root diagram of the algebra  $A_2$  with roots explicitly displayed

$H_1, E_{+1}, E_{-1}$ , with commutation relations,

$$[H_1, H_1] = 0; \quad [H_1, E_{\pm 1}] = \pm E_{\pm 1}; \quad [E_{+1}, E_{-1}] = H_1 \quad (2.9)$$

This algebra, being the angular momentum algebra  $su(2) \sim so(3)$ , is of great interest in quantum mechanics, where it is usually written in terms of the physical angular momentum operators  $J_x, J_y, J_z$ . By taking the combinations

$$J_z = J_z \quad , \quad J_{\pm} = J_x \pm iJ_y \quad (2.10)$$

one obtains the Cartan-Weyl form, with commutation relations

$$[J_z, J_z] = 0; \quad [J_z, J_{\pm}] = \pm J_{\pm}; \quad [J_+, J_-] = 2J_z \quad (2.11)$$

The factor of two in the last commutator is due to a different normalization of the elements of the algebra. One can also see that, in this case, the Abelian Cartan subalgebra is composed of only one element,  $J_z$ , while the Weyl elements are  $J_+$  and  $J_-$ . These elements are called raising and lowering operators.

Another important construction is that of the rank  $l = 2$  algebra,  $A_2$  (Fig. 2.5). This algebra has 8 elements,

$$H_1, H_2, \quad E_{\pm\alpha}, \quad E_{\pm\beta}, \quad E_{\pm(\alpha+\beta)} \quad (2.12)$$

with commutation relations

$$\begin{aligned} [H_1, E_{\pm\alpha}] &= \pm \frac{1}{2\sqrt{3}} E_{\pm\alpha} & [H_2, E_{\pm\alpha}] &= \pm \frac{1}{2} E_{\pm\alpha} \\ [H_1, E_{\pm\beta}] &= \pm \frac{1}{2\sqrt{3}} E_{\pm\beta} & [H_2, E_{\pm\beta}] &= \pm \frac{1}{2} E_{\pm\beta} \\ [H_1, E_{\pm(\alpha+\beta)}] &= \pm \frac{1}{\sqrt{3}} E_{\pm(\alpha+\beta)} & [H_2, E_{\pm(\alpha+\beta)}] &= 0 \end{aligned}$$

$$\begin{aligned}
 [E_\alpha, E_{-\alpha}] &= \frac{H_1}{2\sqrt{3}} + \frac{H_2}{2} & [E_\beta, E_{-\beta}] &= \frac{H_1}{2\sqrt{3}} - \frac{H_2}{2} \\
 [E_{(\alpha+\beta)}, E_{-(\alpha+\beta)}] &= \frac{H_1}{\sqrt{3}} = \frac{H_1}{\sqrt{3}} & [E_\alpha, E_\beta] &= \frac{1}{\sqrt{6}} E_{\alpha+\beta} \\
 [E_\alpha, E_{(\alpha+\beta)}] &= 0 & [E_\beta, E_{(\alpha+\beta)}] &= 0 \\
 [E_\alpha, E_{-(\alpha+\beta)}] &= -\frac{1}{\sqrt{6}} E_{-\beta} & [E_\beta, E_{-(\alpha+\beta)}] &= \frac{1}{\sqrt{6}} E_{-\alpha} \\
 [H_i, H_j] &= 0 \quad (i, j = 1, 2) & & (2.13)
 \end{aligned}$$

The algebra has two Cartan elements and six Weyl elements (raising and lowering operators).


### 2.4 Dynkin Diagrams

The root diagrams for rank  $l \geq 3$  cannot be displayed easily. The Russian mathematician Dynkin devised a method to display root diagrams of all semi-simple Lie algebras. [E.B. Dynkin, *The Structure of Semisimple Lie Algebras*, Usp. Mat. Nauk (N.S.), 2, 59 (1947). Transl. in Am. Math. Soc. Transl. (1), 9, 308 (1962).] We begin by introducing the notion of positive root.

**Definition 1.** *Positive roots,  $\alpha^{(+)}$ , are those for which, in some arbitrary frame, its first coordinate different from zero is positive.*

The number of positive roots is half of non-null roots.

**Example 1.** *Consider the root diagram of  $B_2$  The roots are  $(1, 0)$   $(1, 1)$   $(0, 1)$   $(-1, 1)$   $(-1, 0)$   $(-1, -1)$   $(0, -1)$   $(1, -1)$ . The positive roots are  $(1, 0)$   $(1, 1)$   $(0, 1)$   $(1, -1)$ , with sum  $\sum \alpha^{(+)} = (3, 1)$ . We next introduce the notion of simple roots.*



**Fig. 2.6.** Root diagram of the algebra  $B_2$  with roots explicitly displayed



**Definition 2.** A simple root is a positive root which cannot be decomposed into the sum of positive roots.


In the case of  $B_2$  (Fig. 2.6), the two roots  $(1, 0)$  and  $(1, 1)$  can be decomposed as  $(1, 0) = (1, -1) + (0, 1)$  and  $(1, 1) = (1, 0) + (0, 1)$ . The simple roots are thus only  $\alpha \equiv (0, 1)$  and  $\beta \equiv (1, -1)$ . Dynkin showed that the angle between two simple roots can only be  $90^\circ, 120^\circ, 135^\circ, 150^\circ$  and the normalization

$$\frac{(\beta, \beta)}{(\alpha, \alpha)} = \begin{cases} 1 & \vartheta_{\alpha, \beta} = 120^\circ \\ 2 & \vartheta_{\alpha, \beta} = 135^\circ \\ 3 & \vartheta_{\alpha, \beta} = 150^\circ \\ \text{undetermined} & \vartheta_{\alpha, \beta} = 90^\circ \end{cases} \quad (2.14)$$


The root  $\alpha$  is called short and the root  $\beta$  is called long. When plotted as before, the simple root diagram of  $B_2$  appears as in (Fig. 2.7). All information on the Lie algebra can then be condensed into a Dynkin diagram. In this diagram, the angle  $\vartheta$  between roots is indicated by

$$\begin{aligned} \vartheta = 120^\circ & \quad \text{single line} \\ \vartheta = 135^\circ & \quad \text{double line} \\ \vartheta = 150^\circ & \quad \text{triple line} \\ \vartheta = 90^\circ & \quad \text{not joined} \end{aligned} \quad (2.15)$$

A short root is indicated by a filled dot  $\bullet$ , while a long root is indicated by an open dot  $\circ$ .



**Fig. 2.7.** Simple roots of  $B_2$



**Fig. 2.8.** Dynkin diagrams of rank-2 algebras

**Example 2.** *Dynkin diagrams of rank two algebras*

Dynkin diagrams of rank two algebras are shown in (Fig. 2.8).

## 2.5 Classification of (Complex) Semisimple Lie Algebras

All complex semisimple Lie algebras have been classified by Cartan and are given in Table 2.1. Because of its importance, the non-semisimple Lie algebra  $u(n)$  must be included as well. This algebra is of order  $n^2$  and rank  $n$ .

**Table 2.1.** Cartan classification of complex semisimple Lie algebras (and  $u(n)$ )

Name	Label	Cartan	Order ( $r$ )	Rank ( $l$ )
Special Unitary	$su(n)$	$A_l$	$n^2 - 1$	$n - 1$
[Special] Orthogonal	$so(n)$ ( $n$ odd)	$B_l$	$n(n - 1)/2$	$(n - 1)/2$
Symplectic	$sp(n)$ ( $n$ even)	$C_l$	$n(n + 1)/2$	$n/2$
[Special] Orthogonal	$so(n)$ ( $n$ even)	$D_l$	$n(n - 1)/2$	$n/2$
Exceptional	$G_2$	$G_2$	14	2
	$F_4$	$F_4$	52	4
	$E_6$	$E_6$	78	6
	$E_7$	$E_7$	133	7
	$E_8$	$E_8$	248	8
Unitary	$u(n)$	—	$n^2$	$n$

There is no difference between orthogonal and special orthogonal algebras and hence “special” has been put in brackets in the table. The Abelian algebra  $so(2) \sim u(1)$  composed of a single element is included in the Cartan classification under  $so(n)$   $n = \text{even}$ , although strictly speaking it is not possible to apply to it Cartan’s criterion since it does not have any subalgebra except itself.

## 2.6 Real Forms of Complex Semisimple Lie Algebras

In applications, one is often interested in real forms of complex semi-simple Lie algebras. Commonly used real forms and their notation are given in Table 2.2.

## 2.7 Isomorphisms of Complex Semisimple Lie Algebras

Isomorphisms of complex Lie algebras of low rank are given in Table 2.3.

**Table 2.2.** Real forms of complex semisimple Lie algebras

Cartan	Real Forms
$A_l$	$su(n)$ $su(p, q) \quad (p + q = n)$ $sl(n, \mathbb{R})$ $su^*(2n)$
$B_l$	$so(n) \quad (n \text{ odd})$ $so(p, q) \quad (p + q = n)$
$C_l$	$sp(n) \quad (n \text{ even})$ $sp(n, \mathbb{R})$ $sp(p, q) \quad (p + q = n)$
$D_l$	$so(n) \quad (n \text{ even})$ $so(p, q)$ $so^*(n)$

**Table 2.3.** Isomorphisms of complex semisimple Lie algebra

$$\begin{array}{c}
 \overline{A_1 \sim B_1 \sim C_1} \\
 B_2 \sim C_2 \\
 D_2 \sim A_1 \oplus A_1 \\
 \overline{A_3 \sim D_3}
 \end{array}$$

## 2.8 Isomorphisms of Real Lie Algebras

Isomorphisms of real Lie algebras of low rank are given in Table 2.4.

**Table 2.4.** Isomorphisms of real Lie algebras

$A_1 \sim B_1 \sim C_1$	$su(2) \sim so(3) \sim sp(2) \sim su^*(2)$ $su(1, 1) \sim so(2, 1) \sim sp(2, \mathbb{R}) \sim sl(2, \mathbb{R})$
$B_2 \sim C_2$	$so(5) \sim sp(4)$ $so(4, 1) \sim sp(2, 2)$ $so(3, 2) \sim sp(4, \mathbb{R})$
$D_2 \sim A_1 \oplus A_1$	$so(4) \sim su(2) \oplus su(2) \sim so(3) \oplus so(3) \sim sp(2) \oplus sp(2)$ $so^*(4) \sim su(2) \oplus sl(2, \mathbb{R})$ $so(3, 1) \sim sl(2, \mathbb{C})$ $so(2, 2) \sim sl(2, \mathbb{R}) \oplus sl(2, \mathbb{R})$
$A_3 \sim D_3$	$su(4) \sim so(6)$ $su(3, 1) \sim so^*(6)$ $su^*(4) \sim so(5, 1)$ $sl(4, \mathbb{R}) \sim so(3, 3)$ $su(2, 2) \sim so(4, 2)$



with unit entry on the  $\beta$ -th column and the  $\alpha$ -th row. Commutation relations of the algebra are obtained from the basic commutation relations of matrices. Matrix realizations will be discussed in Chap. 10.

- (iii) Boson creation-annihilation operator realization (often called Jordan-Schwinger realization)

This is in terms of bilinear products of  $n$  boson creation,  $b_\alpha^\dagger$ , and annihilation,  $b_\alpha$ , operators acting on a vacuum  $|0\rangle$ ,

$$E_{\alpha\beta} = b_\alpha^\dagger b_\beta. \quad (2.20)$$

The commutation relations of the algebra can be obtained from those of the creation and annihilation operators

$$[b_\alpha, b_\beta^\dagger] = \delta_{\alpha\beta}, \quad (2.21)$$

called Bose commutation relations. Realizations in terms of boson creation and annihilation operators will be discussed in Chap. 7.

For all three realizations, the commutation relations of the elements of the algebra  $E_{\alpha\beta}$  are

$$[E_{\alpha\beta}, E_{\gamma\delta}] = \delta_{\beta\gamma} E_{\alpha\delta} - \delta_{\alpha\delta} E_{\gamma\beta} \quad (2.22)$$

with  $\alpha, \beta = 1, \dots, n$ . They define the Lie algebra of  $u(n)$ . Realizations of other algebras can be obtained by taking appropriate combinations of the elements  $E_{\alpha\beta}$ , since any Lie algebra is, by Ado's theorem, a subalgebra of  $u(n)$ .

## 2.11 Other Realizations of Lie Algebras

In addition to the three realizations of the previous section, others are possible and have been used for applications in physics. Two of these, particularly important in the description of fermionic systems, are:

- (iv) Grassmann differential realization

This is in terms of Grassmann variables,  $\theta_i$  ( $i = 1, \dots, n$ ), and their derivatives,  $\frac{\partial}{\partial\theta_i}$ . The elements are

$$E_{ij} = \theta_i \frac{\partial}{\partial\theta_j}. \quad (2.23)$$

The commutation relations can be obtained from the basic commutation relations

$$\left[ \frac{\partial}{\partial\theta_i}, \theta_j \right]_+ \equiv \frac{\partial}{\partial\theta_i} \theta_j + \theta_j \frac{\partial}{\partial\theta_i} = \delta_{ij}. \quad (2.24)$$

These realizations will not be discussed in these notes. A detailed account is given by Berezin. [F.A. Berezin, *An Introduction to Superanalysis*, D. Riedel Publishing, Dordrecht, 1987.]

(v) Fermion creation-annihilation operators realization

This is in terms of bilinear products of  $n$  fermion creation,  $a_i^\dagger$ , and annihilation,  $a_i$ , operators acting on a vacuum  $|0\rangle$ ,

$$E_{ij} = a_i^\dagger a_j , \quad (2.25)$$

where the creation and annihilation operators,  $a_i^\dagger, a_i$ , satisfy

$$[a_i, a_j^\dagger]_+ \equiv a_i a_j^\dagger + a_j^\dagger a_i = \delta_{ij} , \quad (2.26)$$

called Fermi commutation relations. These realizations will be discussed in Chap. 8.

For both realizations (iv) and (v), the commutation relations of the elements of the algebra  $E_{ij}$  are

$$[E_{ij}, E_{km}] = \delta_{jk} E_{im} - \delta_{im} E_{kj} , \quad (2.27)$$

with  $i, j = 1, \dots, n$ . The commutation relations (2.28) are identical those given by (2.23). They define again the Lie algebra  $u(n)$ .

# 3 Lie Groups

## 3.1 Groups of Transformations

Associated with each Lie algebra there is a Lie group. The notion of groups of transformations precedes that of Lie algebras and is more general than it. A set of elements  $A, B, C \dots$ , forms a group  $G$  if it satisfies the following axioms:

**Axiom 1.** *Among the elements there is an element  $I$  such that*

$$AI = IA = A \tag{3.1}$$

This property is called identity.

**Axiom 2.** *The product  $AB$  gives another element  $C$  in the set*

$$AB = C \tag{3.2}$$

This property is called closure.

**Axiom 3.** *There exists an element  $A^{-1}$  such that*

$$A^{-1}A = AA^{-1} = I \tag{3.3}$$

This property is called inverse.

**Axiom 4.** *The order of multiplication is immaterial*

$$A(BC) = (AB)C \tag{3.4}$$

This property is called associativity.

## 3.2 Groups of Matrices

Particularly important are groups of square matrices

$$A = \begin{pmatrix} \dots \\ \dots \\ \dots \end{pmatrix} \quad (n \times n) . \tag{3.5}$$

These matrices satisfy all the axioms of a Lie group:

(i) The identity  $I$  is the unit matrix

$$I = \begin{pmatrix} 1 & & & 0 \\ & 1 & & \\ & & \ddots & \\ & & & 1 \\ 0 & & & & 1 \end{pmatrix} \quad (3.6)$$

(ii) Matrix multiplication gives closure.

(iii) If  $\det |A| \neq 0$  an inverse  $A^{-1}$  exists.

(iv) Matrix multiplication gives associativity.

Groups of matrices can be written in terms of all number fields,  $R, C, Q, O$ . In these notes, we shall consider only groups of real and complex matrices. The matrix elements of the matrix  $A$  will be denoted by  $A_{ik}$ , with  $i = \text{row index}$  and  $k = \text{column index}$ . We shall also introduce real and complex vectors in  $n$  dimensions. The components of vectors will be denoted by  $x_i$  and  $z_i$ . Standard matrix notation will be used in this chapter (no covariant or contravariant indices).

### 3.3 Properties of Matrices

We begin by recalling in Table 3.1 some basic properties of matrices.

**Table 3.1.** Matrix properties

$A = A^t$	Symmetric
$A = -A^t$	Skew symmetric
$A^t A = I$	Orthogonal
$A = A^*$	Real
$A = -A^*$	Imaginary
$A = A^\dagger$	Hermitian
$A = -A^\dagger$	Skew hermitian
$A^\dagger A = I$	Unitary

In this table,  $A^t$  denotes the transpose matrix,  $A^*$  the complex conjugate matrix, and  $A^\dagger$  the hermitian conjugate matrix,  $A^\dagger = (A^t)^*$ .

A group of transformations transforms the real or complex vector  $\mathbf{x} \equiv (x_1, x_2, \dots, x_n)$  or  $\mathbf{z} \equiv (z_1, z_2, \dots, z_n)$  into the real or complex vector  $\mathbf{x}'$  or  $\mathbf{z}'$ . We shall consider both real and complex transformations



$$\begin{aligned} \mathbf{x}' &= A\mathbf{x} \quad ; \quad x'_i = \sum_k A_{ik}x_k \\ \mathbf{z}' &= B\mathbf{z} \quad ; \quad z'_i = \sum_k B_{ik}z_k \end{aligned} \tag{3.7}$$

where  $A_{ik}$  and  $B_{ik}$  are the matrix elements of the real and complex  $n \times n$  matrices  $A$  and  $B$ .

### 3.4 Continuous Matrix Groups

- (a) General linear groups

These are the most general linear transformations. They are denoted by

$$\begin{aligned} GL(n, C) & \quad r = 2n^2 \\ GL(n, R) & \quad r = n^2 . \end{aligned} \tag{3.8}$$

The number of real parameters that characterize the transformation is given next to its name. The number field  $R, C$  is also explicitly shown.

- (b) Special linear groups

If, on the general linear transformation, the condition

$$\det | A | = +1 \tag{3.9}$$

is imposed, the group is called special linear group, denoted by

$$\begin{aligned} SL(n, C) & \quad r = 2(n^2 - 1) \\ SL(n, R) & \quad r = n^2 - 1 . \end{aligned} \tag{3.10}$$

- (c) Unitary groups

Imposing the condition

$$A^\dagger A = I \tag{3.11}$$

one obtains the unitary groups

$$\begin{aligned} U(n, C) & \equiv U(n) & r &= n^2 \\ U(p, q; C) & \equiv U(p, q) & r &= n^2 . \end{aligned} \tag{3.12}$$

They leave invariant the quantities

$$\begin{aligned} U(n) : & \quad \sum_{i=1}^n z_i z_i^* \\ U(p, q) : & \quad - \sum_{i=1}^p z_i z_i^* + \sum_{j=p+1}^{p+q} z_j z_j^* \end{aligned} \tag{3.13}$$

Unitary groups are over complex numbers  $C$ . It has become common practice to delete the number field from the group notation, that is to use  $U(n)$  instead of  $U(n, C)$ .

## (d) Special unitary groups

The combination of the special condition with the unitary condition

$$A^\dagger A = I, \quad \det |A| = +1 \quad (3.14)$$

gives the special unitary groups

$$\begin{aligned} SU(n, C) &\equiv SU(n) & r = n^2 - 1 \\ SU(p, q; C) &\equiv SU(p, q) & r = n^2 - 1. \end{aligned} \quad (3.15)$$

Again, the number field  $C$  is often deleted. For special unitary groups, there is an (anomalous) case, denoted by  $SU^*(2n)$ ,

$$SU^*(2n) \quad r = n^2 - 1 \quad (3.16)$$

defined by matrices

$$A = \begin{pmatrix} A_1 & A_2 \\ -A_2^* & A_1^* \end{pmatrix}$$

$$A_1, A_2 = n \times n \text{ complex matrices with } \text{Tr } A_1 + \text{Tr } A_1^* = 0. \quad (3.17)$$

## (e) Orthogonal groups

These groups are defined by the orthogonality condition

$$A^t A = I. \quad (3.18)$$

In applications in physics, they are usually over the real number field. The number field is often deleted in the notation and  $O(n, R)$  is often denoted by  $O(n)$

$$\begin{aligned} O(n, C) & \quad r = n(n-1) \\ O(n, R) &\equiv O(n) \quad r = \frac{1}{2} n(n-1). \end{aligned} \quad (3.19)$$

They leave invariant the quantities

$$\begin{aligned} O(n, C) : & \quad \sum_{i=1}^n z_i^2 \\ O(n, R) &\equiv O(n) : \quad \sum_{i=1}^n x_i^2 \end{aligned} \quad (3.20)$$

In addition, one has the groups

$$\begin{aligned} O(p, q; C) & \quad r = n(n-1) \\ O(p, q, R) & \quad r = \frac{1}{2} n(n-1) \end{aligned} \quad (3.21)$$

which leave invariant the quantities

$$\begin{aligned}
 O(p, q; C) : & \quad - \sum_{i=1}^p z_i^2 + \sum_{j=p+1}^{p+q} z_j^2 \\
 O(p, q; R) \equiv O(p, q) : & \quad - \sum_{i=1}^p x_i^2 + \sum_{j=p+1}^{p+q} x_j^2
 \end{aligned} \tag{3.22}$$

(f) Special orthogonal groups

The combination of the special with the orthogonal condition

$$A^t A = I \quad \det |A| = +1 \tag{3.23}$$

gives the special orthogonal groups

$$\begin{aligned}
 SO(n, C) & \quad r = n(n - 1) \\
 SO(n, R) & \quad r = \frac{1}{2}n(n - 1) \\
 SO(p, q; C) & \quad r = n(n - 1) \\
 SO(p, q; R) & \quad r = \frac{1}{2}n(n - 1) .
 \end{aligned} \tag{3.24}$$

Also here there is an (anomalous) case, called  $SO^*(2n)$ , described by matrices

$$A = \begin{pmatrix} A_1 & A_2 \\ -A_2^* & A_1 \end{pmatrix}$$

$A_1, A_2 = n \times n$  complex matrices with  $A_1 = -A_1^\dagger$  and  $A_2 = A_2^\dagger$ . (3.25)

Real orthogonal groups are used both in quantum and in classical mechanics.

(g) Symplectic groups

To define these groups, the vectors  $\mathbf{x}$  and  $\mathbf{y}$  are divided into two pieces,  $\mathbf{x} = (x_1, \dots, x_n; x'_1, \dots, x'_n)$ ,  $\mathbf{y} = (y_1, \dots, y_n; y'_1, \dots, y'_n)$ . Symplectic groups

$$\begin{aligned}
 Sp(2n, C) & \quad r = 2n(2n + 1) \\
 Sp(2n, R) & \quad r = \frac{1}{2}2n(2n + 1)
 \end{aligned} \tag{3.26}$$

are defined as those groups that leave invariant the quantity

$$\sum_{i=1}^n (x_i y'_i - y_i x'_i) \tag{3.27}$$

where the vectors can be either real or complex. If the unitary condition is imposed

$$A^\dagger A = I \tag{3.28}$$

the group is called unitary symplectic

$$USp(2n, C) \equiv Sp(2n) \quad r = \frac{1}{2} 2n(2n + 1) \quad (3.29)$$

and is often denoted by  $Sp(2n)$ .

Both real and complex symplectic groups are used in quantum mechanics, while real symplectic groups are used in classical mechanics (canonical transformations and Hamilton's equations).

## 3.5 Examples of Groups of Transformations

### 3.5.1 The Rotation Group in Two Dimensions, $SO(2)$

As a first example we consider the rotation group in two dimensions  $SO(2) \equiv SO(2, R)$ . Under a general linear transformation the two coordinates  $x, y$  (used here to conform with usual physics notation) transform as

$$\begin{aligned} x' &= a_{11} x + a_{12} y \\ y' &= a_{21} x + a_{22} y \end{aligned} \quad (3.30)$$

The corresponding group,  $GL(2, R)$ , is a 4 parameter group. The invariance of  $x^2 + y^2$

$$a_{11}^2 x^2 + a_{12}^2 y^2 + 2 a_{11} a_{12} xy + a_{21}^2 x^2 + a_{22}^2 y^2 + 2 a_{21} a_{22} xy = x^2 + y^2 \quad (3.31)$$

gives 3 conditions

$$\begin{aligned} a_{11}^2 + a_{21}^2 &= 1 \\ 2a_{11}a_{12} + 2a_{21}a_{22} &= 0 \\ a_{12}^2 + a_{22}^2 &= 1. \end{aligned} \quad (3.32)$$

This leaves only 1 parameter.

**Example 1.** *The group  $SO(2)$  is a one parameter group*

The parameter can be chosen as the angle of rotation,  $\varphi$ ,

$$\begin{aligned} x' &= (\cos \varphi) x - (\sin \varphi) y \\ y' &= (\sin \varphi) x + (\cos \varphi) y \end{aligned} \quad (3.33)$$

as shown in Fig. 3.1.




Fig. 3.1. The angle  $\varphi$  that parametrizes  $SO(2)$

### 3.5.2 The Lorentz Group in One Plus One Dimension, $SO(1, 1)$

A group closely related to the rotation group is the Lorentz group  $SO(1, 1) \equiv SO(1, 1; R)$ . The general linear transformation in space-time,  $x, t$  can be written

$$\begin{aligned} x' &= a_{11}x + a_{12}t \\ t' &= a_{21}x + a_{22}t . \end{aligned} \tag{3.34}$$

Imposing the condition  $x^2 - t^2 = \text{invariant}$ , leaves a 1 parameter group.

**Example 2.** *The group  $SO(1,1)$  is a one parameter group*

A convenient parametrization is in term of the boost,  $\vartheta$ .

$$\begin{aligned} x' &= (\cosh \vartheta)x + (\sinh \vartheta)t \\ t' &= (\sinh \vartheta)x + (\cosh \vartheta)t . \end{aligned} \tag{3.35}$$

By comparing with the previous subsection, one can see that the invariant forms are different (Fig. 3.2).

The group  $SO(2)$  which leaves invariant the form  $x^2 + y^2$  is said to be compact, while the group  $SO(1, 1)$  which leaves invariant  $x^2 - t^2$  is said to be non-compact. The notation is such that the number of plus or minus signs is indicated in  $SO(p, q)$ .




Fig. 3.2. Invariant forms of (a)  $SO(2)$  and (b)  $SO(1, 1)$

### 3.5.3 The Rotation Group in Three Dimensions

As another example consider the rotation group in three dimensions  $SO(3) \equiv SO(3, R)$ . Under a general linear transformation,  $GL(3, R)$ , the coordinates  $x, y, z$  transform as

$$\begin{aligned} x' &= a_{11} x + a_{12} y + a_{13} z \\ y' &= a_{21} x + a_{22} y + a_{23} z \\ z' &= a_{31} x + a_{32} y + a_{33} z \end{aligned} \tag{3.36}$$

This is a 9 parameter group. Orthogonality

$$x'^2 + y'^2 + z'^2 = x^2 + y^2 + z^2 \tag{3.37}$$

gives 6 conditions. We thus have a three parameter group.

**Example 3.** *The group  $SO(3)$  is a three parameter group*


A convenient parametrization is in terms of Euler angles,  $\varphi, \vartheta, \psi$ ,

$$\begin{pmatrix} \cos \varphi \cos \vartheta \cos \psi - \sin \varphi \sin \psi & -\cos \varphi \cos \vartheta \sin \psi - \sin \varphi \cos \psi & \cos \varphi \sin \vartheta \\ \sin \varphi \cos \vartheta \cos \psi + \cos \varphi \sin \psi & -\sin \varphi \cos \vartheta \sin \psi + \cos \varphi \cos \psi & -\sin \varphi \sin \vartheta \\ -\sin \vartheta \cos \psi & \sin \vartheta \sin \psi & \cos \vartheta \end{pmatrix} \tag{3.38}$$

as shown in Fig. 3.3. The rotation matrices (3.39) are usually denoted by  $R(\varphi, \vartheta, \psi)$ .

### 3.5.4 The Special Unitary Group in Two Dimensions, $SU(2)$

This group is denoted by  $SU(2) \equiv SU(2, C)$ . Under a general linear complex transformation,  $GL(2, C)$ , the complex quantities,  $u, v$ , called a spinor, transform as



**Fig. 3.3.** The Euler angles  $\varphi, \vartheta, \psi$  that parametrize  $SO(3)$

$$\begin{aligned} u' &= a_{11} u + a_{12} v \\ v' &= a_{21} u + a_{22} v . \end{aligned} \tag{3.39}$$

This is a 8 parameter group. Call the matrix of the transformation  $A$

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \quad A^\dagger = \begin{pmatrix} a_{11}^* & a_{21}^* \\ a_{12}^* & a_{22}^* \end{pmatrix} \tag{3.40}$$

Unitarity,  $A^\dagger A = 1$ , gives four conditions

$$\begin{aligned} a_{11}^* a_{11} + a_{21}^* a_{21} &= 1 \\ a_{11}^* a_{12} + a_{21}^* a_{22} &= 0 \\ a_{12}^* a_{11} + a_{22}^* a_{21} &= 0 \\ a_{12}^* a_{12} + a_{22}^* a_{22} &= 1 \end{aligned} \tag{3.41}$$

The corresponding group,  $U(2)$ , is a four parameter group. If one imposes a further condition  $\det | A | = +1$ , that is

$$a_{11} a_{22} - a_{12} a_{21} = 1 \tag{3.42}$$

one obtains the three parameter group  $SU(2)$ .

**Example 4.** *The group  $SU(2)$  is a three parameter group*

This group can be parametrized as

$$\begin{aligned} u' &= a_{11} u + a_{12} v \\ v' &= -a_{12}^* u + a_{11}^* v \end{aligned} \tag{3.43}$$

with

$$a_{11} a_{11}^* + a_{12} a_{12}^* = 1 . \tag{3.44}$$

### 3.5.5 Relation Between $SO(3)$ and $SU(2)$

Both  $SO(3)$  and  $SU(2)$  are three parameter groups. It is of importance to find their relationship. Consider the following combination of the complex spinor  $u, v$

$$x_1 = u^2 ; x_2 = uv ; x_3 = v^2 . \tag{3.45}$$

These combinations transform as

$$\begin{aligned} x'_1 &= u'^2 = a_{11}^2 x_1 + 2a_{11} a_{12} x_2 + a_{12}^2 x_3 \\ x'_2 &= u'v' = -a_{11} a_{12}^* x_1 + (a_{11} a_{11}^* - a_{12} a_{12}^*) x_2 + a_{11}^* a_{12} x_3 \\ x'_3 &= v'^2 = a_{12}^{*2} x_1 - 2a_{11}^* a_{12}^* x_2 + a_{11}^{*2} x_3 \end{aligned} \tag{3.46}$$

By introducing the coordinates  $x, y, z$

$$x = (x_1 - x_3)/2; \quad y = (x_1 + x_3)/2i; \quad z = x_2 \quad (3.47)$$

one can see that they transform as

$$\begin{aligned} x' &= \frac{1}{2}(a_{11}^2 - a_{12}^{*2} - a_{12}^2 + a_{11}^{*2})x + \frac{i}{2}(a_{11}^2 - a_{12}^{*2} + a_{12}^2 - a_{11}^{*2})y \\ &\quad + (a_{11}a_{12} + a_{11}^*a_{12}^*)z \\ y' &= \frac{-i}{2}(a_{11}^2 - a_{12}^{*2} - a_{12}^2 + a_{11}^{*2})x + \frac{1}{2}(a_{11}^2 - a_{12}^{*2} + a_{12}^2 + a_{11}^{*2})y \\ &\quad - i(a_{11}a_{12} - a_{11}^*a_{12}^*)z \\ z' &= -(a_{11}^*a_{12} + a_{11}a_{12}^*)x + i(a_{11}^*a_{12} - a_{11}a_{12}^*)y \\ &\quad + (a_{11}a_{12}^* - a_{12}a_{11}^*)z \end{aligned} \quad (3.48)$$

This is a real orthogonal transformation in three dimensions, satisfying

$$x'^2 + y'^2 + z'^2 = x^2 + y^2 + z^2. \quad (3.49)$$

Thus  $SU(2)$  and  $SO(3)$  are related by a change of variables. In order to elucidate the correspondence between  $SU(2)$  and  $SO(3)$ , we consider a rotation of an angle  $\alpha$  around the  $z$ -axis. By inserting the values  $a_{11} = e^{i\alpha/2}, a_{12} = 0$  in the appropriate formulas, we see that this rotation is characterized by matrices

$$\begin{array}{cc} SU(2) & SO(3) \\ \left( \begin{array}{cc} e^{i\alpha/2} & 0 \\ 0 & e^{-i\alpha/2} \end{array} \right) & \left( \begin{array}{ccc} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{array} \right) \end{array} \quad (3.50)$$

A generic rotation, by angles  $\alpha, \beta, \gamma$  is instead characterized by matrices

$$\begin{array}{cc} SU(2) & SO(3) \\ \left( \begin{array}{cc} \cos \frac{\beta}{2} e^{\frac{i}{2}(\alpha+\gamma)} & \sin \frac{\beta}{2} e^{-\frac{i}{2}(\alpha-\gamma)} \\ -\sin \frac{\beta}{2} e^{\frac{i}{2}(\alpha-\gamma)} & \cos \frac{\beta}{2} e^{-\frac{i}{2}(\alpha+\gamma)} \end{array} \right) & R(\alpha, \beta, \gamma) \end{array} \quad (3.51)$$

where  $R(\alpha, \beta, \gamma)$  is given in (3.39). For no rotation,  $R(0, 0, 0)$ , the correspondence is

$$\begin{array}{cc} SU(2) & SO(3) \\ \left( \begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array} \right) & \left( \begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right) \end{array} \quad (3.52)$$

while for rotation of  $2\pi$ ,  $R(2\pi, 0, 0)$ , the correspondence is

$$\begin{array}{cc} SU(2) & SO(3) \\ \left( \begin{array}{cc} -1 & 0 \\ 0 & -1 \end{array} \right) & \left( \begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right) \end{array} \quad (3.53)$$



One can see that there is a two-to-one correspondence, called a homomorphic mapping of  $SU(2)$  into  $SO(3)$ , denoted by  $SU(2) \approx SO(3)$ . One says that  $SU(2)$  is the universal covering group of  $SO(3)$ .

### 3.6 Lie Algebras and Lie Groups

#### 3.6.1 The Exponential Map

The relationship between Lie algebras and Lie groups is of great importance. Let the Lie algebra be  $g$  and the corresponding Lie group  $G$ . The relation is

$$\text{Lie algebra } g \ni X_i \quad (i = 1, \dots, r) \tag{3.54}$$

$$\text{Lie group } G \ni \exp \left( \sum_{i=1}^r \alpha_i X_i \right) \tag{3.55}$$

where the  $\alpha_i$ 's are the parameters of the group and the sum goes over the order of the group. (The  $\alpha_i$ 's here should not be confused with the  $\alpha_i$ 's in Chap. 2 where they denote the components of a root vector). This relationship is called an exponential map and denoted by

$$\begin{array}{c} g \\ \downarrow \text{exp} \\ G \end{array} \tag{3.56}$$

**Example 5.** *The Lie group  $SO(3)$  is*

$$A(\alpha_1, \alpha_2, \alpha_3) = e^{\alpha_1 X_1 + \alpha_2 X_2 + \alpha_3 X_3} \tag{3.57}$$

#### 3.6.2 Definition of Exp

The exponentiation is defined through a power series expansion. For rank one algebras, with only one element  $X$  and one parameter  $\alpha$

$$e^{\alpha X} = 1 + \alpha X + \frac{\alpha^2 X^2}{2!} + \dots = \sum_{p=0}^{\infty} \frac{(\alpha X)^p}{p!} \tag{3.58}$$

The infinitesimal group element is obtained by keeping only the linear term in the expansion

$$e^{\alpha X} \xrightarrow{\alpha \rightarrow 0} 1 + \alpha X \tag{3.59}$$

For algebras of larger rank, one needs to exponentiate non-commuting elements. It is convenient to use matrices.

### 3.6.3 Matrix Exponentials

Let  $A$  be a  $n \times n$  matrix. Then

$$e^A = I + A + \frac{A^2}{2!} + \dots \quad (3.60)$$

Some properties of matrix exponentials are:

- (i) The exponential  $e^A$  converges if the matrix elements  $|a_{ij}|$  have an upper bound, that is the group is compact.
- (ii) If  $A$  and  $B$  commute, then

$$e^{A+B} = e^A e^B. \quad (3.61)$$

- (iii) If  $B$  can be inverted, then

$$B e^A B^{-1} = e^{BAB^{-1}} \quad (3.62)$$

- (iv) If  $\lambda_1, \lambda_2, \dots, \lambda_n$  are eigenvalues of  $A$ , then

$$e^{\lambda_1}, \dots, e^{\lambda_n} \quad (3.63)$$

are eigenvalues of  $e^A$ .

- (v) The exponential series satisfies

$$\begin{aligned} e^{A^*} &= (e^A)^* & (e^{A^t}) &= (e^A)^t \\ e^{A^\dagger} &= (e^A)^\dagger & e^{-A} &= (e^A)^{-1} \end{aligned} \quad (3.64)$$

- (vi) The determinant of  $e^A$  is  $e^{\text{tr}A}$ .
- (vii) If  $A$  is skew symmetric,  $e^A$  is orthogonal. If  $A$  is skew hermitian,  $e^A$  is unitary.
- (viii) The following formula (Campbell-Hausdorff) applies

$$e^{-A} B e^A = B + \frac{1}{1!} [B, A] + \frac{1}{2!} [[B, A], A] + \dots \quad (3.65)$$

# 4 Irreducible Bases (Representations)

## 4.1 Definitions

An irreducible basis (*IRB*) is the basis for the representations of the algebra,  $g$ , (and the associated group,  $G$ ), and the basis upon which the elements of the algebra,  $X_\rho \in g$ , act. It will be denoted by  $\mathcal{B}$ .

Also, if  $\mathcal{V}$  is a linear vector space,

$$\mathcal{B} : \mathcal{V} \oplus (\mathcal{V} \oplus \mathcal{V}) \oplus (\mathcal{V} \oplus \mathcal{V} \oplus \mathcal{V}) \oplus \dots \tag{4.1}$$

The meaning of the term irreducible is that any element  $X_\rho$  acting on  $\mathcal{B}$  does not lead out of  $\mathcal{B}$ .

## 4.2 Abstract Characterization

Irreducible representations are characterized by a set of labels (often called quantum numbers). For a semisimple Lie algebra  $g$ , the number of labels is the rank of the algebra,  $l$ , which is also the number of Cartan commuting elements. The number of labels and their notation in these lecture notes are shown in Table 4.1.

**Table 4.1.** Labels of irreducible representations

Name	Number of Labels	Labels
$u(n)$	$n$	$[\lambda_1, \lambda_2, \dots, \lambda_n]$
$su(n)$	$n - 1$	$[\lambda_1, \lambda_2, \dots, \lambda_{n-1}]$
$so(n)$ ( $n$ odd)	$\nu = (n - 1)/2$	$[\mu_1, \mu_2, \dots, \mu_\nu]$
$sp(n)$ ( $n$ even)	$\nu = n/2$	$[\mu_1, \mu_2, \dots, \mu_\nu]$
$so(n)$ ( $n$ even)	$\nu = n/2$	$[\mu_1, \mu_2, \dots, \mu_\nu]$
$G_2$	2	$[\gamma_1, \gamma_2]$
$F_4$	4	$[\gamma_1, \gamma_2, \gamma_3, \gamma_4]$
$E_6$	6	$[\gamma_1, \gamma_2, \gamma_3, \gamma_4, \gamma_5, \gamma_6]$
$E_7$	7	$[\gamma_1, \gamma_2, \gamma_3, \gamma_4, \gamma_5, \gamma_6, \gamma_7]$
$E_8$	8	$[\gamma_1, \gamma_2, \gamma_3, \gamma_4, \gamma_5, \gamma_6, \gamma_7, \gamma_8]$

We shall consider here representations of the classical compact algebras,  $su(n)$ ,  $so(n)$ ,  $sp(n)$ . There are two types of representations: tensor representations and spinor representations.

## 4.3 Irreducible Tensors

### 4.3.1 Irreducible Tensors with Respect to $GL(n)$

In order to illustrate the notion of irreducible tensors we consider here two vectors  $\mathbf{x} \equiv (x_1, \dots, x_n)$  and  $\mathbf{y} \equiv (y_1, \dots, y_n)$ . Under  $GL(n)$  they transform as

$$x'_i = \sum_k a_{ik} x_k \quad ; \quad y'_i = \sum_k a_{ik} y_k \quad ; \quad (i, k = 1, \dots, n) . \quad (4.2)$$

Consider now the Kronecker product of the two vectors,  $F_{ij} = x_i y_j$ . This product has  $n \times n$  components and transforms as a second rank tensor with respect to  $GL(n)$

$$F'_{ij} = \sum_{k,l} a_{ik} a_{jl} F_{kl} . \quad (4.3)$$

Irreducible tensors with respect to  $GL(n)$  are obtained by taking the symmetric and antisymmetric parts

$$\begin{aligned} S_{ik} &= S_{ki} = \frac{1}{2} (F_{ik} + F_{ki}) , \\ A_{ik} &= -A_{ki} = \frac{1}{2} (F_{ik} - F_{ki}) . \end{aligned} \quad (4.4)$$

The symmetric tensor has  $n(n+1)/2$  components, while the antisymmetric tensor has  $n(n-1)/2$  components. In general, reducibility with respect to  $GL(n)$  (and  $gl(n)$ ) means to classify tensors according to their symmetry under interchange of indices. Young devised a procedure how to find the possible symmetry types of a tensor of rank  $t$  in a  $n$  dimensional space.

(i) Partition  $t$  into  $n$  integers

$$t = \lambda_1 + \dots + \lambda_n \quad (4.5)$$

with

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0 . \quad (4.6)$$

(ii) To each partition there correspond a graph (or tableau)

$$\begin{array}{cccc} \lambda_1 & \square & \square & \dots & \square \\ \lambda_2 & \square & \dots & \square & \\ \dots & \dots & & & \\ \lambda_n & \square & & & \end{array} \quad (4.7)$$

The tensor is symmetric under interchange of the rows and antisymmetric under interchange of columns. The tableau is often denoted by  $[\lambda_1, \dots, \lambda_n]$  and zeros are deleted.

**Example 1.** Consider a second rank tensor  $t = 2$  in a three dimensional space,  $n = 3$ .

The partitions are  $[2], [1, 1]$  with Young diagrams

$$\begin{aligned} \square\square &\equiv [2] && \text{Symmetric tensor} \\ \begin{array}{c} \square \\ \square \end{array} &\equiv [1, 1] && \text{Antisymmetric tensor} \end{aligned} \tag{4.8}$$

### 4.3.2 Irreducible Tensors with Respect to $SU(n)$

The irreducible representations of  $GL(n)$  remain irreducible when we go to  $U(n)$ . If we go to the unimodular groups  $SU(n)$ , the representations corresponding to the patterns  $[\lambda_1, \lambda_2, \dots, \lambda_n]$  and  $[\lambda_1 + s, \lambda_2 + s, \dots, \lambda_n + s]$ ,  $s = \text{integer}$ , are equivalent. Thus, for  $SU(n)$ , we need to consider only patterns with one less row  $[\lambda_1 - \lambda_n, \lambda_2 - \lambda_n, \dots, \lambda_{n-1} - \lambda_n]$  obtained from the pattern  $[\lambda_1, \lambda_2, \dots, \lambda_n]$  by subtracting the last integer  $\lambda_n$ .

### 4.3.3 Irreducible Tensors with Respect to $O(n)$ . Contractions

When we go from  $U(n)$  to  $O(n)$ , the representations in terms of tensors of a given symmetry are no longer irreducible. There is a new operation, called contraction, which commutes with orthogonal transformations. For  $O(n)$ , the elements of the transformation matrix  $a_{ij}$  satisfy

$$\sum_i a_{ij} a_{ik} = \delta_{jk} . \tag{4.9}$$

Contraction of a second rank tensor  $F_{ij}$  gives

$$\tau = \sum_i F_{ii} . \tag{4.10}$$

In general, contraction of a rank- $t$  tensor, gives a tensor of rank  $t - 2$ . Any tensor can be decomposed into a traceless part, plus the rest.

**Example 2.** Decomposition of a rank-2 tensor in  $n = 3$  dimensions

	$GL(3)$	$O(3)$	$\dim$	$L$	
		$S_{ik} - \frac{1}{3}S_{ii}\delta_{ik} = \Sigma_{ik}$	5	2	
	$S_{ik}$				
	$F_{ik}$	$\frac{1}{3}S_{ii}\delta_{ik} = \tau_{ik}$	1	0	
			3	1	
	$A_{ik}$				

(4.11)

The decomposition is thus

$$F_{ik} = A_{ik} + \Sigma_{ik} + \tau_{ik} . \tag{4.12}$$

The number of components of the tensors  $A, \Sigma, \tau$  is shown in the column labeled  $\dim$ .

## 4.4 Tensor Representations of Classical Compact Algebras

### 4.4.1 Unitary Algebras $u(n)$

Irreducible representations of  $u(n)$  are characterized by  $n$  integers, satisfying the conditions

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0 . \tag{4.13}$$

A graphical representation is provided by the Young tableau introduced previously for  $U(n)$

$$\begin{array}{c} \lambda_1 \square \square \dots \square \\ \lambda_2 \square \square \dots \square \\ \dots \\ \lambda_n \square \end{array} , \tag{4.14}$$

also written as  $[\lambda_1, \lambda_2, \dots, \lambda_n]$ .

### 4.4.2 Special Unitary Algebras $su(n)$

Irreducible representations of  $su(n)$  are characterized by  $n - 1$  integers, satisfying the relations

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_{n-1} \geq 0 . \tag{4.15}$$

Because of the special condition,  $S$ , some representations become equivalent.

**Equivalence relation 1** Start from the representations of  $u(n)$  and subtract the last integer  $[\lambda_1, \lambda_2, \dots, \lambda_n] \equiv [\lambda_1 - \lambda_n, \lambda_2 - \lambda_n, \dots, \lambda_{n-1} - \lambda_n, 0]$ .

This equivalence relation was already quoted in Sect. 4.3.2.

**Equivalence relation 2** Start from the representations of  $u(n)$  and use  $[\lambda_1, \lambda_2, \dots, \lambda_n] \equiv [\lambda_1 - \lambda_n, \lambda_1 - \lambda_{n-1}, \dots, \lambda_1 - \lambda_2, 0]$ .

This equivalence relation, when written at the level of  $su(n)$  is  $[\lambda_1, \lambda_2, \dots, \lambda_n] \equiv [\lambda_1, \lambda_1 - \lambda_{n-1}, \dots, \lambda_1 - \lambda_2]$  and is sometimes called particle-hole conjugation.

The equivalence relations 1 and 2 are used extensively.

**Example 2.** The first equivalence relation for  $su(3)$  gives

$$[4, 3, 1] \equiv [3, 2] . \tag{4.16}$$

**Example 3.** The second equivalence relation for  $su(3)$  gives

$$[3, 2] \equiv [3, 1] . \tag{4.17}$$

#### 4.4.3 Orthogonal Algebras $so(n)$ , $n = \text{Odd}$

The irreducible representations are labeled by  $\nu = (n - 1)/2$  integers, satisfying

$$\mu_1 \geq \mu_2 \geq \dots \geq \mu_\nu \geq 0 . \tag{4.18}$$

#### 4.4.4 Orthogonal Algebras $so(n)$ , $n = \text{Even}$

The irreducible representations are labeled by  $\nu = n/2$  integers, satisfying

$$\mu_1 \geq \mu_2 \geq \dots \geq |\mu_\nu| \geq 0 . \tag{4.19}$$

The last integer  $\mu_\nu$  can be here positive, negative or zero. If  $\mu_\nu \neq 0$ , there are two irreducible representations, called mirror conjugate, with  $\mu_\nu = \pm |\mu_\nu|$ .

#### 4.4.5 Symplectic Algebras $sp(n)$ , $n = \text{Even}$

The irreducible representations are labeled by  $\nu = n/2$  integers, satisfying

$$\mu_1 \geq \mu_2 \geq \dots \geq \mu_\nu \geq 0 . \tag{4.20}$$

**Example 4.** Unitary algebras

$$\begin{aligned} u(2) & \quad [\lambda_1, \lambda_2] , \quad \lambda_1 \geq \lambda_2 \geq 0 \\ u(3) & \quad [\lambda_1, \lambda_2, \lambda_3] , \quad \lambda_1 \geq \lambda_2 \geq \lambda_3 \geq 0 \end{aligned} \tag{4.21}$$

**Example 5.** Special unitary algebras

$$\begin{aligned} su(2) & \quad [\lambda_1 - \lambda_2] = [f_1] , \quad f_1 \geq 0 \\ su(3) & \quad [\lambda_1 - \lambda_3, \lambda_2 - \lambda_3] \equiv [f_1, f_2] , \quad f_1 \geq f_2 \geq 0 \end{aligned} \tag{4.22}$$

**Example 6.** *Orthogonal algebras*

$$\begin{aligned}
 so(2) \quad & [\mu_1] \equiv M, \quad | \mu_1 | \geq 0 \\
 so(3) \quad & [\mu_1] \equiv L, \quad \mu_1 \geq 0 \\
 so(4) \quad & [\mu_1, \mu_2] \equiv (\omega_1, \omega_2); \mu_1 \geq | \mu_2 | \geq 0 . \\
 so(5) \quad & [\mu_1, \mu_2] \equiv (\tau_1, \tau_2); \mu_1 \geq \mu_2 \geq 0
 \end{aligned} \tag{4.23}$$

Often, in applications, the abstract labels are replaced by letters related to their physical interpretation, especially for orthogonal algebras and groups. In Example 6, the letter  $M$  ( $z$ -component of the angular momentum) is used to denote the representations of  $so(2)$ , and the letter  $L$  (angular momentum) to denote the representations of  $so(3)$ .

### 4.5 Spinor Representations

As discussed in Chap. 3, the group  $SO(3)$  is doubly connected. It turns out that all orthogonal groups in odd number of dimensions  $SO(2\nu + 1)$  are doubly connected, while the orthogonal groups in even number of dimensions,  $SO(2n)$ , are four-fold connected. As a result, orthogonal groups and algebras have another type of representations, called spinor representations, characterized by half-integer labels. (The additional two-fold connecteness of  $SO(2\nu)$  produces mirror conjugate representations already discussed in the previous section.) [A.O. Barut and R. Raczka, *Theory of Group Representations and Applications*, World Scientific, Singapore, 1986].

#### 4.5.1 Orthogonal Algebras $so(n)$ , $n = \text{Odd}$

Spinor representations of  $so(n)$ ,  $n = \text{odd}$ , are characterized by  $\nu = (n - 1)/2$  half-integers

$$\mu_1 \geq \mu_2 \geq \dots \geq \mu_\nu \geq \frac{1}{2} . \tag{4.24}$$

#### 4.5.2 Orthogonal Algebras $so(n)$ , $n = \text{Even}$

Spinor representations of  $so(n)$ ,  $n = \text{odd}$ , are characterized by  $\nu = n/2$  half-integers

$$\mu_1 \geq \mu_2 \geq \dots \geq | \mu_\nu | \geq \frac{1}{2} . \tag{4.25}$$

As in the case of tensor representations, when  $n = \text{even}$ , there are two irreducible representations, called mirror conjugate, with  $\mu_\nu = \pm | \mu_\nu |$ .

When spinor representations are included the algebras and groups are denoted  $spin(n)$  and  $Spin(n)$  respectively.

**Example 7.** *Spinor representations*

$$\begin{aligned}
 spin(2) \quad & [\mu_1] \equiv M_J = \pm \frac{1}{2}, \pm \frac{3}{2}, \dots \\
 spin(3) \quad & [\mu_1] \equiv J = \frac{1}{2}, \frac{3}{2}, \dots
 \end{aligned} \tag{4.26}$$



## 4.6 Fundamental Representations

Any irreducible representation can be written as

$$[\lambda] = \sum_{i=1}^l f_i [\lambda^i], \tag{4.27}$$

where the  $f_i$ 's are non-negative integers and the  $[\lambda^i]$ 's are called fundamental representations. Here  $[\lambda]$  is a short-hand notation for the Young tableau characterizing the representation and  $[\lambda^i]$  is a short-hand for the Young tableau characterizing the fundamental representations. The index  $i$  runs from 1 to the values given in Table 4.1 under "number of labels". The fundamental representations of the classical Lie algebras are listed below.

### 4.6.1 Unitary Algebras

There are  $n$  fundamental representations here

$$\begin{aligned} &u(n) \\ &[1, 0, 0, \dots, 0] \\ &[1, 1, 0, \dots, 0] \ . \\ &\dots \\ &[1, 1, 1, \dots, 1] \end{aligned} \tag{4.28}$$

### 4.6.2 Special Unitary Algebras

There are  $n - 1$  fundamental representations

$$\begin{aligned} &su(n) \\ &[1, 0, 0, \dots, 0] \\ &[1, 1, 0, \dots, 0] \ . \\ &\dots \\ &[1, 1, 1, \dots, 1] \end{aligned} \tag{4.29}$$

### 4.6.3 Orthogonal Algebras, $n = \text{Odd}$

There are  $\nu = (n - 1)/2$  fundamental representations, one of which is a spinor representation

$$\begin{aligned} &spin(n), n = \text{odd} \\ &[1, 0, \dots, 0] \\ &[1, 1, \dots, 0] \ . \\ &\dots \\ &[1, 1, \dots, 1, 0] \\ &[\frac{1}{2}, \frac{1}{2}, \dots, \frac{1}{2}, \frac{1}{2}] \end{aligned} \tag{4.30}$$

### 4.6.4 Orthogonal Algebras, $n = \text{Even}$

There are  $\nu = n/2$  fundamental representations, two of which are spinor representations

$$\begin{aligned}
 & \text{spin}(n), n = \text{even} \\
 & \begin{aligned} & [1, 0, \dots, 0] \\ & [1, 1, \dots, 0] \\ & \dots \end{aligned} \\
 & \begin{aligned} & [1, 1, \dots, 1, 0, 0] \\ & [\frac{1}{2}, \frac{1}{2}, \dots, \frac{1}{2}, \frac{1}{2}] \\ & [\frac{1}{2}, \frac{1}{2}, \dots, \frac{1}{2}, -\frac{1}{2}] \end{aligned} \end{aligned} \tag{4.31}$$

### 4.6.5 Symplectic Algebras

There are  $\nu = n/2$  fundamental representations.

$$\begin{aligned}
 & sp(n), n = \text{even} \\
 & \begin{aligned} & [1, 0, \dots, 0] \\ & [1, 1, \dots, 0] \\ & \dots \\ & [1, 1, \dots, 1] \end{aligned} \end{aligned} \tag{4.32}$$

## 4.7 Chains of Algebras

For applications, it is necessary to characterize uniquely the basis, in other words to provide a complete set of quantum numbers (labels). This is done by introducing a chain of algebras,

$$\left| \begin{array}{cccc} g & \supset & g' & \supset & g'' & \supset & \dots \\ \downarrow & & \downarrow & & \downarrow & & \\ [\lambda] & & [\lambda'] & & [\lambda''] & & \end{array} \right\rangle \tag{4.33}$$

Since in applications in quantum mechanics, the representations are interpreted as quantum mechanical states, a notation often used is that introduced by Dirac, called bra-ket notation. A ket is denoted by  $| \rangle$  and a bra by  $\langle |$ .

A crucial problem of representation theory is to find the irreducible representations of an algebra  $g'$  contained in a given representation of  $g$  (often called the branching problem).

## 4.8 Canonical Chains

The branching problem was solved completely by Gel'fand and Cetlin in a series of articles in the 1950's, for a particular chain of algebras, called the canonical chain of unitary and orthogonal algebras. [I.M. Gel'fand and

M.L. Cetlin, *Finite-Dimensional Representations of a Group of Unimodular Matrices*, Dokl. Akad. Nauk SSSR 71, 8–25, 825–828 (1950); *Finite-Dimensional Representations of Groups of Orthogonal Matrices*, ibid. 71, 1017–1020 (1950).]

### 4.8.1 Unitary Algebras

The canonical chain is

$$u(n) \supset u(n-1) \supset u(n-2) \supset \dots \supset u(1). \tag{4.34}$$

The labels (quantum numbers) are conveniently arranged into a pattern called Gel'fand pattern

$$\begin{array}{ccccccc}
 \lambda_{1,n} & & \lambda_{2,n} & & & \lambda_{n-1,n} & \lambda_{n,n} \\
 & \lambda_{1,n-1} & & \dots & & \dots & \lambda_{n-1,n-1} \\
 & & \dots & & \dots & & \\
 & & & \lambda_{1,2} & \lambda_{2,2} & & \\
 & & & & \lambda_{1,1} & & 
 \end{array} . \tag{4.35}$$

The entries in this pattern are  $\lambda_{i,j}$  where  $i$  labels the entries in the Young tableau, and  $j$  the algebra in the chain. For example,  $\lambda_{n,n}$  is the  $n$ -th entry in the Young tableau of  $u(n)$ .

The solution to the branching problem is that the labels must satisfy the inequalities

$$\begin{aligned}
 \lambda_{1,n} &\geq \lambda_{1,n-1} \geq \lambda_{2,n} \geq \dots \geq \lambda_{n,n} \geq 0 \\
 &\dots \\
 \lambda_{1,2} &\geq \lambda_{1,1} \geq \lambda_{2,2} .
 \end{aligned} \tag{4.36}$$

These inequalities hold for any two rows in (4.35) and are often called triangular inequalities.

**Example 8.** *Representations of  $u(4)$*

The complete basis is labelled by

$$\begin{array}{cccc}
 \lambda_{1,4} & \lambda_{2,4} & \lambda_{3,4} & \lambda_{4,4} \\
 & \lambda_{1,3} & \lambda_{2,3} & \lambda_{3,3} \\
 & & \lambda_{1,2} & \lambda_{2,2} \\
 & & & \lambda_{1,1}
 \end{array} \tag{4.37}$$

In the applications discussed in Chap. 7, one needs the branching of the totally symmetric representation  $[N]$  of  $u(4)$  with ket  $|N_1; N_2; N_3; N_4\rangle$  and pattern

$$\begin{array}{cccc}
 N_4 & 0 & 0 & 0 \\
 & N_3 & 0 & 0 \\
 & & N_2 & 0 \\
 & & & N_1
 \end{array} . \tag{4.38}$$

Use of Gel'fand inequalities gives  $N_3 = N_4, N_4 - 1, \dots, 0; N_2 = N_3, N_3 - 1, \dots, 0; N_1 = N_2, N_2 - 1, \dots, 0$ .

### 4.8.2 Orthogonal Algebras

The canonical chain is

$$so(n) \supset so(n-1) \supset so(n-2) \supset \dots \supset so(2). \quad (4.39)$$

For  $n = 2k + 2 = \text{even}$ , the Gel'fand pattern is

$$\begin{array}{ccccccc}
 \mu_{1,2k+1} & & \mu_{2,2k+1} & & \cdots & & \mu_{k,2k+1} & & \mu_{k+1,2k+1} \\
 & & \mu_{1,2k} & & & & & & \mu_{k,2k} \\
 & & \mu_{1,2k-1} & & & & & & \mu_{k,2k-1} \\
 & & & \cdots & & & & & \\
 & & & & \mu_{1,4} & & \mu_{2,4} & & \\
 & & & & \mu_{1,3} & & \mu_{2,3} & & \\
 & & & & & & \mu_{1,2} & & \\
 & & & & & & \mu_{1,1} & & 
 \end{array} \quad (4.40)$$

There is an alternation between even and odd algebras, ending with  $so(5) \supset so(4) \supset so(3) \supset so(2)$ . The triangular inequalities are:

$$\begin{aligned}
 \mu_{1,2k+1} &\geq \mu_{1,2k} \geq \mu_{2,2k+1} \geq \dots \geq \mu_{k,2k} \geq | \mu_{k+1,2k+1} | \\
 \mu_{1,2k} &\geq \mu_{1,2k-1} \geq \mu_{2,2k} \geq \dots \geq \mu_{k,2k} \geq | \mu_{k,2k-1} | \\
 \mu_{1,2k-1} &\geq \mu_{1,2k-2} \geq \dots \geq \mu_{k-1,2k-2} \geq | \mu_{k,2k-1} | \cdot
 \end{aligned} \quad (4.41)$$

For  $n = 2k + 1 = \text{odd}$ , the Gel'fand pattern is

$$\begin{array}{ccccccc}
 & & \mu_{1,2k} & & & & \mu_{k,2k} \\
 & & \mu_{1,2k-1} & & & & \mu_{k,2k-1} \\
 & & & \mu_{1,2k-2} & & & \mu_{k-1,2k-2} \\
 & & & \cdots & & & \cdots \\
 & & & & \mu_{1,4} & & \mu_{2,4} \\
 & & & & \mu_{1,3} & & \mu_{2,3} \\
 & & & & & & \mu_{1,2} \\
 & & & & & & \mu_{1,1}
 \end{array} \quad (4.42)$$

There is alternation between odd and even algebras, still ending with  $so(5) \supset so(4) \supset so(3) \supset so(2)$  with inequalities

$$\begin{aligned}
 \mu_{1,2k} &\geq \mu_{1,2k-1} \geq \dots \geq \mu_{k,2k} \geq | \mu_{k,2k-1} | \\
 \mu_{1,2k-1} &\geq \mu_{1,2k-2} \geq \dots \geq \mu_{k-1,2k-2} \geq | \mu_{k,2k-1} | \cdot
 \end{aligned} \quad (4.43)$$

#### Example 9. Branching of $so(4)$

For the ket

$$\left\langle \begin{array}{ccc} so(4) & \supset & so(3) \supset so(2) \\ \downarrow & & \downarrow \quad \downarrow \\ [\mu_1, \mu_2] & & J \quad M \end{array} \right\rangle \tag{4.44}$$

the Gel'fand pattern is

$$\begin{array}{ccc} \mu_1 & & \mu_2 \\ & J & \\ & M & \end{array} \tag{4.45}$$

with branching

$$\mu_1 \geq J \geq |\mu_2|, \quad +J \geq M \geq -J. \tag{4.46}$$

### 4.9 Isomorphisms of Spinor Algebras

Because of the isomorphisms discussed in Chap. 2, one has

$$\begin{aligned} spin(2) &\sim u(1) \\ spin(3) &\sim su(2) \\ spin(4) &\sim su(2) \oplus su(2) \\ spin(5) &\sim sp(4) \\ spin(6) &\sim su(4) \end{aligned} \tag{4.47}$$

After *spin*(6), the spinor algebras are no longer isomorphic to other (non-orthogonal) classical Lie algebras.

It is of interest to find the relation between the quantum numbers labelling the representations of two isomorphic algebras.

**Example 10.** *The case  $spin(3) \sim su(2)$*

$$\begin{array}{ccc} spin(3) & \sim & su(2) \\ \downarrow & & \downarrow \\ J & & \lambda \end{array} . \tag{4.48}$$

The relation in this case is simply  $J = \frac{\lambda}{2}$ ,

$$su(2) : \overbrace{\square \square \dots \square}^{\lambda} \equiv spin(3) : J = \frac{\lambda}{2} \tag{4.49}$$

**Example 11.** *The case  $spin(6) \sim su(4)$*

$$\begin{array}{ccc} spin(6) & \sim & su(4) \\ \downarrow & & \downarrow \\ [\sigma_1, \sigma_2, \sigma_3] & & [\lambda_1, \lambda_2, \lambda_3] \\ \sigma_1 \geq \sigma_2 \geq |\sigma_3| \geq 0 & & \lambda_1 \geq \lambda_2 \geq \lambda_3 \geq 0 \end{array} \tag{4.50}$$

The relation in this case is

$$\lambda_1 = \sigma_1 + \sigma_2, \quad \lambda_2 = \sigma_1 - \sigma_3, \quad \lambda_3 = \sigma_2 - \sigma_3. \quad (4.51)$$

The correspondence between the fundamental representations is

$$\begin{array}{c}
 \hline
 \begin{array}{cc}
 \textit{spin}(6) & \textit{su}(4)
 \end{array} \\
 \hline
 \begin{array}{ccc}
 [\frac{1}{2}, \frac{1}{2}, \frac{1}{2}] & [1, 0, 0] & \square \\
 \hline
 [\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}] & [1, 1, 1] & \square \\
 \hline
 [1, 0, 0] & [1, 1, 0] & \square
 \end{array}
 \end{array}
 \cdot \quad (4.52)$$

### 4.10 Nomenclature for $u(n)$

In physics textbooks, the representations of  $u(n)$  are often referred according to the particles they describe. The totally symmetric representations

$$\overbrace{\square \square \dots \square}^N \quad [N, 0, 0, \dots, 0] = [N, \dot{0}] \equiv [N] \quad (4.53)$$

describe bosons and are often referred to as bosonic. The notation  $[N, \dot{0}]$  is rarely used. The totally antisymmetric representations

$$\begin{array}{c}
 \square \\
 \square \\
 \dots \\
 \square
 \end{array}
 \quad [1, 1, \dots, 1] \equiv [\dot{1}] \quad (4.54)$$

describe fermions and are often referred to as fermionic. The representations with mixed symmetry

$$\begin{array}{c}
 \square \square \dots \square \\
 \square \dots \square \\
 \vdots \\
 \square
 \end{array} \quad (4.55)$$

describe particles with internal degrees of freedom.

### 4.11 Dimensions of the Representations

The dimensions of the representations are often used to check the branching rules. A general formula was provided by Weyl and is called the Weyl formula.

### 4.11.1 Dimensions of the Representations of $u(n)$

The dimension of the representation  $[\lambda] \equiv [\lambda_1, \lambda_2, \dots, \lambda_n]$  of  $u(n)$  is given by the formula

$$\dim[\lambda] = \prod_{i < j} \frac{(\ell_i - \ell_j)}{(\ell_i^0 - \ell_j^0)} \quad \ell_j^0 = n - j, \quad \ell_j = \lambda_j + n - j, \quad (4.56)$$

with  $i, j = 1, \dots, n$ .

**Example 12.** *Dimensions of the representations of  $u(3)$*

For  $u(3)$ ,

$$\dim[\lambda] = \frac{(\ell_1 - \ell_2)(\ell_1 - \ell_3)(\ell_2 - \ell_3)}{(\ell_1^0 - \ell_2^0)(\ell_1^0 - \ell_3^0)(\ell_2^0 - \ell_3^0)}. \quad (4.57)$$

From (4.56)

$$\dim[\lambda_1, \lambda_2, \lambda_3] = \frac{1}{2} (\lambda_1 - \lambda_2 + 1) (\lambda_1 - \lambda_3 + 2) (\lambda_2 - \lambda_3 + 1). \quad (4.58)$$

For example, the dimension of the representation  $[2, 1, 0]$  is

$$\dim[2, 1, 0] = 8 \quad (4.59)$$

It has become customary in particle physics to denote the representation with its dimension. This notation is ambiguous, as often there are different representations with the same dimension. The Gel'fand patterns of the basis states of the representation  $[2, 1]$  of  $u(3)$  and  $su(3)$  are

$$\begin{aligned} & \left| \begin{array}{ccc} 2 & 1 & 0 \\ & 2 & 1 \\ & & 2 \end{array} \right\rangle \left| \begin{array}{ccc} 2 & 1 & 0 \\ & 2 & 1 \\ & & 1 \end{array} \right\rangle \\ & \left| \begin{array}{ccc} 2 & 1 & 0 \\ & 2 & 0 \\ & & 2 \end{array} \right\rangle \left| \begin{array}{ccc} 2 & 1 & 0 \\ & 2 & 0 \\ & & 1 \end{array} \right\rangle \left| \begin{array}{ccc} 2 & 1 & 0 \\ & 2 & 0 \\ & & 0 \end{array} \right\rangle \\ & \left| \begin{array}{ccc} 2 & 1 & 0 \\ & 1 & 1 \\ & & 1 \end{array} \right\rangle \\ & \left| \begin{array}{ccc} 2 & 1 & 0 \\ & 1 & 0 \\ & & 1 \end{array} \right\rangle \left| \begin{array}{ccc} 2 & 1 & 0 \\ & 1 & 0 \\ & & 0 \end{array} \right\rangle \end{aligned} \quad (4.60)$$

In this application, the basis states correspond to particles. The eight states of the representation 8 for baryons correspond to particles called  $p, n, \Sigma^+, \Sigma^0, \Sigma^-, \Xi^0, \Xi^-$ .

### 4.11.2 Dimensions of the Representations of $su(n)$

The dimensions of the representations of  $su(n)$  can be obtained from those of  $u(n)$  by equivalences. Thus, for the representation  $[\lambda_1, \lambda_2, \dots, \lambda_{n-1}]$ , formula (4.56) applies with  $\lambda_n = 0$ .

### 4.11.3 Dimensions of the Representations of $A_n \equiv su(n+1)$

An alternative formula for the dimensions of the representations of  $su(n)$ , which makes the connection with those of the orthogonal and symplectic algebras given below clear, can be written for the Cartan algebras  $A_n \equiv su(n+1)$ . This formula can be simply derived from (4.54).

One first constructs the quantities

$$g_i = n - i + 1, \quad m_i = \lambda_i + g_i \quad (i = 1, \dots, n). \quad (4.61)$$

The dimensions of the representations of  $A_n$  are

$$\dim[\lambda] = \prod_i \left( \frac{m_i}{g_i} \right) \prod_{i < j} \left( \frac{m_i - m_j}{g_i - g_j} \right). \quad (4.62)$$

### 4.11.4 Dimensions of the Representations of $B_n \equiv so(2n+1)$

One constructs the quantities

$$g_i = n - i + \frac{1}{2}, \quad m_i = \mu_i + g_i \quad (i = 1, \dots, n). \quad (4.63)$$

The dimensions of the representations of  $B_n$  are

$$\dim[\mu] = \prod_i \left( \frac{m_i}{g_i} \right) \prod_{i < j} \left( \frac{m_i - m_j}{g_i - g_j} \right) \prod_{i < j} \left( \frac{m_i + m_j}{g_i + g_j} \right). \quad (4.64)$$

#### Example 13. Dimensions of the representations of $so(3)$ and $so(5)$

The dimensions of the representations of  $so(3)$  and  $so(5)$  are given by

$$\begin{aligned} so(3) \quad & \dim[\mu_1] = (2\mu_1 + 1) \\ so(5) \quad & \dim[\mu_1, \mu_2] = \frac{1}{6} (\mu_1 - \mu_2 + 1) (\mu_1 + \mu_2 + 2) (2\mu_1 + 3) (2\mu_2 + 1). \end{aligned} \quad (4.65)$$



### 4.11.5 Dimensions of the Representations of $C_n \equiv sp(2n)$

Here

$$g_i = n - i + 1, \quad m_i = \mu_i + g_i \quad (i = 1, \dots, n). \quad (4.66)$$

The dimensions of the representations of  $C_n$  is

$$\dim [\mu] = \prod_i \binom{m_i}{g_i} \prod_{i < j} \binom{m_i - m_j}{g_i - g_j} \prod_{i < j} \binom{m_i + m_j}{g_i + g_j}. \quad (4.67)$$

**Example 14.** *Dimensions of the representations of  $sp(4)$*

The dimensions of the representations of  $sp(4)$  are given by

$$sp(4) \dim [\mu_1, \mu_2] = \frac{1}{6} (\mu_1 - \mu_2 + 1) (\mu_1 + \mu_2 + 3) (\mu_1 + 2) (\mu_2 + 1). \quad (4.68)$$

### 4.11.6 Dimensions of the Representations of $D_n \equiv so(2n)$

Here

$$g_i = n - 1, \quad m_i = \mu_i + g_i \quad (i = 1, \dots, n). \quad (4.69)$$

The dimensions of the representations of  $D_n$  are

$$\dim [\mu] = \prod_{i < j} \binom{m_i - m_j}{g_i - g_j} \prod_{i < j} \binom{m_i + m_j}{g_i + g_j}. \quad (4.70)$$

**Example 15.** *Dimension of the representations of  $so(4)$*

The dimensions of the representations of  $so(4)$  are

$$so(4) \dim [\mu_1, \mu_2] = (\mu_1 - \mu_2 + 1) (\mu_1 + \mu_2 + 1). \quad (4.71)$$

## 4.12 Action of the Elements of $g$ on the Basis $\mathcal{B}$

The action of the elements of  $g$  on the basis is of great interest in physics. We consider here the case of  $u(n)$ . The elements of  $u(n)$ , when written in the double index notation of Chap. 2, satisfy the commutation relations

$$[E_{ij}, E_{kl}] = \delta_{jk} E_{il} - \delta_{il} E_{kj} \quad i, j, k, l = 1, \dots, n. \quad (4.72)$$

Consider only  $E_{k,k}, E_{k,k-1}$  and  $E_{k-1,k}$ , since the action of the others can be obtained from the commutators of these. The action is

$$\begin{aligned}
 E_{k,k} | \lambda \rangle &= (r_k - r_{k-1}) | \lambda \rangle \\
 E_{k,k-1} | \lambda \rangle &= \sum_{j=1}^{k-1} a_{k-1}^j | \lambda' \rangle \\
 E_{k-1,k} | \lambda \rangle &= \sum_{j=1}^{k-1} b_{k-1}^j | \lambda'' \rangle
 \end{aligned}
 \tag{4.73}$$

where  $E_{k,k}$  is called the diagonal element (or operator),  $E_{k,k-1}$  the lowering element (or operator),  $E_{k-1,k}$  the raising element (or operator). In this equation,  $| \lambda \rangle$  denotes a generic representation  $| \lambda_1, \lambda_2, \dots, \lambda_n \rangle$ . The coefficients  $r_k$  are given by

$$r_0 = 0, \quad r_k = \sum_{j=1}^k \lambda_{j,k}, \quad k = 1, \dots, n, \tag{4.74}$$

while the coefficients  $a$  and  $b$  are given by

$$\begin{aligned}
 a_{k-1}^j &= \left[ \frac{\prod_{i=1}^k (\lambda_{i,k} - \lambda_{j,k-1} - i + j + 1) \prod_{i=1}^{k-2} (\lambda_{i,k-2} - \lambda_{j,k-1} - i + j)}{\prod_{\substack{i \neq j \\ i=1}}^{k-1} (\lambda_{i,k-1} - \lambda_{j,k-1} - i + j + 1) \prod_{\substack{i \neq j \\ i=1}}^{k-1} (\lambda_{i,k-1} - \lambda_{j,k-1} - i + j)} \right]^{1/2} \\
 b_{k-1}^j &= \left[ \frac{\prod_{i=1}^k (\lambda_{i,k} - \lambda_{j,k-1} - i + j) \prod_{i=1}^{k-2} (\lambda_{i,k-2} - \lambda_{j,k-1} - i + j - 1)}{\prod_{\substack{i \neq j \\ i=1}}^{k-1} (\lambda_{i,k-1} - \lambda_{j,k-1} - i + j) \prod_{\substack{i \neq j \\ i=1}}^{k-1} (\lambda_{i,k-1} - \lambda_{j,k-1} - i + j - 1)} \right]^{1/2}
 \end{aligned}
 \tag{4.75}$$

The representation  $| \lambda' \rangle$  is obtained from  $| \lambda \rangle$  by replacing  $\lambda_{j, k-1}$  by  $\lambda_{j, k-1} - 1$ . (This is the reason why the operator  $E_{k,k-1}$  is called lowering operator).  $| \lambda'' \rangle$  is obtained from  $| \lambda \rangle$  by replacing  $\lambda_{j, k-1}$  by  $\lambda_{j, k-1} + 1$ . ( $E_{k-1,k}$  is called raising operator).

These results were obtained by Baird and Biederharn in the 1960's. [G.E. Baird and L.C. Biedenarn, *On the Representations of Semisimple Lie Groups-II*, J. Math. Phys. 4, 1449 (1963).]

**Example 16.** *The algebra  $u(2)$*

The states are

$$\left| \begin{matrix} \lambda_{1,2} & \lambda_{2,2} \\ & \lambda_{1,1} \end{matrix} \right\rangle. \tag{4.76}$$

The algebra contains four elements  $E_{1,1}, E_{1,2}, E_{2,1}, E_{2,2}$ . The action of the elements on the basis is

$$\begin{aligned}
 E_{1,1} \left| \begin{array}{c} \lambda_{1,2} \quad \lambda_{2,2} \\ \lambda_{1,1} \end{array} \right\rangle &= \lambda_{1,1} \left| \begin{array}{c} \lambda_{1,2} \quad \lambda_{2,2} \\ \lambda_{1,1} \end{array} \right\rangle \\
 E_{2,2} \left| \begin{array}{c} \lambda_{1,2} \quad \lambda_{2,2} \\ \lambda_{1,1} \end{array} \right\rangle &= (\lambda_{1,2} + \lambda_{2,2} - \lambda_{1,1}) \left| \begin{array}{c} \lambda_{1,2} \quad \lambda_{2,2} \\ \lambda_{1,1} \end{array} \right\rangle \\
 E_{2,1} \left| \begin{array}{c} \lambda_{1,2} \quad \lambda_{2,2} \\ \lambda_{1,1} \end{array} \right\rangle &= [(\lambda_{1,2} - \lambda_{1,1} + 1) (\lambda_{1,1} - \lambda_{2,2})]^{1/2} \left| \begin{array}{c} \lambda_{1,2} \quad \lambda_{2,2} \\ \lambda_{1,1} - 1 \end{array} \right\rangle \\
 E_{1,2} \left| \begin{array}{c} \lambda_{1,2} \quad \lambda_{2,2} \\ \lambda_{1,1} \end{array} \right\rangle &= [(\lambda_{1,1} - \lambda_{1,2}) (\lambda_{2,2} - \lambda_{1,1} - 1)]^{1/2} \left| \begin{array}{c} \lambda_{1,2} \quad \lambda_{2,2} \\ \lambda_{1,1} + 1 \end{array} \right\rangle
 \end{aligned} \tag{4.77}$$

**Example 17.** *The algebra  $su(2)$*

In Gel'fand notation the basis states are

$$\left| \begin{array}{c} \lambda_{1,2} = 2J \\ \lambda_{1,1} = M + J \\ \lambda_{2,2} = 0 \end{array} \right\rangle \tag{4.78}$$

In the usual notation in quantum mechanics the basis states are written as  $|J, M\rangle$ . The action of the elements on the basis is

$$\begin{aligned}
 E_{1,1} |J, M\rangle &= (M + J) |J, M\rangle \\
 E_{2,2} |J, M\rangle &= (-M + J) |J, M\rangle .
 \end{aligned} \tag{4.79}$$

From these one obtains

$$\frac{1}{2} (E_{1,1} - E_{2,2}) |J, M\rangle = M |J, M\rangle \tag{4.80}$$

The action of the raising and lowering operators is

$$\begin{aligned}
 E_{2,1} |J, M\rangle &= [J(J+1) - M(M-1)]^{1/2} |J, M-1\rangle \\
 E_{1,2} |J, M\rangle &= [J(J+1) - M(M+1)]^{1/2} |J, M+1\rangle .
 \end{aligned} \tag{4.81}$$

In quantum mechanics textbooks, the elements of the algebra are denoted by  $J_z, J_+, J_-$  with

$$\begin{aligned}
 J_z &= \frac{1}{2} (E_{1,1} - E_{2,2}) \\
 J_- &= E_{1,2} \\
 J_+ &= E_{2,1}
 \end{aligned} \tag{4.82}$$

In this notation, the action of the elements of the algebra is

$$\begin{aligned}
 J_z |J, M\rangle &= M |J, M\rangle \\
 J_- |J, M\rangle &= [J(J+1) - M(M-1)]^{1/2} |J, M-1\rangle \\
 J_+ |J, M\rangle &= [J(J+1) - M(M+1)]^{1/2} |J, M+1\rangle .
 \end{aligned} \tag{4.83}$$

### 4.13 Tensor Products

With the representations  $[\lambda_1, \lambda_2, \dots, \lambda_n]$  one can form tensor products. The outer product of two tensors is denoted by

$$[\lambda'_1, \lambda'_2, \dots, \lambda'_n] \otimes [\lambda''_1, \lambda''_2, \dots, \lambda''_n] = \sum \oplus [\lambda_1, \lambda_2, \dots, \lambda_n]. \quad (4.84)$$

A crucial problem is to find what are the representations contained in the product. For  $u(n)$  and  $su(n)$  the representations contained in the product can be simply obtained using a set of rules, known also as Young calculus.

**Rule 1** *Product by a symmetric representation*

Consider the product of a generic representation  $[\lambda'_1, \lambda'_2, \dots, \lambda'_n]$  by a symmetric representation  $[\lambda''_1, 0, \dots, 0]$ , for Example

$$\begin{array}{c} \square \square \otimes \square \square \\ \square \end{array} \quad (4.85)$$

Replace the second factor by  $a$ 's

$$\begin{array}{c} \square \square \otimes a a \\ \square \end{array} \quad (4.86)$$

Place the  $a$ 's in all possible ways but no two  $a$ 's in the same column

$$\begin{array}{c} \square \square a a \oplus \square \square a \oplus \square \square a \oplus \square \square \\ \square \qquad \qquad \square a \qquad \square \qquad \square a \\ \qquad \qquad \qquad \qquad \qquad a \qquad \qquad a \end{array} \quad (4.87)$$

**Example 18.** *Product of fundamental representations*

$$\square \otimes \square = \square \square \oplus \square \quad (4.88)$$

When written in the notation (4.84)

$$[1] \otimes [1] = [2] \oplus [1, 1] . \quad (4.89)$$

**Rule 2** *Product by a generic representation*

Consider the product of a generic representation  $[\lambda'_1, \lambda'_2, \dots, \lambda'_n]$  by a generic representation  $[\lambda''_1, \lambda''_2, \dots, \lambda''_n]$ , for Example

$$\begin{array}{c} \square \square \otimes \square \square \\ \square \qquad \square \end{array} \quad (4.90)$$

Replace the second factor by  $a$ 's,  $b$ 's, . . .



Consider, for example, the product  $(1) \otimes (1)$ . Using the relation  $\lambda = 2J$ , this product can be converted to

$$su(2) : \quad [2] \otimes [2] . \tag{4.95}$$

This product can be performed using the rules 1 and 3,

$$\square\square \otimes aa = \square\square a a \oplus \begin{array}{c} \square\square a \\ a \end{array} \oplus \begin{array}{c} \square\square \\ a a \end{array} \tag{4.96}$$

that is

$$[2] \otimes [2] = [4] \oplus [2] \oplus [0] . \tag{4.97}$$

This can be converted back to  $so(3)$  using  $J = \lambda/2$ , with result

$$(1) \otimes (1) = (2) \oplus (1) \oplus (0) . \tag{4.98}$$

**Example 22.** *Multiplication rules of  $so(6)$*

Consider as another example, the product  $(2, 0, 0) \otimes (1, 0, 0)$ . Using the relations  $\lambda_1 = \sigma_1 + \sigma_2, \lambda_2 = \sigma_1 - \sigma_3, \lambda_3 = \sigma_2 - \sigma_3$  given in (4.51), this product can be converted to

$$su(4) : \quad [2, 2, 0] \otimes [1, 1, 0] . \tag{4.99}$$

This product can be performed using the rules for  $su(4)$

$$\begin{array}{c} \square\square \otimes a = \square\square a \oplus \begin{array}{c} \square\square a \\ b \end{array} \oplus \begin{array}{c} \square\square \\ a \\ b \end{array} \\ \square\square \quad b \quad \square\square b \quad \square\square \quad \square\square \end{array} , \tag{4.100}$$

that is

$$[2, 2, 0] \otimes [1, 1, 0] = [3, 3, 0] \oplus [3, 2, 1] \oplus [1, 1, 0] . \tag{4.101}$$

This can now be converted back to  $so(6)$  using  $\sigma_1 = \frac{\lambda_1 + \lambda_2 - \lambda_3}{2}, \sigma_2 = \frac{\lambda_1 - \lambda_2 + \lambda_3}{2}, \sigma_3 = \frac{\lambda_1 - \lambda_2 - \lambda_3}{2}$ , with result

$$(2, 0, 0) \otimes (1, 0, 0) = (3, 0, 0) \oplus (2, 1, 0) \oplus (1, 0, 0) . \tag{4.102}$$

### 4.14 Non-Canonical Chains

Gel'fand canonical chains of unitary and orthogonal algebras provide a complete solution to the branching problem for these algebras. However, in most problems in physics, one needs to consider non-canonical chains. For example, quite often the rotation algebra  $so(3)$  needs to be considered as a subalgebra of  $u(n)$ . Also, the branching problem for symplectic algebras  $sp(n)$ , not addressed by Gel'fand and Cetlin, needs be considered.

**Example 23.** *The chain  $u(6) \supset so(6) \supset so(5) \supset so(3) \supset so(2)$*

This chain has two non-canonical steps,  $u(6) \supset so(6)$  and  $so(5) \supset so(3)$  and illustrates two of the most important cases often encountered:  $u(n) \supset o(n)$ ;  $u(n) \supset sp(n)$ ;  $o(n) \supset o(n - 2)$ ;  $sp(n) \supset sp(n - 2)$ . The other two cases,  $u(n) \supset sp(n)$  and  $sp(n) \supset sp(n - 2)$ , are also encountered, especially the former, as discussed in Sect. 8.4.

The decomposition of irreducible representations of  $g$  into representations of  $g'$  for non-canonical chains

$$\left| \begin{array}{cc} g \supset g' \\ \downarrow \quad \downarrow \\ [\lambda] \quad [\mu] \end{array} \right\rangle \tag{4.103}$$

is one of the most difficult problems in group theory. A method often used is the so-called building-up process. The decomposition is constructed by taking successive products of the fundamental representations of  $g$  and  $g'$ . If one of the algebras is  $su(n)$ , equivalences are used. The method is illustrated by considering a problem of interest in nuclear physics.

**Example 24.** *Decomposition of representations of  $u(3)$  and  $su(3)$  into representations of  $so(3)$*

In order to decompose representations of  $u(3)$  and  $su(3)$  into representations of  $so(3)$  it is convenient to consider the decomposition  $u(3) \supset so(3)$  and use equivalence relations to obtain  $su(3) \supset so(3)$ . It is also convenient, for clarity, to use Young notation to label representations of  $u(3)$  and  $su(3)$  and angular momentum notation to label representations of  $so(3)$ . In this notation, the fundamental representation [1] of  $u(3)$  has Young tableau  $\square$ , while the fundamental representation (1) of  $so(3)$  has  $L = 1$ . The representation  $\square$  of  $u(3)$  contains only the representation (1) of  $so(3)$ , as one can see by noting that  $\dim[\square] = 3 = \dim(1)$  and that there are no other representations of  $so(3)$  with  $\dim = 3$ . (The dimensions of the representations play an important role in the building-up process. The dimension of the representations of  $u(3)$  is given in (4.56), while that of  $so(3)$  is given in (4.64).) The decomposition of the fundamental representation of  $u(3)$  into representations of  $so(3)$  is thus

$$\left| \begin{array}{cc} u(3) \supset so(3) \\ \downarrow \quad \downarrow \\ [1] \equiv \square \quad (1) \end{array} \right\rangle . \tag{4.104}$$

Consider now the products of representations of  $u(3)$  obtained by using the rules of Sect. 13 and of  $so(3)$  obtained either by using the rules of Sect. 13 or, equivalently, by using the angular momentum rule

$$| L_1 - L_2 | \leq L \leq | L_1 + L_2 | \tag{4.105}$$

which results from the isomorphism  $so(3) \sim su(2)$ , as described in Example 18. The product of the fundamental representation  $[1] \equiv \square$  of  $u(3)$  with itself, and of the fundamental representation (1) of  $so(3)$  with itself gives

$$\begin{aligned} u(3) : \quad & \square \otimes \square = \square\square \oplus \begin{matrix} \square \\ \square \end{matrix} \\ so(3) : \quad & (1) \otimes (1) = (0) \oplus (1) \oplus (2). \end{aligned} \tag{4.106}$$

The representation  $[1, 1]$  of  $u(3)$  contains only the representation (1) of  $so(3)$  as one can see by noting that  $\dim[1, 1] = 3 = \dim(1)$  and that there is no other representation of  $so(3)$  with  $\dim = 3$ . Thus

$$\left| \begin{array}{cc} u(3) \supset so(3) \\ \downarrow \quad \downarrow \\ [1, 1] \quad (1) \end{array} \right\rangle. \tag{4.107}$$

The remaining representations  $(0) \oplus (2)$  of  $so(3)$  must belong to the representation  $\square\square \equiv [2]$  of  $u(3)$ . This is verified by a dimensional check,  $\dim[2] = 6$ ,  $\dim((0) \oplus (2)) = 1 + 5 = 6$ . Thus

$$\left| \begin{array}{cc} u(3) \supset so(3) \\ \downarrow \quad \downarrow \\ [2] \quad (0) \oplus (2) \end{array} \right\rangle. \tag{4.108}$$

When going from  $u(3)$  to  $su(3)$ , by virtue of equivalence relation 2, we have  $[1, 1] \equiv [1]$ , again showing that the representation [1] of  $su(3)$  contains only (1) of  $so(3)$ .

Next, consider the products

$$\begin{aligned} u(3) : \quad & \begin{matrix} \square & & \square \\ \square & \otimes & \square \end{matrix} = \begin{matrix} \square & \square \\ \square & \square \end{matrix} \oplus \begin{matrix} \square \\ \square \end{matrix} \\ so(3) : \quad & (1) \otimes (1) = (0) \oplus (1) \oplus (2). \end{aligned} \tag{4.109}$$

The dimension of the representation  $[1, 1, 1]$  of  $u(3)$  is 1 and therefore

$$\left| \begin{array}{cc} u(3) \supset so(3) \\ \downarrow \quad \downarrow \\ [1, 1, 1] \quad (0) \end{array} \right\rangle. \tag{4.110}$$

Consequently  $[2, 1]$  must contain the remaining representations  $(1) \oplus (2)$

$$\left| \begin{array}{cc} u(3) \supset so(3) \\ \downarrow \quad \downarrow \\ [2, 1] \quad (1) \oplus (2) \end{array} \right\rangle. \tag{4.111}$$

The dimensional check is  $\dim[2, 1] = 8$ ,  $\dim((1) \oplus (2)) = 3 + 5 = 8$ . When going from  $u(3)$  to  $su(3)$ , use of the equivalence relation 1 gives



$$\begin{array}{l} \square \\ \square \equiv [1, 1, 1] \equiv [0] \\ \square \end{array} \tag{4.112}$$

A tedious but straightforward procedure, sometimes called plethism, gives then Table 4.2. In this table, for clarity, both the representations of  $u(3)$  and of  $su(3)$  (with their equivalences) are given. Also the number  $t$ , the rank of the tensor in (4.5), is given. This number is important in applications since it denotes the total number of particles

$$t = \lambda_1 + \dots + \lambda_n \tag{4.113}$$

Consider now the representation  $[4, 2]$  of  $su(3)$ . From the table, one can see that it contains two  $L = 2$  representations of  $so(3)$ . Thus the decomposition is not unique and one needs an additional quantum number to distinguish the two  $L = 2$  representations. The identification and use of this quantum number (missing label) is one of the most subtle points of representation

**Table 4.2.** Decomposition of representations of  $u(3)$  and  $su(3)$  into representations of  $so(3)$

$t$	$u(3)$	$su(3)$	$so(3)$
0	[0]	[0]	0
1	[1]	[1]	1
2	[2]	[2]	0, 2
	[1, 1]	[1, 1] $\equiv$ [1]	1
3	[3]	[3]	1, 3
	[2, 1]	[2, 1]	1, 2
	[1, 1, 1]	[1, 1, 1] $\equiv$ [0]	0
4	[4]	[4]	0, 2, 4
	[3, 1]	[3, 1]	1, 2, 3
	[2, 2]	[2, 2] $\equiv$ [2]	0, 2
	[2, 1, 1]	[2, 1, 1] $\equiv$ [1]	1
5	[5]	[5]	1, 3, 5
	[4, 1]	[4, 1]	1, 2, 3, 4
	[3, 2]	[3, 2] $\equiv$ [3, 1]	1, 2, 3
	[3, 1, 1]	[3, 1, 1] $\equiv$ [2]	0, 2
	[2, 2, 1]	[2, 2, 1] $\equiv$ [1, 1]	1
6	[6]	[6]	0, 2, 4, 6
	[5, 1]	[5, 1]	1, 2, 3, 4, 5
	[4, 2]	[4, 2]	0, 2 <sup>2</sup> , 3, 4
	[4, 1, 1]	[4, 1, 1] $\equiv$ [3]	1, 3
	[3, 3]	[3, 3] $\equiv$ [3]	1, 3
	[3, 2, 1]	[3, 2, 1] $\equiv$ [2, 1]	1, 2
	[2, 2, 2]	[2, 2, 2] $\equiv$ [0]	0

theory. To find how many missing labels there are in a given problem, go to the canonical chain (which has no missing labels) and count the total number of labels.

**Example 25.** *Missing labels for  $su(3) \supset so(3)$*

Consider the non-canonical chain

$$\left| \begin{array}{ccc} su(3) & \supset & so(3) \supset so(2) \\ \downarrow & & \downarrow \quad \downarrow \\ [\lambda_1, \lambda_2] & & L \quad M \end{array} \right\rangle \quad (4.114)$$

and its associated canonical chain

$$\left| \begin{array}{ccc} su(3) & \supset & u(2) \supset u(1) \\ \downarrow & & \downarrow \quad \downarrow \\ [\lambda_{1,3}, \lambda_{2,3}, 0] & & [\lambda_{1,2}, \lambda_{2,3}] \quad [\lambda_{1,1}] \end{array} \right\rangle. \quad (4.115)$$

The non-canonical chain has 4 labels, while the canonical chain has 5. There is thus one missing label in the non-canonical chain.

Extensive tables for the non-canonical reductions

$$\left\{ \begin{array}{l} u(7) \supset so(7) \\ u(5) \supset so(5) \\ u(3) \supset so(3) \end{array} \right\} \left\{ \begin{array}{l} u(8) \supset sp(8) \\ u(6) \supset sp(6) \\ u(4) \supset sp(4) \end{array} \right\} \quad (4.116)$$

$$\left\{ \begin{array}{l} u(6) \supset so(6) \\ u(4) \supset so(4) \end{array} \right\}$$

exist. [M. Hamermesh, *Group Theory and Application to Physical Problems*, Addison-Wesley, Reading (1962).] Tables are also available for the reduction  $so(5) \supset so(3)$ . [F. Iachello and A. Arima, *The Interacting Boson Model*, Cambridge University Press, 1987.]

Although not of the type mentioned at the beginning of the section, it is worth noting that tables of non-canonical reductions  $sp(8) \supset spin(3)$ ,  $sp(6) \supset spin(3)$ ,  $sp(4) \supset spin(3)$  are also available. [R.H. Flowers, *Studies in  $j$ - $j$  coupling. I. Classification of nuclear and atomic states*, Proc. Roy. Soc. A212, 248 (1952).]

# 5 Casimir Operators and Their Eigenvalues

## 5.1 Definitions

An operator which commutes with all the elements of a Lie algebra,  $g$ , is called an invariant, or Casimir operator,  $C$

$$[C, X_\rho] = 0 \quad \text{for any } X_\rho \in g . \quad (5.1)$$

The operator is called of order  $p$ , if it is built from products of  $p$  elements

$$C_p = \sum_{\alpha_1, \alpha_2, \dots, \alpha_p} f^{\alpha_1 \alpha_2 \dots \alpha_p} X_{\alpha_1} X_{\alpha_2} \dots X_{\alpha_p} . \quad (5.2)$$

It lies in the enveloping algebra of  $g$ ,  $T(g)$ .

## 5.2 Independent Casimir Operators

The number of independent Casimir operators,  $C$ , of a Lie algebra  $g$ , is equal to the rank  $l$  of  $g$ , and hence equal to the number of labels that characterize the irreducible representations of  $g$ . As mentioned in Chap. 1, if  $C$  is a Casimir operator, so is  $aC$ ,  $C^2$ ,  $\dots$

### 5.2.1 Casimir Operators of $u(n)$

The algebra of  $u(n)$  has  $n$  independent Casimir operators of order  $1, 2, \dots, n$

$$C_1, C_2, \dots, C_n . \quad (5.3)$$

A construction of these operators was given in Chap. 1 in terms of the structure constants  $c_{\alpha\beta}^\gamma$ . For the algebra  $u(n)$ , if the elements are denoted by  $E_{ij}$  ( $i, j = 1, \dots, n$ ), that is if the algebra is realized as in Chap. 2, Sect. 10, the Casimir operators of order  $p$  can be written as

$$C_p = E_{i_1 i_2} E_{i_2 i_3} \dots E_{i_{p-1} i_p} E_{i_p i_1} \quad p = 1, 2, \dots, n \quad (5.4)$$

(summation over repeated indices). In particular

$$C_1 = E_{i_1 i_1} , \quad (5.5)$$

or, displaying explicitly the summation

$$C_1 = \sum_{i=1}^n E_{ii} . \quad (5.6)$$

Using the form (5.4), one can show that the Casimir operators so defined are independent. [A.O. Barut and R. Raczka, *Theory of Group Representations and Applications*, World Scientific, Singapore, 1986.]

### 5.2.2 Casimir Operators of $su(n)$

The independent Casimir operators of  $su(n)$  are of order  $2, 3, \dots, n$

$$C_2, C_3, \dots, C_n \quad (5.7)$$

that is the same as  $u(n)$  but with  $C_1$  omitted. The elements of  $su(n)$  are obtained from those of  $u(n)$  by keeping the off-diagonal elements the same, replacing the diagonal ones with

$$\tilde{E}_{ii} = E_{ii} - \frac{1}{n} \sum_{j=1}^n E_{jj} . \quad (5.8)$$

and deleting  $\tilde{E}_{nn}$ . The linear Casimir operator of  $su(n)$  is

$$C_1 = \sum_{i=1}^n \tilde{E}_{ii} = 0 , \quad (5.9)$$

and can thus be omitted.

### 5.2.3 Casimir Operators of $so(n)$ , $n = \text{Odd}$

The independent Casimir operators of  $so(n)$ ,  $n = \text{odd}$  are

$$C_2, C_4, C_6, \dots, C_{2\nu-2}, C_{2\nu}; \quad \nu = \frac{n-1}{2} . \quad (5.10)$$

These operators are all of even order.

### 5.2.4 Casimir Operators of $so(n)$ , $n = \text{Even}$

The independent Casimir operators of  $so(n)$ ,  $n = \text{even}$  are

$$C_2, C_4, C_6, \dots, C_{2\nu-2}, C'_\nu; \quad \nu = \frac{n}{2} . \quad (5.11)$$

The operators  $C$  are of even order, while  $C'$  is of even or odd order depending on whether  $\nu$  is even or odd. Comparing with the previous case (5.10) one can see that there is a peculiarity here, since the operator of order  $2\nu$  is replaced by an operator of order  $\nu$ . The operator  $C'_\nu$  is needed to distinguish between the mirror conjugate representations  $[\lambda_1, \lambda_2, \dots, +\lambda_\nu]$  and  $[\lambda_1, \lambda_2, \dots, -\lambda_\nu]$ .

**Example 1.** *Casimir operators of  $so(4)$  and  $so(6)$*

The Casimir operators of  $so(4)$  are of order

$$C_2, C'_2 . \tag{5.12}$$

The Casimir operators of  $so(6)$  are of order

$$C_2, C_4, C'_3 . \tag{5.13}$$

**5.2.5 Casimir Operators of  $sp(n)$ ,  $n = \text{Even}$**

The independent Casimir operators of  $sp(n)$ ,  $n = \text{even}$  are

$$C_2, C_4, C_6, \dots, C_{2\nu-2}, C_{2\nu}; \quad \nu = \frac{n}{2} . \tag{5.14}$$

These operators are all of even order.

**5.2.6 Casimir Operators of the Exceptional Algebras**

The independent Casimir operators of the exceptional algebras are of order

$$\begin{array}{ll} G_2 & C_2, C_6 \\ F_4 & C_2, C_6, C_8, C_{12} \\ E_6 & C_2, C_5, C_6, C_8, C_9, C_{12} \\ E_7 & C_2, C_6, C_8, C_{10}, C_{12}, C_{14}, C_{18} \\ E_8 & C_2, C_8, C_{12}, C_{14}, C_{18}, C_{20}, C_{24}, C_{30} \end{array} \tag{5.15}$$

**5.3 Complete Set of Commuting Operators**

For any given Lie algebra  $g$ , one is often interested in constructing a complete set of commuting operators. This is done by considering the decomposition into subalgebras  $g \supset g' \supset g'' \supset \dots$ . In view of problems connected with missing labels, the construction of a complete set of commuting operators is straightforward only for canonical chains. The construction parallels that of the labels given Chap. 4 (Gel'fand construction) and applies to the canonical chains  $u(n) \supset u(n-1) \supset \dots \supset u(2) \supset u(1)$  and  $so(n) \supset so(n-1) \supset \dots \supset so(2)$ . The construction for the symplectic algebras  $sp(n)$  is more complex and it will not be discussed here.

### 5.3.1 The Unitary Algebra $u(n)$

The commuting Casimir operators can be arranged into a triangular pattern

$$\begin{array}{ccccccc}
 C_{1,n} & & C_{2,n} & & & C_{n-1,n} & & C_{n,n} \\
 & C_{1,n-1} & & & & & C_{n-1,n-1} & \\
 & & \dots & & & \dots & & \\
 & & & C_{1,2} & & C_{2,2} & & \\
 & & & & C_{1,1} & & & 
 \end{array} \tag{5.16}$$

where  $C_{1,n}$  denotes  $C_1(u(n)), \dots$

### 5.3.2 The Orthogonal Algebra $so(n)$ , $n = \text{Odd}$

The complete set of commuting operators is, for  $n = 2k + 1$ ,

$$\begin{array}{ccccccc}
 C_{2,2k+1} & C_{4,2k+1} & & & & C_{2k-2,2k+1} & C_{2k,2k+1} \\
 C_{2,2k} & C_{4,2k} & & & & C_{2k-2,2k} & C'_{k,2k} \\
 & & \dots & & & & \\
 & & & C_{2,4} & & C'_{2,4} & \\
 & & & & C_{2,3} & & \\
 & & & & C'_{1,2} & & 
 \end{array} \tag{5.17}$$

### 5.3.3 The Orthogonal Algebra $so(n)$ , $n = \text{Even}$

The complete set of commuting operators is for  $n = 2k + 2$ ,

$$\begin{array}{ccccccc}
 C_{2,2k+2} & & C_{4,2k+2} & & & C_{2k,2k+2} & & C'_{k+1,2k+2} \\
 & C_{2,2k+1} & & & & & C_{2k,2k+1} & \\
 & & \dots & & & \dots & & \\
 & & & C_{2,4} & & C'_{2,4} & & \\
 & & & & C_{2,3} & & & \\
 & & & & C'_{1,2} & & & 
 \end{array} \tag{5.18}$$

## 5.4 Eigenvalues of Casimir Operators

The eigenvalues of the Casimir operators of all classical Lie algebras in a representation  $|\lambda\rangle$ , denoted in short by

$$\langle \lambda_1, \dots, \lambda_n | C_p | \lambda_1, \dots, \lambda_n \rangle = \langle C_p \rangle \tag{5.19}$$

were worked out in a series of papers by Perelomov and Popov in the 1960's. [A.M. Perelomov and V.S. Popov, *Casimir Operators for  $u(n)$  and  $su(n)$* , Sov. J. of Nucl. Phys. 3, 676–680 (1966); *Casimir Operators of Orthogonal and Symplectic Groups*, Sov. J. of Nucl. Phys. 3, 819–824 (1966).] The algorithm to obtain the eigenvalues is described below.

**5.4.1 The Algebras  $u(n)$  and  $su(n)$**

To find the eigenvalues of the Casimir operators,  $C_p$ , construct the quantities

$$S_k = \sum_{i=1}^n (\ell_i^k - \rho_i^k), \quad \rho_i = n - i, \quad \ell_i = m_i + n - i \quad (5.20)$$

with

$$m_i = \begin{cases} \lambda_i & \text{for } u(n) \\ \lambda_i - \frac{\lambda}{n} & \text{for } su(n) \end{cases}, \quad \lambda = \sum_{i=1}^n \lambda_i. \quad (5.21)$$

Construct the function

$$\varphi(z) = \sum_{k=2}^{\infty} a_k z^k, \quad a_k = \sum_{j=1}^{k-1} \frac{(k-1)!}{j!(k-j)!} S_j. \quad (5.22)$$

Define the quantities  $B_p$  by

$$\exp\{-\varphi(z)\} = 1 - \sum_{p=0}^{\infty} z^{p+1}, \quad B_0 = 0. \quad (5.23)$$

Then, the eigenvalue of  $C_p$  in the representation  $[\lambda_1, \lambda_2, \dots, \lambda_n]$  is  $\langle C_p \rangle = B_p - n B_{p-1}$ . This algorithm gives the following eigenvalues of Casimir operators of order,  $p \leq 3$ . For  $u(n)$

$$\begin{aligned} \langle C_1 \rangle &= S_1 \\ \langle C_2 \rangle &= S_2 - (n-1)S_1 \\ \langle C_3 \rangle &= S_3 - \left(n - \frac{3}{2}\right)S_2 - \frac{1}{2}S_1^2 - (n-1)S_1 \\ &\dots \end{aligned} \quad (5.24)$$

and for  $su(n)$

$$\begin{aligned} \langle C_1 \rangle &= 0 \\ \langle C_2 \rangle &= S_2 \\ \langle C_3 \rangle &= S_3 - \left(n - \frac{3}{2}\right)S_2 \\ &\dots \end{aligned} \quad (5.25)$$

**Example 2.** *Eigenvalue of the quadratic Casimir operator of  $su(2)$  in the representation  $[\lambda_1]$*

From

$$S_2 = \sum_{i=1}^2 (\ell_i^2 - \rho_i^2)$$

$$\begin{aligned}
\rho_1 &= 1, \rho_2 = 0 \\
\ell_1 &= m_1 + 1, \ell_2 = m_2 \\
m_1 &= \lambda_1 - \frac{\lambda_1}{2}, m_2 = -\frac{\lambda_1}{2}
\end{aligned} \tag{5.26}$$

one obtains

$$\langle C_2 \rangle = \frac{1}{2} \lambda_1 (\lambda_1 + 2) . \tag{5.27}$$

**Example 3.** *Eigenvalue of the quadratic Casimir operator of  $su(3)$  in the representation  $[\lambda_1, \lambda_2]$*

From

$$\begin{aligned}
S_2 &= \sum_{i=1}^3 (\ell_i^2 - \rho_i^2) \\
\rho_1 &= 2, \rho_2 = 1, \rho_3 = 0 \\
\ell_1 &= m_1 + 2, \ell_2 = m_2 + 1, \ell_3 = m_3 \\
m_1 &= \lambda_1 - \frac{\lambda_1 + \lambda_2}{3}, m_2 = \lambda_2 - \frac{\lambda_1 + \lambda_2}{3}, m_3 = -\frac{\lambda_1 + \lambda_2}{3}
\end{aligned} \tag{5.28}$$

one obtains

$$\langle C_2 \rangle = \frac{6}{9} [\lambda_1^2 + \lambda_2^2 - \lambda_1 \lambda_2 + 3\lambda_1] . \tag{5.29}$$

The coefficient  $\frac{6}{9}$  is usually omitted, since Casimir operators are defined up to a multiplicative constant.

**Example 4.** *Eigenvalue of the quadratic Casimir operator of  $su(5)$  in the representation  $[\lambda_1, \lambda_2, \lambda_3, \lambda_4]$*

From

$$\begin{aligned}
S_2 &= \sum_{i=1}^5 (\ell_i^2 - \rho_i^2) \\
\rho_1 &= 4, \rho_2 = 3, \rho_3 = 2, \rho_4 = 1, \rho_5 = 0 \\
\ell_1 &= m_1 + 4, \ell_2 = m_2 + 3, \ell_3 = m_3 + 2, \ell_4 = m_4 + 1, \ell_5 = m_5 \\
m_1 &= \lambda_1 - \frac{\lambda}{5}, m_2 = \lambda_2 - \frac{\lambda}{5}, m_3 = \lambda_3 - \frac{\lambda}{5}, m_4 = \lambda_4 - \frac{\lambda}{5}, m_5 = -\frac{\lambda}{5}
\end{aligned} \tag{5.30}$$

one obtains

$$\langle C_2 \rangle = \lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2 - \frac{\lambda^2}{5} + 8\lambda_1 + 6\lambda_2 + 4\lambda_3 + 2\lambda_4 - 4\lambda , \tag{5.31}$$

with

$$\lambda = \lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 . \tag{5.32}$$



**Example 5.** *Eigenvalue of the linear and quadratic Casimir operators of  $u(5)$  in the representation  $[\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5]$*

Straightforward application of the formulas above give

$$\langle C_1 \rangle = S_1 = \lambda \tag{5.33}$$

$$\begin{aligned} \langle C_2 \rangle &= S_2 - 4S_1 \\ &= \lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2 + \lambda_5^2 + 4\lambda_1 + 2\lambda_2 - 2\lambda_4 . \end{aligned} \tag{5.34}$$

For example, for the symmetric representation  $[N, 0, 0, 0, 0]$ ,

$$\langle C_2 \rangle = N(N + 4) . \tag{5.35}$$

### 5.4.2 The Orthogonal Algebra $so(2n + 1)$

To find the eigenvalues, construct the quantities

$$\begin{aligned} S_k &= \sum_{i=-n}^{+n} (\ell_i^k - \rho_i^k); & \rho_i &= \ell_i - f_i \\ \begin{cases} \ell_i = f_i + n + i - \vartheta_{0i} \\ \ell_{-i} = -\ell_i + 2n - 1 & (i \neq 0) \\ \ell_0 = n \end{cases} & & \vartheta_{ji} &= \begin{cases} 1 & j < i \\ 0 & j \geq i \end{cases} \\ f_{-i} &= -f_i, & f_0 &= 0 \\ S_0 &= S_1 = 0 . \end{aligned} \tag{5.36}$$

Construct the function

$$\varphi(z) = \sum_{k=3}^{\infty} a_k z^k, \quad a_k = \sum_{j=2}^{k-1} \frac{(k-1)!}{j!(k-j)!} S_j . \tag{5.37}$$

Define the quantities  $B_p$  by

$$\exp(-\varphi(z)) = 1 - \sum_{p=2}^{\infty} B_p z^{p+1}, \quad B_0 = B_1 = 0 . \tag{5.38}$$

Then, the expectation value of  $C_p$  in the representation  $[f_n, f_{n-1}, \dots, f_1]$  (note the inverse order of the labels) is

$$\langle C_p \rangle = (2n + 1) \delta_{p0} + B_p - \left( n + \frac{1}{2} \right) B_{p-1} - \sum_{q=1}^{p-1} \left[ B_q - \left( n + \frac{1}{2} \right) B_{q-1} \right] n^{p-q} . \tag{5.39}$$

Finally, convert  $[f_n, f_{n-1}, \dots, f_1]$  to the standard notation  $[\mu_1, \mu_2, \dots, \mu_n]$ . This algorithm gives

$$\begin{aligned}
\langle C_0 \rangle &= 2n + 1 \\
\langle C_1 \rangle &= 0 \\
\langle C_2 \rangle &= S_2 \\
&\dots
\end{aligned}
\tag{5.40}$$

**Example 6.** *Eigenvalue of the quadratic Casimir operator of  $so(3)$  in the representation  $[\mu_1]$*

From

$$\begin{aligned}
\langle C_2 \rangle = S_2 &= \sum_{i=-1}^{+1} [\ell_i^2 - (\ell_i - f_i)^2] \\
\left\{ \begin{array}{l} \ell_1 = f_1 + 1 + 1 - 1 = f_1 + 1 \\ \ell_{-1} = -f_1 - 1 + 2 - 1 = -f_1 \\ \ell_0 = 1 \end{array} \right. , & \quad \begin{array}{l} f_{-1} = -f_1 \\ f_0 = 0 \end{array}
\end{aligned}
\tag{5.41}$$

one obtains

$$\langle C_2 \rangle = 2f_1(f_1 + 1) \tag{5.42}$$

Converting to the standard notation  $[\mu_1]$

$$\langle C_2 \rangle = 2\mu_1(\mu_1 + 1) \tag{5.43}$$

Note that these constructions are consistent with the isomorphisms of Chap. 4. Example 4.10 gives for the isomorphism  $su(2) \sim so(3)$ , the relation  $\mu_1 = \frac{\lambda_1}{2}$ . Inserting this relation into (5.43) gives

$$\langle C_2(su(2)) \rangle = \frac{1}{2} \lambda_1(\lambda_1 + 2) \tag{5.44}$$

as in (5.27). The additional normalization factor of two in (5.43) is usually omitted and the label  $\mu_1$  is called  $J$  in quantum mechanics textbooks. The eigenvalues of the quadratic Casimir operator of the rotation algebra (and group) is then written as

$$\langle C_2(so(3)) \rangle = J(J + 1) . \tag{5.45}$$

**Example 7.** *Eigenvalue of the quadratic Casimir operator of  $so(5)$  in the representation  $[\mu_1, \mu_2]$*

From

$$\langle C_2 \rangle = S_2 = \sum_{i=-2}^{+2} (2\ell_i f_i - f_i^2) .$$

$$\left\{ \begin{array}{l} \ell_2 = f + 3 \\ \ell_1 = f_1 + 1 \\ \ell_0 = 2 \\ \ell_{-2} = -f_2 \\ \ell_{-1} = -f_1 \end{array} \right. \left\{ \begin{array}{l} f_{-2} = -f_2 \\ f_{-1} = f_1 \\ f_0 = 0 \end{array} \right. \quad (5.46)$$

one obtains

$$\langle C_2 \rangle = 2 f_2(f_2 + 3) + 2 f_1(f_1 + 1) . \quad (5.47)$$

Converting to the standard notation  $[\mu_1, \mu_2]$

$$\langle C_2 \rangle = 2[\mu_1(\mu_1 + 3) + \mu_2(\mu_2 + 1)] . \quad (5.48)$$

### 5.4.3 The Symplectic Algebra $sp(2n)$

For  $sp(2n)$ , construct the quantities

$$S_k = \sum_{i=-n, i \neq 0}^{+n} (\ell_i^k - \rho_i^k), \quad \rho_i = \ell_i - f_i$$

$$\left\{ \begin{array}{l} \ell_i = f_i + n + i \\ \ell_{-i} = -\ell_i + 2n \end{array} \right. \quad (5.49)$$

$$f_{-i} = -f_i, \quad S_0 = S_1 = 0$$

and the function  $\varphi(z)$  as before. Then

$$\langle C_p \rangle = 2n \delta_{p0} + (B_p - n B_{p-1}) - \sum_{q=1}^{p-1} (B_q - n B_{q-1}) \left( n + \frac{1}{2} \right)^{p-q} . \quad (5.50)$$

Finally, convert  $[f_n, f_{n-1}, \dots, f_1]$  to the standard notation  $[\mu_1, \mu_2, \dots, \mu_\nu]$ . The Casimir operators of lowest order,  $p \leq 2$ , are

$$\begin{aligned} \langle C_0 \rangle &= 2n \\ \langle C_1 \rangle &= 0 \\ \langle C_2 \rangle &= S_2 \\ &\dots \end{aligned} \quad (5.51)$$

The constant  $2n$  does no count as an independent Casimir operator.

**Example 8.** *Eigenvalue of the quadratic Casimir operator of  $sp(2)$  in the representation  $[\mu_1]$*

From

$$\begin{aligned} \langle C_2 \rangle = S_2 &= \sum_{i=-1, i \neq 0}^1 (f_i^2 - 2\ell_i f_i) \\ \ell_1 = f_1 + 2, \ell_{-1} &= -f_1, f_{-1} = -f_1 \end{aligned} \tag{5.52}$$

one obtains

$$\langle C_2 \rangle = -2f_1(f_1 + 2). \tag{5.53}$$

Converting to the standard notation  $[\mu_1]$

$$\langle C_2 \rangle = -2\mu_1(\mu_1 + 2). \tag{5.54}$$

Note once more the consistency with the isomorphisms  $sp(2) \sim su(2)$  apart from an overall minus sign.

#### 5.4.4 The Orthogonal Algebra $so(2n)$

For these algebras, construct the quantities

$$\begin{aligned} S_k &= \sum_{i=-n, i \neq 0}^{+n} (\ell_i^k - \rho_i^k), \quad \rho_i = \ell_i - f_i \\ \begin{cases} \ell_i = f_i + n + i - (1 + \varepsilon_i) \\ \ell_{-i} = -\ell_i + 2n - 2 \end{cases} & \quad \varepsilon_i = \begin{cases} 1 & i > 0 \\ -1 & i < 0 \\ 0 & i = 0 \end{cases} \\ f_{-i} = -f_i, \quad S_0 = S_1 = 0 & \end{aligned} \tag{5.55}$$

and the function  $\varphi(z)$  as before. Then

$$\langle C_p \rangle = 2n \delta_{p0} + (B_p - n B_{p-1}) - \sum_{q=1}^{p-1} (B_q - n B_{q-1}) \left( n - \frac{1}{2} \right)^{p-q}. \tag{5.56}$$

Finally, convert  $[f_n, f_{n-1}, \dots, f_1]$  to  $[\mu_1, \mu_2, \dots, \mu_n]$ . The eigenvalues of Casimir operators of lowest order,  $p \leq 2$ , are

$$\begin{aligned} \langle C_0 \rangle &= 2n \\ \langle C_1 \rangle &= 0 \\ \langle C_2 \rangle &= S_2 \\ &\dots \end{aligned} \tag{5.57}$$

**Example 9.** *Eigenvalue of the quadratic Casimir operator of  $so(2)$  in the representation  $[\mu_1]$*

From

$$S_2 = \sum_{i=-1, i \neq 0}^{+1} (2l_i f_i - f_i^2)$$

$$\begin{cases} \ell_1 = f_1 \\ \ell_{-1} = -f_1 \end{cases} \quad f_{-1} = -f_1 \tag{5.58}$$

one obtains

$$\langle C_2 \rangle = 2 f_1^2 . \tag{5.59}$$

Converting to standard notation  $[\mu_1]$

$$\langle C_2 \rangle = 2\mu_1^2 . \tag{5.60}$$

Since  $so(2)$  is Abelian it has also a linear invariant. The quadratic invariant  $C_2$ , is the square of the linear invariant  $C_1$ . In quantum mechanics textbooks, the label  $\mu_1 = M$  and the factor of two is omitted.

**Example 10.** *Eigenvalue of the quadratic Casimir operator of  $so(4)$  in the representation  $[\mu_1, \mu_2]$*

From

$$S_2 = \sum_{i=-2, i \neq 0}^{+2} (2l_i f_i - f_i^2)$$

$$\begin{cases} \ell_2 = f_2 + 2 \\ \ell_{-2} = -f_2 \end{cases} \quad \begin{cases} \ell_1 = f_1 + 1 \\ \ell_{-1} = -f_1 + 1 \end{cases} \quad \begin{cases} f_{-1} = -f_1 \\ f_{-2} = -f_2 \end{cases} \tag{5.61}$$

one obtains

$$\langle C_2 \rangle = 2 f_2(f_2 + 2) + 2f_1^2 . \tag{5.62}$$

Converting to standard notation  $[\mu_1, \mu_2]$

$$\langle C_2 \rangle = 2 [\mu_1(\mu_1 + 2) + \mu_2^2] . \tag{5.63}$$

**Example 11.** *Eigenvalue of the quadratic Casimir operator of  $so(6)$  in the representation  $[\mu_1, \mu_2, \mu_3]$*

In this case one has

$$\langle C_2 \rangle = 2 f_3(f_3 + 4) + 2 f_2(f_2 + 2) + 2f_1^2 \tag{5.64}$$

and, in standard notation,  $[\mu_1, \mu_2, \mu_3]$ ,

$$\langle C_2 \rangle = 2 [\mu_1(\mu_1 + 4) + \mu_2(\mu_2 + 2) + \mu_3^2] . \tag{5.65}$$

Note that for all algebras  $so(2n)$  the dependence on the last quantum number  $\mu_\nu$  is always quadratic, so that the eigenvalue for  $+\mu_\nu$  is the same as for  $-\mu_\nu$ .

## 5.5 Eigenvalues of Casimir Operators of Order One and Two

The eigenvalues of Casimir operators of order  $p \leq 2$  of all classical Lie algebras are summarized in Table 5.1. In this table, all labels are denoted by  $\lambda_i$ .

**Table 5.1.** Eigenvalues of Casimir operators of order  $p \leq 2$  of all classical Lie algebras

Algebra	Labels	Order	$\langle C_2 \rangle$
$u(n)$	$[\lambda_1, \lambda_2, \dots, \lambda_n]$	1	$\sum_{i=1}^n \lambda_i = \lambda$
$u(n)$	$[\lambda_1, \lambda_2, \dots, \lambda_n]$	2	$\sum_{i=1}^n \lambda_i (\lambda_i + n + 1 - 2i)$
$su(n)$	$[\lambda_1, \lambda_2, \dots, \lambda_{n-1}, 0]$	2	$\sum_{i=1}^n (\lambda_i - \frac{\lambda}{n}) (\lambda_i - \frac{\lambda}{n} + 2n - 2i)$
$so(2n + 1)$	$[\lambda_1, \lambda_2, \dots, \lambda_n]$	2	$\sum_{i=1}^n 2 \lambda_i (\lambda_i + 2n + 1 - 2i)$
$so(2n)$	$[\lambda_1, \lambda_2, \dots, \lambda_n]$	2	$\sum_{i=1}^n 2 \lambda_i (\lambda_i + 2n - 2i)$
$sp(2n)$	$[\lambda_1, \lambda_2, \dots, \lambda_n]$	2	$\sum_{i=1}^n 2 \lambda_i (\lambda_i + 2n + 2 - 2i)$

# 7 Boson Realizations

## 7.1 Boson Operators

Realizations of Lie algebras in terms of boson operators are of great interest for applications to a variety of problems in physics, most notably to oscillator problems in quantum mechanics and to algebraic models of rotation-vibration spectra of molecules (vibron model) and nuclei (interaction boson model).

Let  $b_\alpha (\alpha = 1, \dots, n)$  be a set of boson operators, satisfying the commutation relations

$$[b_\alpha, b_{\alpha'}^\dagger] = \delta_{\alpha\alpha'} \quad [b_\alpha, b_{\alpha'}] = [b_\alpha^\dagger, b_{\alpha'}^\dagger] = 0. \quad (7.1)$$

The operators  $b_\alpha$  are often called annihilation (or destruction) operators, while their hermitian conjugates  $b_\alpha^\dagger$  are called creation operators. The unitary algebra  $u(n)$  can be constructed by taking bilinear products of creation and annihilation operators

$$g \doteq G_{\alpha\beta} = b_\alpha^\dagger b_\beta \quad \alpha, \beta = 1, \dots, n. \quad (7.2)$$

The algebra is composed of  $n^2$  elements satisfying

$$[G_{\alpha\beta}, G_{\gamma\delta}] = \delta_{\beta\gamma} G_{\alpha\delta} - \delta_{\alpha\delta} G_{\gamma\beta}. \quad (7.3)$$

(When written in this form the algebra is the real form  $gl(n, R) \sim u(n)$ .) A basis for the representations of  $u(n)$  is

$$\mathcal{B} : \frac{1}{\mathcal{N}} \underbrace{b_\alpha^\dagger b_{\alpha'}^\dagger \cdots}_{N \text{ times}} |0\rangle \equiv |N\rangle \quad (7.4)$$

Here  $|0\rangle$  denotes a vacuum state, such that

$$b_\alpha |0\rangle = 0, \quad (7.5)$$

and  $\mathcal{N}$  is a normalization. In this construction, the basis states all belong to the totally symmetric irreducible representation

$$[N] \equiv \underbrace{\square \square \cdots \square}_N \quad (7.6)$$

of  $u(n)$  written in (7.4) as  $|N\rangle$ . The dimension of the basis is the dimension of the representation  $[N]$ , given in Chap. 4,

$$\dim[N] = \prod_{j=2,\dots,n} \left( \frac{N+j-1}{j-1} \right). \quad (7.7)$$

Thus, with boson operators it is possible to construct the Lie algebra  $u(n)$ , but only its symmetric representations (Bose-Einstein basis).

Any classical Lie algebra is a subalgebra of  $u(n)$  (Ado's theorem) and thus can be written as a linear combination of the elements  $G_{\alpha\beta}$ . In the following sections an explicit construction of the Lie algebras  $u(1)$ ,  $u(2)$ ,  $u(3)$ ,  $u(4)$ ,  $u(5)$ ,  $u(6)$  and  $u(7)$  and their subalgebras will be given.

## 7.2 The Unitary Algebra $u(1)$

This algebra can be simply constructed with one boson operator  $b$ , satisfying

$$[b, b^\dagger] = 1 \quad [b, b] = [b^\dagger, b^\dagger] = 0. \quad (7.8)$$

The algebra is Abelian and composed of the single element

$$g \doteq b^\dagger b. \quad (7.9)$$

This operator is called the number operator,  $\hat{N}$ . (A hat will be placed above the operators to distinguish them from their eigenvalues.) The basis, written as

$$\left| \begin{array}{c} u(1) \\ \downarrow \\ N \end{array} \right\rangle \quad (7.10)$$

is

$$\mathcal{B}: \quad \frac{1}{\sqrt{N!}} b^{\dagger N} |0\rangle \equiv |N\rangle. \quad (7.11)$$

(The capital letter  $N$  is used here, rather than the commonly used lowercase letter  $n$ , not to confuse it with the number of dimensions in  $u(n)$ .) In the case of  $u(1)$ , all representations are symmetric and thus can be constructed with boson operators. The boson operators satisfy

$$b^\dagger |N\rangle = \sqrt{N+1} |N+1\rangle; b |N\rangle = \sqrt{N} |N-1\rangle; b |0\rangle = 0. \quad (7.12)$$

**Example 1.** *The one dimensional harmonic oscillator in quantum mechanics*



The algebra  $u(1)$  can be used to describe the one-dimensional harmonic oscillator. The boson operators are related to the dimensionless coordinate,  $x$ , and momentum,  $p_x = -i\frac{d}{dx}$ , by

$$b = \frac{1}{\sqrt{2}}(x + ip_x), \quad b^\dagger = \frac{1}{\sqrt{2}}(x - ip_x). \quad (7.13)$$

The quantum mechanical Hamiltonian is

$$H = \frac{1}{2}(p_x^2 + x^2) = \frac{1}{2}\left(-\frac{d^2}{dx^2} + x^2\right). \quad (7.14)$$

When written in terms of elements of the Lie algebra  $u(1)$ , it reads

$$H = b^\dagger b + \frac{1}{2} = \hat{N} + \frac{1}{2}. \quad (7.15)$$

The representations of  $u(1)$  are labelled by the integer  $N = 0, 1, \dots$ . The eigenvalues of  $H$  are thus

$$E(N) = N + \frac{1}{2}, \quad N = 0, 1, \dots \quad (7.16)$$

The basis states are

$$|N\rangle = \frac{1}{\sqrt{N!}}b^{\dagger N}|0\rangle. \quad (7.17)$$

## 7.3 The Algebras $u(2)$ and $su(2)$

In this case, the index  $\alpha = 1, 2$ . The algebra  $u(2)$  has four elements

$$g \doteq b_1^\dagger b_1, b_1^\dagger b_2, b_2^\dagger b_1, b_2^\dagger b_2. \quad (7.18)$$

Beginning with  $u(2)$ , the study of the algebraic structure of  $u(n)$  requires several steps:

1. The enumeration of all possible subalgebra chains and their branchings
2. The construction of the basis  $\mathcal{B}$  for all chains
3. The construction of all invariant operators and their eigenvalues

### 7.3.1 Subalgebra Chains

There are two possible subalgebra chains:

*Subalgebra I:*  $u(2) \supset u(1)$

This is the trivial Abelian subalgebra  $u(1)$

$$g' \doteq b_1^\dagger b_1, \quad (7.19)$$

leading to the canonical chain

$$\left| \begin{array}{c} u(2) \supset u(1) \\ \downarrow \quad \downarrow \\ N \quad n_1 \end{array} \right\rangle, \quad (7.20)$$

with Gel'fand pattern

$$\left\langle \begin{array}{cc} N & 0 \\ & n_1 \end{array} \right\rangle. \quad (7.21)$$

Here the labels of the representations are written under the algebras. Two notations are often used for canonical chains: a bra-ket notation in which only non-zero quantum numbers are displayed, and the Gel'fand notation of Chap. 4 where all quantum numbers, including zeros, are displayed. The zero in the latter notation arises from the fact that only symmetric representations can be constructed with boson operators. Using the rules of Chap. 4, one can find easily the branching

$$n_1 = 0, 1, \dots, N. \quad (7.22)$$

The operator

$$\hat{N} = b_1^\dagger b_1 + b_2^\dagger b_2 = \hat{n}_1 + \hat{n}_2 \quad (7.23)$$

commutes with all elements of the algebra  $g$  and thus is an invariant operator of  $u(2)$

$$C_1(u(2)) = \hat{N}. \quad (7.24)$$

Powers of  $\hat{N}$  also commute with all elements of  $g$  and are thus also invariant operators. The quadratic Casimir operator is usually defined as

$$C_2(u(2)) = \hat{N}(\hat{N} + 2). \quad (7.25)$$

The basis is

$$\mathcal{B} : \frac{1}{\sqrt{n_1!n_2!}} (b_2^\dagger)^{n_2} (b_1^\dagger)^{n_1} |0\rangle \equiv |n_1, n_2\rangle. \quad (7.26)$$

Since  $n_2 = N - n_1$ , this can be rewritten as

$$\frac{1}{\sqrt{n_1!(N - n_1)!}} (b_2^\dagger)^{N - n_1} (b_1^\dagger)^{n_1} |0\rangle. \quad (7.27)$$

The dimension of the irreducible representations is  $N + 1$ .

**Example 2.** *The two dimensional harmonic oscillator in quantum mechanics*

The algebra  $u(2)$  can be used to describe the two-dimensional harmonic oscillator in (dimensionless) Cartesian coordinates,  $x$  and  $y$ . The creation and annihilation operators are related to coordinates and momenta by

$$\begin{aligned} b_1 &= \frac{1}{\sqrt{2}}(x + ip_x), & b_2^\dagger &= \frac{1}{\sqrt{2}}(x - ip_x) \\ b_2 &= \frac{1}{\sqrt{2}}(y + ip_y), & b_1^\dagger &= \frac{1}{\sqrt{2}}(y - ip_y). \end{aligned} \quad (7.28)$$

The Hamiltonian of the isotropic oscillator is

$$H = \frac{1}{2}(p_x^2 + x^2 + p_y^2 + y^2). \quad (7.29)$$

It can be rewritten in terms of elements of  $u(2)$  as

$$H = \hat{N} + 1 \quad (7.30)$$

with eigenvalues

$$E(N) = N + 1, \quad N = 0, 1, \dots \quad (7.31)$$

Because of these properties, the algebra  $u(2)$  is called the degeneracy algebra of the two dimensional harmonic oscillator.

*Subalgebra II:  $u(2) \supset so(2)$*

It is convenient to introduce the operators

$$\underbrace{\hat{F}_+ = b_1^\dagger b_2, \hat{F}_- = b_2^\dagger b_1, \hat{F}_z = \frac{1}{2}(b_2^\dagger b_2 - b_1^\dagger b_1)}_{su(2)}, \quad \hat{N} = b_2^\dagger b_2 + b_1^\dagger b_1. \quad (7.32)$$

The operators  $\hat{F}_+, \hat{F}_-, \hat{F}_z$  satisfy the commutation relations of  $su(2)$  in the Cartan-Weyl form

$$[\hat{F}_+, \hat{F}_-] = 2\hat{F}_z; \quad [\hat{F}_z, \hat{F}_\pm] = \pm \hat{F}_\pm. \quad (7.33)$$

To make connection with the usual form, introduce

$$\hat{F}_x = \frac{1}{2}[b_1^\dagger b_2 + b_2^\dagger b_1]; \quad \hat{F}_y = \frac{1}{2i}[b_1^\dagger b_2 - b_2^\dagger b_1]. \quad (7.34)$$

The operators  $\hat{F}_x, \hat{F}_y, \hat{F}_z$  satisfy the usual commutation relations

$$[\hat{F}_x, \hat{F}_y] = i\hat{F}_z; \quad [\hat{F}_y, \hat{F}_z] = i\hat{F}_x; \quad [\hat{F}_z, \hat{F}_x] = i\hat{F}_y. \quad (7.35)$$

The invariant Casimir operator is

$$\hat{F}^2 = \hat{F}_x^2 + \hat{F}_y^2 + \hat{F}_z^2 = \frac{1}{2}(\hat{F}_+ \hat{F}_- + \hat{F}_- \hat{F}_+) + \hat{F}_z^2 = \hat{F}_- \hat{F}_+ + \hat{F}_z(\hat{F}_z + 1). \quad (7.36)$$

This operator commutes with all elements

$$[\hat{F}^2, \hat{F}_\pm] = [\hat{F}^2, \hat{F}_z] = 0 \quad (7.37)$$

and thus

$$\hat{F}^2 = C_2(su(2)) . \tag{7.38}$$

When written in terms of boson operators

$$\hat{F}^2 = \frac{1}{4} \hat{N}(\hat{N} + 2) = \frac{1}{4} (b_2^\dagger b_2 + b_1^\dagger b_1)(b_2^\dagger b_2 + b_1^\dagger b_1 + 2) . \tag{7.39}$$

The invariant  $\hat{F}^2$  is related to that introduced previously in (7.25) by multiplication by  $\frac{1}{4}$ . This factor is due to the definition of the operators  $\hat{F}$ .

The basis is written in bra-ket notation as

$$\left| \begin{array}{cc} u(2) \supset so(2) \\ \downarrow \quad \downarrow \\ N \quad M \end{array} \right\rangle . \tag{7.40}$$

The intermediate step  $su(2)$  in the chain  $u(2) \supset su(2) \supset so(2)$  may or may not be written down, since, for totally symmetric (bosonic) representations, no additional label (quantum number) is required when going from  $u(n)$  to  $su(n)$ . Since one goes from a unitary to an orthogonal algebra, the chain  $u(2) \supset so(2)$  is non-canonical. The rules of Chap. 4 give the branching

$$M = \pm N, \pm(N - 2), \dots, \pm 1 \text{ or } 0, N = \text{odd or even} . \tag{7.41}$$

Note the  $\pm$  sign, which arises from the fact that  $so(2)$  is an orthogonal algebra in an even number of dimensions. The dimension of the representation is  $N + 1$  (as in Chain I). The basis can be converted to the standard form by introducing the quantum numbers

$$F = \frac{N}{2}, \quad F_z = \frac{M}{2} . \tag{7.42}$$

In this form the branching is the familiar quantum mechanics result

$$F_z = -F, -F + 1, \dots, F - 1, F . \tag{7.43}$$

The dimension of the representation is  $(2F + 1)$ . The basis is written as

$$\left| \begin{array}{cc} su(2) \supset so(2) \\ \downarrow \quad \downarrow \\ F \quad F_z \end{array} \right\rangle \tag{7.44}$$

and can be constructed with boson operators as

$$|F, F_z\rangle = \sqrt{\frac{1}{(F + F_z)!(F - F_z)!}} (b_2^\dagger)^{F + F_z} (b_1^\dagger)^{F - F_z} |0\rangle . \tag{7.45}$$

This construction of  $su(2)$  is called the Jordan-Schwinger construction. [J. Schwinger, in *Quantum Theory of Angular Momentum*, L.C. Biedenharn and H. van Dam, eds., Academic Press, New York, 1965, p. 229].

Note that, since  $u(1) \sim so(2)$ , there is no difference between subalgebra chains I and II. They can be converted into each other by

$$\begin{aligned} \hat{F}_z &= \frac{1}{2}(\hat{n}_2 - \hat{n}_1) = \frac{\hat{N}}{2} - \hat{n}_1; & F_z &= \frac{N}{2} - n_1 \\ F_z &= \frac{N}{2}, \frac{N}{2} - 1, \dots, -\frac{N}{2} = F, F - 1, \dots, -F. \end{aligned} \quad (7.46)$$

The situation is summarized in the graph

$$\begin{array}{c} u(2) \\ | \\ u(1) \sim so(2) \end{array} \quad (7.47)$$

called a lattice of algebras, a concept which will become clear in the following sections.

The algebras  $u(2)$  and  $su(2)$  constructed with boson operators play a crucial role in applications to problems in physics: (i)  $u(2)$  is the degeneracy algebra of the two dimensional harmonic oscillator; (ii)  $su(2) \sim so(3)$  is the angular momentum algebra.

## 7.4 The Algebras $u(n), n \geq 3$

Starting with  $u(3)$ , the possibility arises to have the angular momentum algebra as subalgebra of  $u(n)$ . For applications to problems with rotational invariance, it is convenient to introduce another form of the Lie algebra  $u(n)$ , called the Racah form. This is obtained from the Lie algebra of (7.2) by a change of basis. The Racah form can be constructed by introducing boson operators that transform as tensor operators under  $so(3)$ . Since this is a generic method, it is of interest to provide a general definition.

### 7.4.1 Racah Form

Introduce boson creation  $b_{l,m}^\dagger$  and annihilation  $b_{l,m}$  operators satisfying commutation relations

$$\begin{aligned} [b_{l,m}, b_{l',m'}^\dagger] &= \delta_{ll'} \delta_{mm'} \\ [b_{l,m}, b_{l',m'}] &= [b_{l,m}^\dagger, b_{l',m'}^\dagger] = 0 \end{aligned} \quad (7.48)$$

If the operators  $b_{l,m}^\dagger$  satisfy the commutation relations (6.5) with the elements of the algebra  $so(3)$ , the explicit expression of which is defined below, they transform as spherical tensors of rank  $l$  under  $so(3)$  rotations. In order to construct adjoint operators that transform as spherical tensors under  $so(3)$ , one must use the results of Chap. 6, Example 9. The operators  $b_{l,m}^\dagger$  and  $\tilde{b}_{l,m}$ ,

$$b_{l,m}^\dagger; \quad \tilde{b}_{l,m} = (-1)^{l-m} b_{l,-m} \quad (7.49)$$

transform as spherical tensors.

The coupled bilinear products

$$\begin{aligned} G_\kappa^{(k)}(l, l') &\equiv \left[ b_l^\dagger \times \tilde{b}_{l'} \right]_\kappa^{(k)} \\ &= \sum_{m, m'} \langle l m l' m' | k \kappa \rangle b_{l,m}^\dagger \tilde{b}_{l',m'} \end{aligned} \quad (7.50)$$

with  $|l + l'| \geq k \geq |l - l'|$  generate the Lie algebra  $u(n)$ . Here  $n = \sum_i (2l_i + 1)$ , where  $l_i$  are the values of  $l$  used in the construction of the Lie algebra. The commutation relations of the operators  $G_\kappa^k$  are

$$\begin{aligned} \left[ G_\kappa^{(k)}(l, l'), G_{\kappa'}^{(k')}(l'', l''') \right] &= \sum_{\kappa'', \kappa'''} (2k + 1)^{1/2} (2k' + 1)^{1/2} \langle k \kappa \kappa' \kappa'' | k'' \kappa'' \rangle \\ &\quad \times \left[ (-)^{l+l'''+k''} \left\{ \begin{matrix} l' & l''' & k' \\ k'' & k & l \end{matrix} \right\} \delta_{l'l''} G_{\kappa''}^{(k'')} (l, l''') \right. \\ &\quad \left. - (-)^{l'+l''+k+k'} \left\{ \begin{matrix} l'' & l & k' \\ k & k'' & l' \end{matrix} \right\} \delta_{l''l'''} G_{\kappa'''}^{(k''')} (l'', l') \right]. \end{aligned} \quad (7.51)$$

This form of the algebra and of the commutation relations is known as Racah's form. [G. Racah, *Group Theory and Spectroscopy*, Lecture Notes in Physics, Princeton University, 1951.]

### 7.4.2 Tensor Coupled Form of the Commutators

In deriving the preceding formula (7.51) and the corresponding formula for fermions, discussed in Chap. 8, it is useful to introduce a tensor coupled form of the commutators of the Lie algebras  $u(n)$ . The tensor commutator of two tensor operators  $G^{(e)}$  and  $G^{(f)}$  is defined by

$$\left[ G^{(e)}, G^{(f)} \right]_\xi^{(g)} = \sum_{\varepsilon, \varphi} \langle e \varepsilon f \varphi | g \xi \rangle \left[ G_\varepsilon^{(e)}, G_\varphi^{(f)} \right]. \quad (7.52)$$

This  $\left[ G^{(e)}, G^{(f)} \right]^{(g)}$  is itself a tensor operator and, at the tensor level, can be written as

$$\left[ G^{(e)}, G^{(f)} \right]^{(g)} = (G^{(e)} \times G^{(f)})^{(g)} - (-)^{g-e-f} (G^{(f)} \times G^{(e)})^{(g)}. \quad (7.53)$$

The tensor coupled form of the commutator was introduced in [J.B. French, *Multipole and Sum-rule Methods in Spectroscopy in Proceedings of the International School of Physics "Enrico Fermi"*, Course XXXVI, C. Bloch ed.,

Academic Press, New York, 1966, p. 278] and developed in [J.-Q. Chen, B.-Q. Chen, and A. Klein, *Factorization of Commutators: The Wick Theorem for Coupled Operators*, Nucl. Phys. A554, 61 (1993)].

The commutator formula for the generators of  $u(n)$  can be written as

$$\begin{aligned} \left[ G^{(e)}(a, b), G^{(f)}(c, d) \right]_{\xi}^{(g)} &= (-)^{a+d+g} (-)^{2a+2d+2g} \hat{e}\hat{f} \left\{ \begin{matrix} b & d & f \\ g & e & a \end{matrix} \right\} \delta_{bc} G_{\xi}^{(g)}(a, d) \\ &\quad - (-)^{b+c+e+f} (-)^{2c+2d+2g} \hat{e}\hat{f} \left\{ \begin{matrix} c & a & f \\ e & g & b \end{matrix} \right\} \delta_{ad} G_{\xi}^{(g)}(c, b), \end{aligned} \tag{7.54}$$

where  $\hat{j} = (2j + 1)^{1/2}$ , valid for  $a, b, c$  and  $d$  all integer (bosonic) or all half-integer (fermionic). For the bosonic case  $(-)^{2a+2d+2g} = (-)^{2c+2d+2g} = +1$ , while for the fermionic case, to be discussed in Chap. 8,  $(-)^{2a+2d+2g} = (-)^{2c+2d+2g} = -1$ . The uncoupled commutators (7.51) can be obtained from (7.52) by multiplying by the Clebsch-Gordan coefficient of  $so(3)$ ,  $\langle e\varepsilon f\varphi \mid g\xi \rangle$ , and summing over  $g$  and  $\xi$

$$\left[ G_{\varepsilon}^{(e)}, G_{\varphi}^{(f)} \right] = \sum_{g, \xi} \langle e\varepsilon f\varphi \mid g\xi \rangle \left[ G^{(e)}, G^{(f)} \right]_{\xi}^{(g)}. \tag{7.55}$$

### 7.4.3 Subalgebra Chains Containing $so(3)$

A generic subalgebra chain (called a classification scheme) for  $u(2l + 1)$  constructed with boson operators can be obtained as follows:

- (i) Exclude the element with  $k = 0, \kappa = 0$ ; this gives  $su(2l + 1)$ .
- (ii) Retain only elements with  $k = \text{odd}$ ; this gives the Lie algebra  $so(2l + 1)$ .
- (iii) Retain the elements with  $k = 1$ ; this gives the algebra  $so(3)$ .
- (iv) Retain the element with  $k = 1, \kappa = 0$ ; this gives the algebra  $so(2)$ .

A generic subalgebra chain for a single value of  $l$  is

$$\begin{aligned} u(2l + 1) &\supset su(2l + 1) \supset so(2l + 1) \\ &\supset \dots \supset so(3) \supset so(2). \end{aligned} \tag{7.56}$$

Dots have been inserted between  $so(2l + 1)$  and  $so(3)$  since, for large  $l$ , there may be intermediate steps.

## 7.5 The Algebras $u(3)$ and $su(3)$

To construct these algebras, one needs three boson operators  $b_{\alpha} (\alpha = 1, 2, 3)$ . The 9 elements of  $u(3)$  are

$$g \doteq b_{\alpha}^{\dagger} b_{\beta} \quad (\alpha, \beta = 1, 2, 3). \tag{7.57}$$

The algebraic analysis of  $u(3)$  encounters a new feature, namely that there are here two subalgebra chains that are distinct (non-isomorphic), in contrast with the previous case in which the subalgebra chains were isomorphic. While the first of these chains, the canonical chain, can be studied as in Sect. 3, the second chain, which includes the angular momentum algebra  $so(3)$  as a subalgebra, is best studied by introducing Racah form. In addition, several constructions of the same chain are possible. We begin with the constructions mostly used in quantum mechanical applications.

### 7.5.1 Subalgebra Chains

#### The Canonical Chain

*Subalgebra I:*  $u(3) \supset u(2) \supset u(1)$

The canonical chain can be constructed by introducing three (singlet) boson operators  $b_1, b_2, b_3$ . The elements of the algebras in the chain are

$$\begin{array}{ccc}
 u(3) & u(2) & u(1) \\
 b_1^\dagger b_1 & b_1^\dagger b_1 & b_1^\dagger b_1 \\
 b_1^\dagger b_2, b_2^\dagger b_1 & b_1^\dagger b_2, b_2^\dagger b_1 & \\
 b_1^\dagger b_3, b_3^\dagger b_1 & & \\
 b_2^\dagger b_2 & b_2^\dagger b_2 & \\
 b_2^\dagger b_3, b_3^\dagger b_2 & & \\
 b_3^\dagger b_3 & & .
 \end{array} \tag{7.58}$$

The elements of  $u(2)$  and  $u(1)$  are obtained from  $u(3)$  by deleting successively the boson operators  $b_3$  and  $b_2$ . The basis is

$$\mathcal{B} : \frac{1}{\sqrt{n_1!n_2!n_3!}} (b_1^\dagger)^{n_1} (b_2^\dagger)^{n_2} (b_3^\dagger)^{n_3} |0\rangle \equiv |n_1, n_2, n_3\rangle , \tag{7.59}$$

with

$$n_1 + n_2 + n_3 = N . \tag{7.60}$$

The labels can be arranged in the usual Dirac ket-notation

$$\left| \begin{array}{ccc}
 u(3) \supset u(2) \supset u(1) \\
 \downarrow \quad \downarrow \quad \downarrow \\
 N \quad n \quad n_1
 \end{array} \right\rangle , \tag{7.61}$$

with branching

$$\begin{aligned}
 n &= 0, \dots, N, \\
 n_1 &= 0, \dots, n .
 \end{aligned} \tag{7.62}$$

The Gel'fand pattern is



$$\left\langle \begin{array}{ccc} N & 0 & 0 \\ & n & 0 \\ & & n_1 \end{array} \right\rangle \equiv \left\langle \begin{array}{ccc} n_1 + n_2 + n_3 & 0 & 0 \\ & n_1 + n_2 & 0 \\ & & n_1 \end{array} \right\rangle. \quad (7.63)$$

The algebra has a linear invariant

$$C_1(u(3)) = \hat{N} = b_1^\dagger b_1 + b_2^\dagger b_2 + b_3^\dagger b_3, \quad (7.64)$$

and higher order invariants,  $C_2(u(3))$  and  $C_3(u(3))$ , which are combinations of powers of  $\hat{N}$ .

**Example 3.** *The harmonic oscillator in three dimensions in Cartesian coordinates*

The algebra of  $u(3)$  and its canonical chain describe the harmonic oscillator in three dimensions and in Cartesian coordinates. The dimensionless coordinates,  $x, y, z$ , and momenta,  $p_x, p_y, p_z$ , are related to the creation and annihilation operators by

$$\begin{aligned} b_1 &= \frac{1}{\sqrt{2}}(x + ip_x), & b_1^\dagger &= \frac{1}{\sqrt{2}}(x - ip_x) \\ b_2 &= \frac{1}{\sqrt{2}}(y + ip_y), & b_2^\dagger &= \frac{1}{\sqrt{2}}(y - ip_y) \\ b_3 &= \frac{1}{\sqrt{2}}(z + ip_z), & b_3^\dagger &= \frac{1}{\sqrt{2}}(z - ip_z). \end{aligned} \quad (7.65)$$

The Hamiltonian of the isotropic oscillator is

$$H = \frac{1}{2}(p_x^2 + p_y^2 + p_z^2 + x^2 + y^2 + z^2). \quad (7.66)$$

Its algebraic form is

$$H = \sum_{i=1}^3 b_i^\dagger b_i + \frac{3}{2} = \hat{N} + \frac{3}{2}, \quad (7.67)$$

with eigenvalues

$$E(N) = N + \frac{3}{2}, \quad N = 0, 1, \dots. \quad (7.68)$$

The eigenvalues depend only on the quantum number  $N = n_3 + n_2 + n_1$  and not on the other quantum numbers  $n = n_2 + n_1$  and  $n_1$ . The canonical chain is also useful to describe the anisotropic oscillator, with Hamiltonian

$$H = \frac{1}{2}(p_x^2 + p_y^2 + p_z^2 + \omega_x^2 x^2 + \omega_y^2 y^2 + \omega_z^2 z^2) \quad (7.69)$$

and eigenvalues

$$\begin{aligned} E(n_1, n_2, n_3) &= \omega_x n_1 + \omega_y n_2 + \omega_z n_3 + \frac{3}{2}, \\ n_1, n_2, n_3 &= 0, 1, \dots \end{aligned} \quad (7.70)$$

The algebra  $u(3)$  is the degeneracy algebra of the three dimensional harmonic oscillator.

**The Non-Canonical Chain and its Racah Form**

*Subalgebra II:*  $u(3) \supset so(3) \supset so(2)$

In order to construct the non-canonical chain  $u(3) \supset so(3) \supset so(2)$ , it is convenient to introduce spherical boson operators (tensor operators with respect to  $so(3)$ )

$$p_{\pm 1}^\dagger = \mp \frac{1}{\sqrt{2}} (b_1^\dagger \pm i b_2^\dagger), \quad p_0^\dagger = b_3^\dagger \tag{7.71}$$

denoted by  $p_\mu^\dagger (\mu = 0, \pm 1)$  and their adjoint  $\tilde{p}_\mu = (-1)^{1-\mu} p_{-\mu}$ . The three boson operators  $p_\mu^\dagger (\mu = \pm 1, 0)$  transform as the representation  $l = 1$  of  $so(3)$ . In the generic construction of the elements of the algebra given in the preceding section, there is thus only one value of  $l$ .

The elements of  $u(3)$  in Racah form are

$$\begin{aligned} G_0^{(0)}(pp) &= (p^\dagger \times \tilde{p})_0^{(0)} \quad 1 \\ G_\kappa^{(1)}(pp) &= (p^\dagger \times \tilde{p})_\kappa^{(1)} \quad 3 \\ G_\kappa^{(2)}(pp) &= (p^\dagger \times \tilde{p})_\kappa^{(2)} \quad 5 \end{aligned} \tag{7.72}$$

for a total of nine elements. The number of elements  $(2k + 1)$  is written next to them.

Subalgebras of  $u(3)$  can be then constructed as follows: (i) Deleting the element  $G_0^{(0)}$  gives  $su(3)$ . (ii) Keeping only the elements with  $k = 1$  gives  $so(3)$ . (iii) Keeping only  $G_0^{(1)}$  gives  $so(2)$ .

$$\begin{array}{cccc} u(3) & su(3) & so(3) & so(2) \\ \\ (p^\dagger \times \tilde{p})_0^{(0)} & & & \\ (p^\dagger \times \tilde{p})_\kappa^{(1)} & (p^\dagger \times \tilde{p})_\kappa^{(1)} & (p^\dagger \times \tilde{p})_\kappa^{(1)} & (p^\dagger \times \tilde{p})_0^{(1)} \\ (p^\dagger \times \tilde{p})_0^{(2)} & (p^\dagger \times \tilde{p})_0^{(2)} & & \end{array} \tag{7.73}$$

The basis can be written as

$$\left| \begin{array}{ccc} u(3) \supset so(3) \supset so(2) \\ \downarrow \quad \downarrow \quad \downarrow \\ N \quad L \quad M_L \end{array} \right\rangle . \tag{7.74}$$

Using the rules of Chap. 4, one obtains the branching

$$\begin{array}{l} L = N, N - 2, \dots, 1 \text{ or } 0 \quad (N = \text{odd or even}) \\ M_L = -L, \dots, +L \end{array} . \tag{7.75}$$

Again the intermediate step  $su(3)$  in  $u(3) \supset su(3) \supset so(3) \supset so(2)$  may or may not be written down, since for symmetric representations no new quantum number is needed when going from  $u(3)$  to  $su(3)$ .

**Example 4.** *The three dimensional harmonic oscillator in spherical coordinates*

The chain  $u(3) \supset so(3) \supset so(2)$  can be used to describe the harmonic oscillator in spherical coordinates,  $r, \vartheta, \varphi$ . These coordinates are related to the Cartesian coordinates by the familiar relations  $x = r \cos \varphi \sin \vartheta$ ,  $y = r \sin \varphi \sin \vartheta$ ,  $z = r \cos \vartheta$ . The Hamiltonian is still given by (7.66), written now as

$$H = \frac{1}{2} (p^2 + r^2) . \tag{7.76}$$

The algebraic form of this Hamiltonian is still

$$H = \hat{N} + \frac{3}{2} , \tag{7.77}$$

with eigenvalues

$$E(N) = N + \frac{3}{2}, \quad N = 0, 1, \dots \tag{7.78}$$

The eigenstates, given in (7.74), can be written in the form

$$|N, L, M_L\rangle = \frac{1}{\mathcal{N}} (p^\dagger)_{L, M_L}^N |0\rangle \tag{7.79}$$

where the product of  $N$  operators  $p^\dagger$  is coupled to  $L$  and  $M_L$ , and  $\mathcal{N}$  is a normalization constant.

### 7.5.2 Lattice of Algebras

The situation for the branchings of  $u(3)$  can be summarized in the following lattice of algebras

$$\begin{array}{ccc}
 & u(3) & \\
 \swarrow & & \searrow \\
 u(2) & & so(3) \\
 \swarrow & & \searrow \\
 & u(1) \sim so(2) & 
 \end{array}
 \quad
 \begin{array}{ccc}
 & A_2 & \\
 \swarrow & & \searrow \\
 A_1 & & B_1 \\
 \swarrow & & \searrow \\
 & D_1 & 
 \end{array}
 , \tag{7.80}$$

where Cartan's notation is used on the right-hand side. The lattice must be read from top to bottom.

### 7.5.3 Boson Calculus of $u(3) \supset so(3)$

Spherical boson operators are extensively used in physics. It is therefore of interest to develop a boson calculus, for manipulations of tensor operators with respect to  $so(3)$ . The basic commutation relations are

$$[p_\mu, p_{\mu'}^\dagger] = \delta_{\mu \mu'} \quad [\tilde{p}_\mu, p_{\mu'}^\dagger] = (-1)^{1-\mu} \delta_{\mu', -\mu} . \tag{7.81}$$

The elements of the Lie algebra  $u(3)$  can be explicitly constructed using (7.50). The element  $[p^\dagger \times \tilde{p}]_0^{(0)}$  is

$$\begin{aligned} [p^\dagger \times \tilde{p}]_0^{(0)} &= \sum_{\mu, \mu'} \langle 1 \mu \ 1 \mu' \mid 00 \rangle p_\mu^\dagger \tilde{p}_{\mu'} \\ &= \sum_{\mu, \mu'} (-1)^{1-\mu} \frac{1}{\sqrt{3}} \delta_{\mu, -\mu'} p_\mu^\dagger \tilde{p}_{\mu'} = \sum_{\mu} (-1)^{1-\mu} \frac{1}{\sqrt{3}} p_\mu^\dagger \tilde{p}_{-\mu} \\ &= \frac{1}{\sqrt{3}} \sum_{\mu} p_\mu^\dagger p_\mu = \frac{1}{\sqrt{3}} \hat{N}. \end{aligned} \quad (7.82)$$

This element is thus proportional to the number operator  $\hat{N}$  that counts the number of bosons. The other elements are

$$[p^\dagger \times \tilde{p}]_\kappa^{(k)} = \sum_{\mu, \mu'} \langle 1\mu 1\mu' \mid k\kappa \rangle p_\mu^\dagger \tilde{p}_{\mu'}. \quad (7.83)$$

An important ingredient is often the tensor product of two creation or annihilation operators. The tensor product of two creation or annihilation operators can only have even rank,  $\lambda = \text{even}$ .

*Proof.* Consider the product

$$\begin{aligned} [p^\dagger \times p^\dagger]_\mu^{(\lambda)} &= \sum_{\mu_1 \mu_2} \langle 1\mu_1 \ 1\mu_2 \mid \lambda\mu \rangle p_{\mu_1}^\dagger p_{\mu_2}^\dagger \\ &= \sum_{\mu_1 \mu_2} \langle 1\mu_1 \ 1\mu_2 \mid \lambda\mu \rangle p_{\mu_2}^\dagger p_{\mu_1}^\dagger = \sum_{\mu_1 \mu_2} \langle 1\mu_2 \ 1\mu_1 \mid \lambda\mu \rangle p_{\mu_1}^\dagger p_{\mu_2}^\dagger \\ &= \sum_{\mu_1 \mu_2} (-1)^{2-\lambda} \langle 1\mu_1 \ 1\mu_2 \mid \lambda\mu \rangle p_{\mu_1}^\dagger p_{\mu_2}^\dagger. \end{aligned} \quad (7.84)$$

Equating the first and last term one finds

$$(-1)^{2-\lambda} = 1; \quad \lambda = \text{even} = 0, 2. \quad (7.85)$$

The elements of the algebra  $u(3) \supset so(3)$  are often denoted by

$$\begin{aligned} \hat{N} &= \sqrt{3}[p^\dagger \times \tilde{p}]_0^{(0)} \\ \hat{L}_\kappa &= \sqrt{2}[p^\dagger \times \tilde{p}]_\kappa^{(1)}. \\ \hat{Q}_\kappa &= [p^\dagger \times \tilde{p}]_\kappa^{(2)}. \end{aligned} \quad (7.86)$$

In applications to the harmonic oscillator problem in quantum mechanics,  $\hat{N}$  represents the number operator that counts the oscillator quanta. The explicit expression for the components  $\hat{L}_\kappa$  is

$$\begin{aligned}
\hat{L}_0 &= \left( p_{+1}^\dagger \tilde{p}_{-1} - p_{-1}^\dagger \tilde{p}_{+1} \right), \\
\hat{L}_{-1} &= \left( p_1^\dagger \tilde{p}_0 - p_0^\dagger \tilde{p}_{+1} \right), \\
\hat{L}_{+1} &= \left( p_0^\dagger \tilde{p}_{+1} - p_{+1}^\dagger \tilde{p}_0 \right).
\end{aligned} \tag{7.87}$$

With the normalization (7.86), the three components  $\hat{L}_\kappa$  of the operator  $\hat{L}$  satisfy the commutation relations of angular momentum

$$[\hat{L}_0, \hat{L}_{\pm 1}] = \pm \hat{L}_{\pm 1}, \quad [\hat{L}_{-1}, \hat{L}_{+1}] = \hat{L}_0. \tag{7.88}$$

*Proof.* The commutator  $[\hat{L}_{-1}, \hat{L}_{+1}]$  is

$$\begin{aligned}
2[(p^\dagger \times \tilde{p})_{-1}^{(1)}, (p^\dagger \times \tilde{p})_{+1}^{(1)}] &= -[p_{-1}^\dagger \tilde{p}_0 + p_0^\dagger \tilde{p}_{-1}, p_0^\dagger \tilde{p}_{+1} + p_{+1}^\dagger \tilde{p}_0] \\
&= -(p_{-1}^\dagger \tilde{p}_{+1} - p_{+1}^\dagger \tilde{p}_{-1}) \\
&= \sqrt{2}(p^\dagger \times \tilde{p})_0^{(1)} = \hat{L}_0.
\end{aligned} \tag{7.89}$$

The last operator,  $\hat{Q}$ , is a tensor operator of rank 2 (quadrupole tensor).

#### 7.5.4 Matrix Elements of Operators in $u(3) \supset so(3)$

Boson calculus is used to evaluate matrix elements of operators. The basis is written as

$$\mathcal{B}: \quad |N, L, M_L\rangle = \frac{1}{\mathcal{N}} (p^\dagger)_{L, M_L}^N |0\rangle, \tag{7.90}$$

where the  $N$  boson operators have been coupled to  $L, M_L$ . There are no missing labels here. An explicit expression in terms of solid harmonics in  $p^\dagger$  is also available. [O.S. van Roosmalen, *Algebraic Description of Nuclear and Molecular Rotation-Vibration Spectra*, Ph.D. Thesis, University of Groningen, The Netherlands, 1982.]

Reduced matrix elements of the boson operators in this basis are

$$\begin{aligned}
\langle N+1, L-1 \| p^\dagger \| N, L \rangle &= [(N-L+2)L]^{1/2} \\
\langle N+1, L+1 \| p^\dagger \| N, L \rangle &= [(N+L+3)(L+1)]^{1/2}.
\end{aligned} \tag{7.91}$$

The reduced matrix elements of the annihilation operators can be obtained by using the relation

$$\langle \alpha', L' \| \tilde{b}_l \| \alpha, L \rangle = (-1)^{L-L'+l} \langle \alpha, L \| b_l^\dagger \| \alpha', L' \rangle, \tag{7.92}$$

where  $\alpha$  denotes additional labels. The matrix elements of the elements of the algebra are

$$\langle N, L, M_L | \hat{N} | N, L, M_L \rangle = N \tag{7.93}$$

and

$$\begin{aligned} \langle N, L \parallel \hat{L} \parallel N, L \rangle &= [L(L+1)(2L+1)]^{1/2} \\ \langle N, L \parallel \hat{Q} \parallel N, L \rangle &= (2N+3) \left[ \frac{L(L+1)(2L+1)}{6(2L-1)(2L+3)} \right]^{1/2} \\ \langle N, L+2 \parallel \hat{Q} \parallel N, L \rangle &= \left[ \frac{(N-L)(N+L+3)(L+1)(L+2)}{(2L+3)} \right]^{1/2}. \end{aligned} \quad (7.94)$$

Matrix elements of polynomials in the elements of the algebra can be obtained by using the reduction formulas of Chap. 6.

### 7.5.5 Tensor Calculus of $u(3) \supset so(3)$

An alternative way to calculate matrix elements of operators is making use of tensor calculus. The three boson creation operators  $p_{\pm 1}^\dagger, p_0$  transform under  $u(3)$  as the 3-dimensional representation  $[1, 0, 0]$  (or  $(1, 0)$  of  $su(3)$ ). They also transform as the representation  $L = 1$  under  $so(3)$ , with component  $\kappa = \pm 1, 0$ . A commonly used notation for these tensors is

$$T_{1,\kappa}^{[1,0,0]} \equiv \square, \quad (7.95)$$

that is the creation operator transform as the fundamental representation of  $u(3)$ , often denoted as a *particle*. The boson annihilation operators  $\tilde{p}_{\pm 1}, \tilde{p}_0$  transform instead as the 3-dimensional conjugate representation  $[1, 1, 0]$  (or  $(1, 1)$  of  $su(3)$ ),

$$T_{1,\kappa}^{[1,1,0]} \equiv \begin{array}{c} \square \\ \square \end{array}. \quad (7.96)$$

This is often denoted as an *antiparticle* (or *hole*).

The elements of the algebra, that is the bilinear products  $(p^\dagger \times \tilde{p})_\kappa^{(k)}$ , can be simply obtained by taking tensor products, as in Chap. 4,

$$\square \otimes \begin{array}{c} \square \\ \square \end{array} = \begin{array}{cc} \square & \square \\ \square & \square \end{array} \oplus \begin{array}{c} \square \\ \square \end{array}. \quad (7.97)$$

The representations on the right hand side contain

$$\begin{array}{ccc} [2, 1, 0] & \oplus & [1, 1, 1] \\ \downarrow & & \downarrow \\ (p^\dagger \times \tilde{p})_\kappa^{(1)} & & (p^\dagger \times \tilde{p})_\kappa^{(0)} \end{array}. \quad (7.98)$$

$$(p^\dagger \times \tilde{p})_\kappa^{(2)}$$

The dimensions are:  $\dim [2, 1, 0] = 8, \dim [1, 1, 1] = 1$  for a total of 9 elements. Note that the elements of the algebra do not transform as its fundamental

representation. Under reduction of  $u(3)$  to  $su(3)$ , the representation  $[2, 1, 0]$  becomes  $(2, 1)$  with  $\dim(2, 1) = 8$ , while the representation  $[1, 1, 1]$  becomes  $(0, 0)$  with  $\dim(0, 0) = 1$ .

The calculation of matrix elements of a generic tensor  $T$

$$\left\langle [n', 0, 0], L', M'_L \mid T_{k, \kappa}^{[n_1, n_2, n_3]} \mid [n, 0, 0], L, M_L \right\rangle \quad (7.99)$$

is done using the techniques of Chap. 6.

The algebra  $u(3)$  (and its subalgebra  $su(3)$ ) constructed with boson operators occupy a special role in physics since  $u(3)$  is the degeneracy algebra of the three-dimensional harmonic oscillator.

### 7.5.6 Other Boson Constructions of $u(3)$

In addition to the construction in terms of boson operators  $b_1, b_2, b_3$  (canonical chain) and in terms of vector bosons,  $p_\mu (\mu = \pm 1, 0)$ , there is another boson construction of  $u(3)$  of practical interest. This construction is in terms of a singlet boson,  $\sigma$ , and a doublet  $\tau_x, \tau_y$ . The doublet can be rewritten as

$$\begin{aligned} \tau_\pm &= \frac{1}{\sqrt{2}} (\tau_x \pm i\tau_y) \\ \tau_\pm^\dagger &= \frac{1}{\sqrt{2}} (\tau_x^\dagger \mp i\tau_y^\dagger) . \end{aligned} \quad (7.100)$$

The boson operators  $\tau_\pm$  are called circular boson operators. The algebra  $u(3)$  is composed of nine elements:

$$\begin{aligned} \hat{n} &= (\tau_+^\dagger \tau_+ + \tau_-^\dagger \tau_-) \\ \hat{l} &= (\tau_+^\dagger \tau_+ - \tau_-^\dagger \tau_-) \\ \hat{Q}_+ &= \sqrt{2} (\tau_+^\dagger \tau_-) \\ \hat{Q}_- &= \sqrt{2} (\tau_-^\dagger \tau_+) \\ \hat{n}_s &= (\sigma^\dagger \sigma) \\ \hat{D}_+ &= \sqrt{2} (\tau_+^\dagger \sigma - \sigma^\dagger \tau_-) \\ \hat{D}_- &= \sqrt{2} (-\tau_-^\dagger \sigma + \sigma^\dagger \tau_+) \\ \hat{R}_+ &= \sqrt{2} (\tau_+^\dagger \sigma + \sigma^\dagger \tau_-) \\ \hat{R}_- &= \sqrt{2} (\tau_-^\dagger \sigma + \sigma^\dagger \tau_+) . \end{aligned} \quad (7.101)$$

This construction has two possible subalgebra chains (as before).

*Subalgebra chain I:*  $u(3) \supset u(2) \supset so(2)$

The subalgebra  $u(2)$  is composed by the four operators

$$u(2) \doteq \hat{n}, \hat{l}, \hat{Q}_+, \hat{Q}_-. \quad (7.102)$$

The subalgebra  $so(2)$  is composed of a single operator

$$so(2) \doteq \hat{l}. \quad (7.103)$$

The basis states in this chain are characterized by the quantum numbers

$$\left| \begin{array}{ccc} u(3) \supset u(2) \supset so(2) \\ \downarrow \quad \downarrow \quad \downarrow \\ N \quad n \quad l \end{array} \right\rangle, \quad (7.104)$$

with branching rules

$$\begin{aligned} n &= N, N-1, \dots, 0; \\ l &= \pm n, \pm(n-2), \dots, \pm 1 \text{ or } 0 \quad (n = \text{odd or even}). \end{aligned} \quad (7.105)$$

*Subalgebra chain II:*  $u(3) \supset so(3) \supset so(2)$

The subalgebra  $so(3)$  is composed by the three operators

$$so(3) \doteq \hat{D}_+, \hat{D}_-, \hat{l}. \quad (7.106)$$

The subalgebra  $so(2)$  is composed by the single operator  $\hat{l}$  as in (7.101).

The basis states in this chain are characterized by the quantum numbers

$$\left| \begin{array}{ccc} u(3) \supset so(3) \supset so(2) \\ \downarrow \quad \downarrow \quad \downarrow \\ N \quad \omega \quad l \end{array} \right\rangle, \quad (7.107)$$

with branchings

$$\begin{aligned} \omega &= N, N-2, \dots, 1 \text{ or } 0 \quad (N = \text{odd or even}); \\ -\omega &\leq l \leq +\omega. \end{aligned} \quad (7.108)$$

In view of the isomorphism  $so(2) \sim u(1)$ , the lattice of algebras for this construction is identical to that given previously in Sect. 4.2.

The construction of  $u(3)$  in terms of a singlet and a doublet has practical applications in the study of vibration-rotation spectra of molecules in two-dimensions (bending vibrations). [F. Iachello and S. Oss, *Algebraic Approach to Molecular Spectra: Two-dimensional Problems*, J. Chem. Phys. 104, 6956 (1996).]

## 7.6 The Unitary Algebra $u(4)$

This algebra can be constructed by considering four boson operators,  $b_\alpha$ ,  $\alpha = 1, 2, 3, 4$ . The 16 elements of the algebra are

$$G_{\alpha\beta} \doteq b_\alpha^\dagger b_\beta. \quad (7.109)$$



### 7.6.1 Subalgebra Chains not Containing $so(3)$

#### The Canonical Chain

The canonical chain  $u(4) \supset u(3) \supset u(2) \supset u(1)$  can be constructed trivially as before in terms of four boson operators  $b_1, b_2, b_3, b_4$ . No more will be said about this chain.

#### The Doublet Chain

Another chain that can be constructed simply is that in which the boson operators are divided into two doublets,  $b_1, b_2$  and  $b_3, b_4$ . The bilinear products of creation and annihilation operators of each doublet generate a  $u(2)$  algebra (Jordan-Schwinger construction), Sect. 2. Denoting by  $u_1(2)$  and  $u_2(2)$  the two  $u(2)$  algebras, one has the chain  $u(4) \supset u_1(2) \oplus u_2(2)$ . From there on, one can use the results of Sect. 2. The complete subalgebra chain is  $u(4) \supset u_1(2) \oplus u_2(2) \supset so_1(2) \oplus so_2(2)$ .

The situation can be summarized in the lattice of algebras

$$\begin{array}{ccc}
 & u(4) & \\
 & / \quad \backslash & \\
 u(3) & & u_1(2) \oplus u_2(2) \\
 | & & | \\
 u(2) & & so_1(2) \oplus so_2(2) \\
 | & & \\
 u(1) & & 
 \end{array} . \tag{7.110}$$

### 7.6.2 Subalgebra Chains Containing $so(3)$

We consider here explicitly non-canonical chains that contain the angular momentum algebra  $so(3)$  as a subalgebra. These are particularly important in problems with rotational invariance and will be discussed here in detail. These chains can be constructed by introducing scalar,  $s^\dagger$ , and vector,  $p_\mu^\dagger$  ( $\mu = 0, \pm 1$ ), boson operators that transform as  $l = 0$  and  $l = 1$  under  $so(3)$ ,

$$\begin{aligned}
 l = 0 : & \quad s^\dagger \\
 l = 1 : & \quad p_\mu^\dagger \quad (\mu = 0, \pm 1) .
 \end{aligned} \tag{7.111}$$

(Although not necessary for the construction of the algebra, in applications in molecular physics the parity of these operators is chosen to be  $\mathcal{P} = (-1)^l$ .) The elements of  $u(4)$  are, in Racah form,

$$\begin{aligned}
 G_0^{(0)}(ss) &= (s^\dagger \times \tilde{s})_0^{(0)} \quad 1 \\
 G_0^{(0)}(pp) &= (p^\dagger \times \tilde{p})_0^{(0)} \quad 1 \\
 G_\kappa^{(1)}(pp) &= (p^\dagger \times \tilde{p})_\kappa^{(1)} \quad 3 \\
 G_\kappa^{(2)}(pp) &= (p^\dagger \times \tilde{p})_\kappa^{(2)} \quad 5 \\
 G_\kappa^{(1)}(ps) &= (p^\dagger \times \tilde{s})_\kappa^{(1)} \quad 3 \\
 G_\kappa^{(1)}(sp) &= (s^\dagger \times \tilde{p})_\kappa^{(1)} \quad 3 \quad , \tag{7.112}
 \end{aligned}$$

for a total of 16 elements. To this algebra the standard procedure of (i) constructing all possible subalgebra chains; (ii) constructing the Casimir invariants and their eigenvalues; (iii) constructing the basis  $\mathcal{B}$ ; is then applied.

### Subalgebras

There are two subalgebra chains that contain  $so(3)$ :

*Subalgebra I:*  $u(4) \supset u(3) \supset so(3) \supset so(2)$

The elements of the subalgebras and their numbers are

$$\begin{aligned}
 u(3) : & \\
 & (p^\dagger \times \tilde{p})_0^{(0)} \quad 1 \\
 & (p^\dagger \times \tilde{p})_\kappa^{(1)} \quad 3 \\
 & (p^\dagger \times \tilde{p})_\kappa^{(2)} \quad 5 \\
 so(3) : & \\
 & (p^\dagger \times \tilde{p})_\kappa^{(1)} \quad 3 \\
 so(2) : & \\
 & (p^\dagger \times \tilde{p})_0^{(1)} \quad 1
 \end{aligned} \tag{7.113}$$

*Subalgebra II:*  $u(4) \supset so(4) \supset so(3) \supset so(2)$

The elements of the subalgebras and their number are

$$\begin{aligned}
 so(4) : & \\
 & (p^\dagger \times \tilde{p})_\kappa^{(1)} \quad 3 \\
 & (p^\dagger \times \tilde{s})_\kappa^{(1)} + (s^\dagger \times \tilde{p})_\kappa^{(1)} \quad 3 \\
 so(3) : & \quad . \\
 & (p^\dagger \times \tilde{p})_\kappa^{(1)} \quad 3 \\
 so(2) : & \\
 & (p^\dagger \times \tilde{p})_0^{(1)} \quad 1
 \end{aligned} \tag{7.114}$$

For this subalgebra chain, a difficulty arises, as there are two  $so(4)$  algebras that can be constructed with the bilinear products of  $s$  and  $p$  boson operators. The second  $so(4)$  algebra, denoted by  $so(4)$  is composed of the following operators



The dot product denotes scalar products with respect to  $so(3)$ , defined in Chap. 6, Sect. 13. A short-hand notation, often used, is

$$\begin{aligned}
 \hat{n}_s &= (s^\dagger \times \tilde{s})_0^{(0)} \\
 \hat{n}_p &= \sqrt{3}(p^\dagger \times \tilde{p})_0^{(0)} \\
 \hat{L} &= \sqrt{2}(p^\dagger \times \tilde{p})_\kappa^{(1)} \\
 \hat{Q} &= (p^\dagger \times \tilde{p})_\kappa^{(2)} \\
 \hat{D} &= (p^\dagger \times \tilde{s} + s^\dagger \times \tilde{p})_\kappa^{(1)} \\
 \hat{D}' &= i(p^\dagger \times \tilde{s} - s^\dagger \times \tilde{p})_\kappa^{(1)}.
 \end{aligned} \tag{7.123}$$

In addition to the number operators  $\hat{n}_s$  and  $\hat{n}_p$ , there are the angular momentum operator  $\hat{L}$ , the quadrupole operator  $\hat{Q}$  and two (hermitian) dipole operators  $\hat{D}$  and  $\hat{D}'$ . (In quantum mechanical applications these operators are related to the coordinate and momentum vectors.) In the notation (7.123), the invariants take a simple and familiar form

$$C_2(so(3)) = \hat{L} \cdot \hat{L} \quad C_2(so(4)) = \hat{L} \cdot \hat{L} + \hat{D} \cdot \hat{D}. \tag{7.124}$$

### Branchings

A crucial problem for application of Lie algebraic methods to problems in physics and chemistry is the classification (or branching) problem. This problem is solved using the techniques developed in Chap. 4.

#### *Branching I*

The branching of the totally symmetric representations of  $u(4)$  for the chain  $u(4) \supset u(3) \supset so(3) \supset so(2)$  is

$$\begin{aligned}
 u(4) \ [f_1, f_2, f_3, f_4] \ [N] &= \overbrace{\square \dots \square}^{N\text{-times}} \equiv [N, 0, 0, 0] \\
 u(3) \ [f'_1, f'_2, f'_3] \ [n_p] &= \overbrace{\square \dots \square}^{n_p\text{-times}} \equiv [n_p, 0, 0] \\
 & \quad n_p = 0, 1, \dots, N \\
 so(3) \ (\mu'_1) & \quad (L) \\
 & \quad L = n_p, n_p - 2, \dots, 1 \text{ or } 0 \text{ (} N = \text{odd or even)} \\
 so(2) \ (\mu''_1) & \quad (M_L) \\
 & \quad M_L = -L, \dots, +L.
 \end{aligned} \tag{7.125}$$

The abstract characterization of the representations of the algebras  $u(n)$ ,  $so(n)$ , of Chap. 4, has been replaced here by quantum numbers with physical meaning:  $N$  is the total boson number,  $n_p$  is the number of  $p$  bosons,  $L$  is

the angular momentum and  $M_L$  is its  $z$ -component. This branching is in part canonical,  $u(4) \supset u(3)$  and  $so(3) \supset so(2)$  and in part non-canonical,  $u(3) \supset so(3)$ . The basis is labelled by the quantum numbers

$$\left| \begin{array}{cccc} u(4) \supset u(3) \supset so(3) \supset so(2) \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ N \quad n_p \quad L \quad M_L \end{array} \right\rangle. \quad (7.126)$$

*Branching II*

For the chain  $u(4) \supset so(4) \supset so(3) \supset so(2)$ , the branching is

$$\begin{aligned} u(4) \ [f_1, f_2, f_3, f_4] \ [N] &= \overbrace{\square \dots \square}^{N\text{-times}} \equiv [N, 0, 0, 0] \\ so(4) \ (\mu_1, \mu_2) \quad (\omega, 0) & \quad \omega = N, N - 2, \dots, 1 \text{ or } 0 \ (N = \text{odd or even}) \\ so(3) \ (\mu'_1) \quad (L) & \quad L = \omega, \omega - 1, \dots, 1, 0 \\ so(2) \ (\mu''_1) \quad (M_L) & \quad M_L = -L, \dots, +L \end{aligned} \quad (7.127)$$

The basis is labelled by the quantum numbers

$$\left| \begin{array}{cccc} u(4) \supset so(4) \supset so(3) \supset so(2) \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ N \quad \omega \quad L \quad M_L \end{array} \right\rangle. \quad (7.128)$$

The branching  $u(4) \supset so(4)$  is non-canonical while the branching  $so(4) \supset so(3) \supset so(2)$  is canonical.

**Eigenvalues of Casimir Operators**

The eigenvalues of the Casimir operators in the appropriate irreducible representations can be obtained using the rules of Chap. 5. For the unitary algebras  $u(4)$  and  $u(3)$  they are trivially given by

$$\begin{aligned} \langle [N] | C_1(u(4)) | [N] \rangle &= N \\ \langle [N] | C_2(u(4)) | [N] \rangle &= N(N + 3) \\ \langle [n_p] | C_1(u(3)) | [n_p] \rangle &= n_p \\ \langle [n_p] | C_2(u(3)) | [n_p] \rangle &= n_p(n_p + 2). \end{aligned} \quad (7.129)$$

For the orthogonal algebras  $so(4)$  and  $so(3)$  they are given by

$$\langle (\omega, 0) | C_2(so(4)) | (\omega, 0) \rangle = \omega(\omega + 2)$$

$$\langle(L) | C_2(so(3)) | (L)\rangle = L(L + 1) . \tag{7.130}$$

These eigenvalues differ from those of Table 5.1 by a factor of 2, due to the different definition of  $C$  given above. Also, note that the eigenvalues of the Casimir operators of an algebra  $g$  depend only on the labels of  $g$  and not on those of the subalgebra chain  $g \supset g' \supset g'' \supset \dots$

### Lattice of Algebras


The non-canonical chains discussed above can be depicted into a lattice of algebras

$$\begin{array}{ccc}
 & u(4) & \\
 & / \quad \backslash & \\
 u(3) & & so(4) \\
 & \backslash \quad / & \\
 & so(3) & \\
 & | & \\
 & so(2) & 
 \end{array}
 \qquad
 \begin{array}{ccc}
 & A_3 & \\
 & / \quad \backslash & \\
 A_2 & & D_2 \\
 & \backslash \quad / & \\
 & B_1 & \\
 & | & \\
 & D_1 & 
 \end{array}
 . \tag{7.131}$$

Cartan’s notation is used on the right hand side. The portion of the branching which ends at  $so(3) \sim B_1$

$$\begin{array}{ccc}
 & A_3 & \\
 & / \quad \backslash & \\
 A_2 & & D_2 \\
 & \backslash \quad / & \\
 & B_1 & 
 \end{array}
 , \tag{7.132}$$

for the representation  $[N] = [4]$  is displayed in Fig. 7.1, route I (left) and route II (right). The algebra  $u(4)$  constructed with  $s$  and  $p$  bosons is known as the



**Fig. 7.1.** Branchings of the representation  $[4]$  of  $u(4)$ . (a) Branching  $u(4) \supset u(3) \supset so(3)$  with labels  $n_p$  and  $L$ . (b) Branching  $u(4) \supset so(4) \supset so(3)$  with labels  $\omega$  and  $L$ . The total number of states on the left is equal to the total number on the right

vibron algebra. [F. Iachello and R.D. Levine, *Algebraic Theory of Molecules*, Oxford University Press, Oxford, England, 1995]. An account of the vibron algebra is also given in [A. Frank and P. van Isacker, *Algebraic Methods in Molecular and Nuclear Structure Physics*, J. Wiley and Sons, New York, 1994].

## 7.7 The Unitary Algebra $u(6)$

This algebra can be constructed by means of six boson operators,  $b_\alpha, \alpha = 1, \dots, 6$ . The 36 elements of the algebra are

$$G_{\alpha\beta} \doteq b_\alpha^\dagger b_\beta \quad \alpha, \beta = 1, \dots, 6. \quad (7.133)$$

### 7.7.1 Subalgebra Chains not Containing $so(3)$

#### The Canonical Chain

The canonical chain  $u(6) \supset u(5) \supset u(4) \supset u(3) \supset u(2) \supset u(1)$  is trivial.

#### The Vector Chain

The algebra  $u(6)$  can be constructed with two vector boson operators,  $p_{1\mu}^\dagger$  and  $p_{2\mu}^\dagger$ , leading to the chain  $u(6) \supset u_1(3) \oplus u_2(3)$ . The algebras  $u_1(3)$  and  $u_2(3)$  can then be decomposed as in Sect. 4.

### 7.7.2 Subalgebra Chains Containing $so(3)$

Of particular interest for applications are the chains that can be constructed with a scalar and a quadrupole boson operator. These chains contain the angular momentum algebra  $so(3)$ . The boson operators  $s^\dagger$  and  $d_\mu^\dagger$  transform as the representation  $l = 0$  and  $l = 2$  of  $so(3)$

$$\begin{aligned} l = 0 : & \quad s^\dagger \\ l = 2 : & \quad d_\mu^\dagger (\mu = 0, \pm 1, \pm 2). \end{aligned} \quad (7.134)$$

The elements of the algebra  $u(6)$  in Racah form are

$$\begin{aligned} G_0^{(0)}(ss) &= (s^\dagger \times \tilde{s})_0^{(0)} & 1 \\ G_0^{(0)}(dd) &= (d^\dagger \times \tilde{d})_0^{(0)} & 1 \\ G_\kappa^{(1)}(dd) &= (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\ G_\kappa^{(2)}(dd) &= (d^\dagger \times \tilde{d})_\kappa^{(2)} & 5 \\ G_\kappa^{(3)}(dd) &= (d^\dagger \times \tilde{d})_\kappa^{(3)} & 7 \\ G_\kappa^{(4)}(dd) &= (d^\dagger \times \tilde{d})_\kappa^{(4)} & 9 \\ G_\kappa^{(2)}(ds) &= (d^\dagger \times \tilde{s})_\kappa^{(2)} & 5 \\ G_\kappa^{(2)}(sd) &= (s^\dagger \times \tilde{d})_\kappa^{(2)} & 5 \end{aligned} \quad (7.135)$$

for a total of 36 operators. The standard procedure is then applied to this algebra.

### Subalgebras

The algebra of  $u(6)$  has three subalgebra chains that contain the angular momentum algebra  $so(3)$ .

*Subalgebra I:*  $u(6) \supset u(5) \supset so(5) \supset so(3) \supset so(2)$

The elements of the subalgebras are

$$\begin{aligned}
 u(5) : \\
 & (d^\dagger \times \tilde{d})_0^{(0)} & 1 \\
 & (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\
 & (d^\dagger \times \tilde{d})_\kappa^{(2)} & 5 \\
 & (d^\dagger \times \tilde{d})_\kappa^{(3)} & 7 \\
 & (d^\dagger \times \tilde{d})_\kappa^{(4)} & 9
 \end{aligned} \tag{7.136}$$

$$\begin{aligned}
 so(5) : \\
 & (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\
 & (d^\dagger \times \tilde{d})_\kappa^{(3)} & 7 \\
 so(3) : & & . \\
 & (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\
 so(2) : & & \\
 & (d^\dagger \times \tilde{d})_0^{(1)} & 1
 \end{aligned} \tag{7.137}$$

*Subalgebra II:*  $u(6) \supset su(3) \supset so(3) \supset so(2)$

The elements of the subalgebras are

$$\begin{aligned}
 u(3) : \\
 & (s^\dagger \times \tilde{s})_0^{(0)} + \sqrt{5}(d^\dagger \times \tilde{d})_0^{(0)} & 1 \\
 & (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\
 & (d^\dagger \times \tilde{s} + s^\dagger \times \tilde{d})_\kappa^{(2)} + \frac{\sqrt{7}}{2}(d^\dagger \times \tilde{d})_\kappa^{(2)} & 5 \\
 su(3) : \\
 & (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\
 & (d^\dagger \times \tilde{s} + s^\dagger \times \tilde{d})_\kappa^{(2)} + \frac{\sqrt{7}}{2}(d^\dagger \times \tilde{d})_\kappa^{(2)} & 5 \\
 so(3) : \\
 & (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\
 so(2) : \\
 & (d^\dagger \times \tilde{d})_0^{(1)} & 1
 \end{aligned} \tag{7.138}$$

This chain is doubled by the inner automorphism of  $u(3)$ , with  $\overline{u(3)}$  given by



$$\begin{aligned}
 \overline{u(3)} : \\
 \begin{array}{rcl}
 (s^\dagger \times \tilde{s})_0^{(0)} + \sqrt{5} (d^\dagger \times \tilde{d})_0^{(0)} & 1 \\
 (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\
 (d^\dagger \times \tilde{s} + s^\dagger \times \tilde{d})_\kappa^{(2)} - \frac{\sqrt{7}}{2} (d^\dagger \times \tilde{d})_\kappa^{(2)} & 5
 \end{array} \cdot \quad (7.139)
 \end{aligned}$$

In applications to nuclear physics, the algebras  $u(3)$  and  $\overline{u(3)}$  are usually called the prolate and oblate algebras.

*Subalgebra III:*  $u(6) \supset so(6) \supset so(5) \supset so(3) \supset so(2)$

The elements of the subalgebras are

$$\begin{aligned}
 so(6) : \\
 \begin{array}{rcl}
 (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\
 (d^\dagger \times \tilde{d})_\kappa^{(3)} & 7 \\
 (d^\dagger \times \tilde{s} + s^\dagger \times \tilde{d})_\kappa^{(2)} & 5
 \end{array} \\
 so(5) : \\
 \begin{array}{rcl}
 (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\
 (d^\dagger \times \tilde{d})_\kappa^{(3)} & 7
 \end{array} \quad (7.140) \\
 so(3) : \\
 \begin{array}{rcl}
 (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\
 so(2) : \\
 (d^\dagger \times \tilde{d})_0^{(1)} & 1
 \end{array}
 \end{aligned}$$

Again this chain is doubled by the inner automorphism of  $so(6)$ , with  $\overline{so(6)}$  given by

$$\begin{aligned}
 \overline{so(6)} : \\
 \begin{array}{rcl}
 (d^\dagger \times \tilde{d})_\kappa^{(1)} & 3 \\
 (d^\dagger \times \tilde{d})_\kappa^{(3)} & 7 \\
 i (d^\dagger \times \tilde{s} - s^\dagger \times \tilde{d})_\kappa^{(2)} & 5
 \end{array} \cdot \quad (7.141)
 \end{aligned}$$

## Invariant Operators

Invariant Casimir operators for all algebras included in the chains of the previous subsection can be constructed explicitly.

(a) Linear operators

(i)  $u(6)$

$$\begin{aligned}
 C_1(u(6)) &= G_0^{(0)}(ss) + \sqrt{5}G_0^{(0)}(dd) = (s^\dagger \times s)_0^{(0)} + \sqrt{5}(d^\dagger \times \tilde{d})_0^{(0)} \\
 &= \hat{n}_s + \hat{n}_d \equiv \hat{N} \quad (7.142)
 \end{aligned}$$

(ii)  $u(5)$

$$C_1(u(5)) = \sqrt{5}(d^\dagger \times \tilde{d})_0^{(0)} = \hat{n}_d \quad (7.143)$$

(iii)  $u(3)$

$$C_1(u(3)) = \hat{N} . \quad (7.144)$$

(b) Quadratic operators

For the unitary algebras  $u(6)$ ,  $u(5)$  and  $u(3)$  the quadratic Casimir operators can be taken to be

$$\begin{aligned} C_2(u(6)) &= \hat{N}(\hat{N} + 5) \\ C_2(u(5)) &= \hat{n}_d(\hat{n}_d + 4) \\ C_2(u(3)) &= \hat{N}(\hat{N} + 2) . \end{aligned} \quad (7.145)$$

For the orthogonal algebras appearing in the branchings of  $u(6)$  described in this section, they are

(i)  $so(3)$

$$C_2(so(3)) = G^{(1)} \cdot G^{(1)} \quad (7.146)$$

with  $G_\kappa^{(1)} \equiv G_\kappa^{(1)}(dd)$ . Introducing the angular momentum operator  $\hat{L}_\kappa = \sqrt{10}G_\kappa^{(1)}$ , this can be rewritten as  $C_2(so(3)) = \hat{L} \cdot \hat{L}$ . (The dot denotes scalar products.)

(ii)  $so(5)$

$$C_2(so(5)) = G^{(1)} \cdot G^{(1)} + G^{(3)} \cdot G^{(3)} \quad (7.147)$$

with  $G_\kappa^{(1)}$  as before and  $G_\kappa^{(3)} \equiv G_\kappa^{(3)}(dd)$ .

(iii)  $su(3)$

Introducing the operator

$$\tilde{G}_\kappa^{(2)} = G_\kappa^{(2)}(ds) + G_\kappa^{(2)}(sd) \mp \frac{\sqrt{7}}{2}G_\kappa^{(2)}(dd) , \quad (7.148)$$

the quadratic Casimir operator of  $su(3)$  can be written as

$$C_2(su(3)) = G^{(1)} \cdot G^{(1)} + \frac{4}{15} \tilde{G}^{(2)} \cdot \tilde{G}^{(2)} . \quad (7.149)$$

If, in addition to the angular momentum operator  $\hat{L}$ , one introduces the quadrupole operator  $\hat{Q} = \sqrt{8}\tilde{G}^{(2)}$ , the Casimir operator of  $su(3)$  can be rewritten as

$$C_2(su(3)) = \frac{1}{30} \left[ 3\hat{L} \cdot \hat{L} + \hat{Q} \cdot \hat{Q} \right] . \quad (7.150)$$

### Branchings

#### Branching I

The branching of representations of  $u(6)$  into representations of its subalgebra present a challenge not encountered when dealing with  $u(n)$ ,  $n \leq 4$ . One needs a hidden quantum number to characterize uniquely the basis. Finding this quantum number is one of the most difficult problems in algebraic theory. The branching  $u(6) \supset u(5) \supset so(5) \supset so(3) \supset so(2)$  is

$$\begin{aligned}
 u(6) \ [N] &\equiv \overbrace{\square \square \dots \square}^{N\text{-times}} \equiv [N, 0, 0, 0, 0, 0] \\
 u(5) \ [n_d] &\equiv \overbrace{\square \square \dots \square}^{n_d\text{-times}} \equiv [n_d, 0, 0, 0, 0] \quad n_d = N, N-1, \dots, 0 \\
 so(5) \ (v) &\equiv (v, 0) \quad v = n_d, n_d - 2, \dots, 1 \text{ or } 0 \\
 &\quad (n_d = \text{odd or even}) \\
 so(3) \ L &\quad \text{Algorithm 1} \\
 so(2) \ M_L &\quad -L \leq M_L \leq +L
 \end{aligned} \tag{7.151}$$

The step from  $so(5)$  to  $so(3)$  is non-canonical and thus requires the development of an algorithm to find the values of  $L$  contained in each representation  $n_d$ .

#### Algorithm 1

Partition  $n_d$  as

$$n_d = 2n_\beta + 3n_\Delta + \lambda \tag{7.152}$$

where

$$n_\beta = (n_d - v)/2; \quad n_\beta = 0, 1, \dots, \frac{n_d}{2} \text{ or } \frac{n_d - 1}{2}. \tag{7.153}$$

Then

$$L = \lambda, \lambda + 1, \lambda + 2, \dots, 2\lambda - 2, 2\lambda. \tag{7.154}$$

[Note that  $2\lambda - 1$  is missing!]. The additional quantum number  $n_\Delta = 0, 1, \dots$  is a missing label (hidden quantum number) needed to characterize uniquely the decompositions of representations of  $so(5)$  into representations of  $so(3)$ . This gives Table 7.1.

The complete classification for branching I is

$$\left. \begin{array}{ccccc}
 u(6) \supset u(5) \supset so(5) \supset so(3) \supset so(2) \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 N \quad n_d \quad v, n_\Delta \quad L \quad M_L
 \end{array} \right\}. \tag{7.155}$$

The total number of labels, including the missing label  $n_\Delta$ , is six.

**Table 7.1.** Decomposition of totally symmetric representations of  $u(5)$  into representations of  $so(3)$

$u(5)$ $n_d$	$\supset$ $so(5)$ $v$	$\supset$ $n_\Delta$	$so(3)$ $L$
0	0	0	0
1	1	0	2
2	2	0	4, 2
	0	0	0
3	3	0	6, 4, 3
		1	0
	1	0	0

*Branching II*

The branching  $u(6) \supset su(3) \supset so(3) \supset so(2)$  is

$$\begin{aligned}
 u(6) \quad [N] &\equiv \overbrace{\square \square \dots \square}^{N\text{-times}} \equiv [N, 0, 0, 0, 0, 0] \\
 su(3) \quad (\lambda, \mu) & \hspace{15em} \text{Algorithm 2} \hspace{5em} (7.156) \\
 so(3) \quad L & \hspace{15em} \text{Algorithm 3} \\
 so(2) \quad M_L & \hspace{15em} -L \leq M_L \leq +L
 \end{aligned}$$

Here, the so-called Elliott quantum numbers,  $\lambda = f_1 - f_2, \mu = f_2$ , are used instead of the entries in the Young tableau  $(f_1, f_2)$  in order to conform with commonly used notation. Both steps from  $u(6)$  to  $su(3)$  and from  $su(3)$  to  $so(3)$  are non-canonical. The step from  $su(3)$  to  $so(3)$  is not fully reducible.

**Algorithm 2**

The algorithm to find the values of  $(\lambda, \mu)$  contained in  $[N]$  is

$$\begin{aligned}
 (2N, 0) \oplus (2N - 4, 2) \oplus \dots & \oplus \left\{ \begin{array}{l} (0, N) \quad N = \text{even} \\ (2, N - 1) \quad N = \text{odd} \end{array} \right\} \oplus \\
 \oplus (2N - 6) \oplus (2N - 10, 2) \oplus \dots & \oplus \left\{ \begin{array}{l} (0, N - 3) \quad N - 3 = \text{even} \\ (2, N - 4) \quad N - 3 = \text{odd} \end{array} \right\} \oplus \\
 \oplus (2N - 12, 0) \oplus (2N - 16, 2) \oplus \dots \oplus & \left\{ \begin{array}{l} (0, N - 6) \quad N - 6 = \text{even} \\ (2, N - 7) \quad N - 6 = \text{odd} \end{array} \right\} \oplus \\
 \oplus \dots & \hspace{15em} (7.157)
 \end{aligned}$$

**Algorithm 3**

The algorithm to find the values of  $L$  contained in  $(\lambda, \mu)$  is

$$L = K, K + 1, K + 2, \dots, (K + \max \{\lambda, \mu\}) \hspace{5em} (7.158)$$

where

$$K = \text{integer} = \min\{\lambda, \mu\}, \min\{\lambda, \mu\} - 2, \dots, 1 \text{ or } 0 \tag{7.159}$$

with exception of  $K = 0$  for which

$$L = \max\{\lambda, \mu\}, \max\{\lambda, \mu\} - 2, \dots, 1 \text{ or } 0 . \tag{7.160}$$

Here  $K$  is the missing label. This gives Table 7.2.

**Table 7.2.** Decomposition of totally symmetric representations of  $u(6) \supset su(3)$  into representations of  $so(3)$

$u(6)$ $N$	$\supset$ $su(3)$ $(\lambda, \mu)$	$\supset$ $K$	$so(3)$ $L$
0	(0, 0)	0	0
1	(2, 0)	0	2, 0
2	(4, 0)	0	4, 2, 0
	(0, 2)	0	2, 0
3	(6, 0)	0	6, 4, 2, 0
	(2, 2)	0	2, 0
		2	2, 3, 4
	(0, 0)	0	0

The complete classification for branching II is

$$\left| \begin{array}{cccc} u(6) \supset & su(3) \supset & so(3) \supset & so(2) \\ \downarrow & \downarrow & \downarrow & \downarrow \\ N & (\lambda, \mu) K & L & M_L \end{array} \right\} . \tag{7.161}$$

Again, a total of six labels is needed.

*Branching III*

The branching  $u(6) \supset so(6) \supset so(5) \supset so(3) \supset so(2)$  is

$$\begin{array}{ll}
 u(6) [N] \equiv \overbrace{\square \square \dots \square}^{N\text{-times}} \equiv [N, 0, 0, 0, 0, 0] & \\
 so(6) \sigma \equiv (\sigma, 0, 0) & \sigma = N, N - 2, \dots, 1 \text{ or } 0 \\
 & (N = \text{odd or even}) \\
 so(5) \tau \equiv (\tau, 0) & \tau = \sigma, \sigma - 1, \dots, 0 \quad . \tag{7.162} \\
 so(3) L & \text{Algorithm 4} \\
 so(2) M_L & -L \leq M_L \leq +L
 \end{array}$$

**Algorithm 4**

The algorithm to find the values of  $L$  contained in each representation  $\tau$  is: Partition  $\tau$  as

$$\tau = 3\nu_\Delta + \lambda, \quad \nu_\Delta = 0, 1, \dots \tag{7.163}$$

and take

$$L = 2\lambda, 2\lambda - 2, \dots, \lambda + 1, \lambda. \tag{7.164}$$

[Note again that  $2\lambda - 1$  is missing!]. The missing label is here  $\nu_\Delta$ .

This gives Table 7.3.

**Table 7.3.** Decomposition of totally symmetric representations of  $u(6) \supset so(6)$  into representations of  $so(3)$

$u(6) \supset so(6) \supset so(5) \supset so(3)$				
$N$	$\sigma$	$\tau$	$\nu_\Delta$	$L$
0	0	0	0	0
1	1	1	0	2
		0	0	0
2	2	2	0	4, 2
		1	0	2
		0	0	0
	0	0	0	0
3	3	3	0	6, 4, 3
			1	0
		2	0	4, 2
		1	0	2
		0	0	0
	1	1	0	2
		0	0	0

The complete classification for branching III is

$$\left| \begin{array}{ccccc} u(6) \supset so(6) \supset so(5) \supset so(3) \supset so(2) \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ N \quad \sigma \quad \tau, \nu_\Delta \quad L \quad M_L \end{array} \right\rangle \tag{7.165}$$

with six quantum numbers as before.

In all three chains, the total number of labels needed to characterize uniquely the totally symmetric representations of  $u(6)$  is six, as one can see by considering the Gel'fand pattern

$$\begin{array}{cccccc} n_1 & 0 & 0 & 0 & 0 & 0 \\ & n_2 & 0 & 0 & 0 & 0 \\ & & n_3 & 0 & 0 & 0 \\ & & & n_4 & 0 & 0 \\ & & & & n_5 & 0 \\ & & & & & n_6 \end{array} \cdot \tag{7.166}$$



$$G_{\alpha\beta} \doteq b_{\alpha}^{\dagger} b_{\beta} \quad (\alpha, \beta = 1, 2, \dots, 7) . \tag{7.169}$$

The algebra  $u(7)$  is interesting for two reasons: (i) It is the unitary algebra of lowest rank that contains as a subalgebra one of the exceptional algebras,  $g_2$ , and (ii) it has applications to the three-body problem in quantum mechanics. The subalgebra chains that describe these two situations are discussed in the following subsections.

### 7.8.1 Subalgebra Chain Containing $g_2$

This chain can be constructed by introducing an octupole boson  $f^{\dagger}$  that transforms as the representation  $l = 3$  of  $so(3)$

$$l = 3: \quad f_{\mu}^{\dagger} \quad (\mu = 0, \pm 1, \pm 2, \pm 3) . \tag{7.170}$$

The elements of the algebra  $u(7)$  in Racah form are

$$\begin{aligned} G_0^{(0)}(ff) &= \left( f^{\dagger} \times \tilde{f} \right)_0^{(0)} & 1 \\ G_{\kappa}^{(1)}(ff) &= \left( f^{\dagger} \times \tilde{f} \right)_0^{(1)} & 3 \\ G_{\kappa}^{(2)}(ff) &= \left( f^{\dagger} \times \tilde{f} \right)_{\kappa}^{(2)} & 5 \\ G_{\kappa}^{(3)}(ff) &= \left( f^{\dagger} \times \tilde{f} \right)_{\kappa}^{(3)} & 7 , \\ G_{\kappa}^{(4)}(ff) &= \left( f^{\dagger} \times \tilde{f} \right)_{\kappa}^{(4)} & 9 \\ G_{\kappa}^{(5)}(ff) &= \left( f^{\dagger} \times \tilde{f} \right)_{\kappa}^{(5)} & 11 \\ G_{\kappa}^{(6)}(ff) &= \left( f^{\dagger} \times \tilde{f} \right)_{\kappa}^{(6)} & 13 \end{aligned} \tag{7.171}$$

for a total of 49 elements.

*Subalgebra I:*  $u(7) \supset so(7) \supset g_2 \supset so(3) \supset so(2)$

The elements are:

$$\begin{aligned} so(7) : & \\ & \left( f^{\dagger} \times \tilde{f} \right)_{\kappa}^{(k)} \quad k = 1, 3, 5 \quad 21 \\ g_2; & \\ & \left( f^{\dagger} \times \tilde{f} \right)_{\kappa}^{(k)} \quad k = 1, 5 \quad 14 \\ so(3) : & \\ & \left( f^{\dagger} \times \tilde{f} \right)_{\kappa}^{(1)} \quad 3 \\ so(2) : & \\ & \left( f^{\dagger} \times \tilde{f} \right)_0^{(1)} \quad 1 \end{aligned} \tag{7.172}$$



**Table 7.4.** Decomposition of totally symmetric representations of  $u(7) \supset so(7)$  into representations of  $so(3)$

$u(7) \supset so(7) \supset g_2 \supset so(3)$			
$N$	$\omega$	$\gamma$	$L$
0	0	0	0
1	1	1	3
2	2	2	6, 4, 2
	0	0	0
3	3	3	9, 7, 6, 5, 4, 3, 1
	1	1	3
4	4	4	12, 10, 9, 8 <sup>2</sup> , 7, 6 <sup>2</sup> , 5 <sup>2</sup> , 4, 3 <sup>2</sup> , 2, 1
	2	2	6, 4, 2
	0	0	0

**Branching**

The branching of representations  $[N]$  of  $u(7)$  is

$$\begin{aligned}
 u(7) \quad [N] &\equiv [N, 0, 0, 0, 0, 0, 0] \\
 so(7) \quad \omega &\equiv (\omega, 0, 0) \quad \omega = N, N - 2, \dots, 1 \text{ or } 0 \quad (N = \text{odd or even})
 \end{aligned}$$

$$\begin{aligned}
 g_2 \quad \gamma &\equiv (\gamma, 0) \text{ Racah algorithm} \\
 so(3) \quad L &\text{ Racah algorithm .} \\
 so(2) \quad M_L &\quad -L \leq M_L \leq +L
 \end{aligned} \tag{7.173}$$

The steps  $so(7) \supset g_2$  and  $g_2 \supset so(3)$  are non-canonical and involve the exceptional group  $g_2$ . The Racah algorithm gives Table 7.4. The algorithm can be found in [G. Racah, *Theory of Complex Spectra-IV*, Phys. Rev. 76, 1352 (1949).] Note that for bosonic representations the algebra  $g_2$  does not provide any new label and the multiplicities must be resolved by additional missing labels. (The algebra  $g_2$  will be also discussed in Chap. 8 for fermionic systems for which it provides new labels.)

**7.8.2 The Triplet Chains**

These chains can be constructed with two three-dimensional vector bosons,  $b_{\rho,m}^\dagger$  ( $m = 0 \pm 1$ ),  $b_{\lambda,m}^\dagger$  ( $m = 0, \pm 1$ ) and a scalar boson  $s^\dagger$ , generically denoted  $c_\alpha^\dagger$  ( $\alpha = 1, \dots, 7$ )

$$\begin{aligned}
 l = 1 : & \quad b_{\rho,m}^\dagger \quad (m = 0, \pm 1) \\
 l = 1 : & \quad b_{\lambda,m}^\dagger \quad (m = 0, \pm 1) \\
 l = 0 : & \quad s^\dagger
 \end{aligned} \tag{7.174}$$

together with the corresponding annihilation operators  $c_\alpha$  ( $\alpha = 1, \dots, 7$ ). These chains are important in the study of the three-body problem in quantum mechanics. The subscript  $\rho, \lambda$  that distinguishes the two vector bosons is used to indicate that these vector bosons represent the second quantized form of the Jacobi variables  $\boldsymbol{\rho}, \boldsymbol{\lambda}$  that characterize the geometric configuration of a three-body system. The Jacobi variables are defined in terms of the coordinates of the three particles  $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3$  as  $\boldsymbol{\rho} = \frac{1}{\sqrt{2}}(\mathbf{r}_1 - \mathbf{r}_2)$  and  $\boldsymbol{\lambda} = \frac{1}{\sqrt{6}}(\mathbf{r}_1 + \mathbf{r}_2 - 2\mathbf{r}_3)$ . The bilinear products

$$G_{\alpha\alpha'} \doteq c_\alpha^\dagger c_{\alpha'} \quad (\alpha, \alpha' = 1, \dots, 7) \quad (7.175)$$

generate  $u(7)$ . The basis states are written as

$$\mathcal{B} : \frac{1}{\mathcal{N}} (b_\rho^\dagger)^{n_\rho} (b_\lambda^\dagger)^{n_\lambda} (s^\dagger)^{N-n_\rho-n_\lambda} |0\rangle, \quad (7.176)$$

where  $\mathcal{N}$  is a normalization constant. The Racah form of  $u(7)$  is obtained by introducing the operators

$$\begin{aligned} \tilde{b}_{\rho,m} &= (-1)^{1-m} b_{\rho,-m} \\ \tilde{b}_{\lambda,m} &= (-1)^{1-m} b_{\lambda,-m} \\ \tilde{s} &= s. \end{aligned} \quad (7.177)$$

It can be written as

$$\begin{aligned} \hat{n}_s &= (s^\dagger \times \tilde{s}) & 1 \\ \hat{D}_{\rho,\mu} &= (b_\rho^\dagger \times \tilde{s} - s^\dagger \times \tilde{b}_\rho)_{\mu}^{(1)} & 3 \\ \hat{D}_{\lambda,\mu} &= (b_\lambda^\dagger \times \tilde{s} - s^\dagger \times \tilde{b}_\lambda)_{\mu}^{(1)} & 3 \\ \hat{A}_{\rho,\mu} &= i (b_\rho^\dagger \times \tilde{s} + s^\dagger \times \tilde{b}_\rho)_{\mu}^{(1)} & 3 \\ \hat{A}_{\lambda,\mu} &= i (b_\lambda^\dagger \times \tilde{s} + s^\dagger \times \tilde{b}_\lambda)_{\mu}^{(1)} & 3 \\ \hat{G}_{S,\mu}^{(\ell)} &= (b_\rho^\dagger \times \tilde{b}_\rho + b_\lambda^\dagger \times \tilde{b}_\lambda)_{\mu}^{(\ell)} & 9 \\ \hat{G}_{A,\mu}^{(\ell)} &= i (b_\rho^\dagger \times \tilde{b}_\lambda - b_\lambda^\dagger \times \tilde{b}_\rho)_{\mu}^{(\ell)} & 9 \\ \hat{G}_{M\rho,\mu}^{(\ell)} &= (b_\rho^\dagger \times \tilde{b}_\lambda + b_\lambda^\dagger \times \tilde{b}_\rho)_{\mu}^{(\ell)} & 9 \\ \hat{G}_{M\lambda,\mu}^{(\ell)} &= (b_\rho^\dagger \times \tilde{b}_\rho - b_\lambda^\dagger \times \tilde{b}_\lambda)_{\mu}^{(\ell)} & 9 \end{aligned} \quad (7.178)$$

with  $\ell = 0, 1, 2$ . This form is not the usual Racah form of the algebra in which the bilinear products are only angular momentum coupled. In applications to the many-body problem, especially in quantum chemistry, it is useful to assign to the creation and annihilation operators properties under

transformations of a discrete group. The most important property is parity, where the transformation is inversion of the coordinates  $\mathbf{r}_i \rightarrow -\mathbf{r}_i$ . The two boson operators  $b_\rho^\dagger$  and  $b_\lambda^\dagger$  are assumed to be odd under parity, while the operator  $s^\dagger$  is assumed to be even. (In general, in this chapter, the parity  $\mathcal{P}$  of the boson operators  $b_{l,m}^\dagger$  is assumed to be  $(-1)^l$ .) For applications to the three-body problem, it is convenient to assign to the elements of the algebra definite transformation properties under permutation of the three-bodies. The boson operators  $s^\dagger, b_{\rho,m}^\dagger, b_{\lambda,m}^\dagger$  are assumed to transform under the transposition  $P(12)$  as

$$P(12) \begin{pmatrix} s^\dagger \\ b_{\rho,m}^\dagger \\ b_{\lambda,m}^\dagger \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} s^\dagger \\ b_{\rho,m}^\dagger \\ b_{\lambda,m}^\dagger \end{pmatrix}, \quad (7.179)$$

and under the cyclic permutation  $P(123)$  as

$$P(123) \begin{pmatrix} s^\dagger \\ b_{\rho,m}^\dagger \\ b_{\lambda,m}^\dagger \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(2\pi/3) & \sin(2\pi/3) \\ 0 & -\sin(2\pi/3) & \cos(2\pi/3) \end{pmatrix} \begin{pmatrix} s^\dagger \\ b_{\rho,m}^\dagger \\ b_{\lambda,m}^\dagger \end{pmatrix}. \quad (7.180)$$

The permutation group  $S_3$  is a six element discrete group which is isomorphic to the dihedral group  $D_3$ . In order to characterize the transformation properties of the elements of the algebra (and of the states) one can use the label of either group, given by

$S_3$	$D_3$	
$S \equiv \square \square \square$	$A_1$	
$M \equiv \begin{matrix} \square & \square \\ & \square \end{matrix}$	$E$	·
$A \equiv \begin{matrix} \square \\ \square \end{matrix}$	$A_2$	

(7.181)

This notation is used in (7.178). The construction of bosonic algebras with definite transformation properties under discrete groups is discussed in [R. Bijker and A. Leviatan, *Algebraic Treatment of Collective Excitations in Baryon Spectroscopy in Symmetries in Science VII: Spectrum Generating Algebras and Dynamic Symmetries in Physics*, Edited by B. Gruber and T. Otsuka, Plenum Press, New York, 1994].

### Subalgebras II

Since the number of elements of  $u(7)$  is relatively large, there are several possible subalgebra chains, four of which are included here.

*Subalgebra II.1*

This is the chain

$$u(7) \supset u(6) \supset so(6) \supset so_\rho(3) \oplus so_\lambda(3) \supset so(3) \supset so(2) . \quad (7.182)$$

States in this chain are characterized by the quantum numbers

$$\left| \begin{array}{ccccccc} u(7) \supset u(6) \supset so(6) \supset so_\rho(3) \oplus so_\lambda(3) \supset so(3) \supset so(2) \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ N \quad n \quad \gamma \quad L_\rho \quad L_\lambda \quad L \quad M_L \end{array} \right\rangle \quad (7.183)$$

with branching

$$\begin{array}{llll} u(7) & N \equiv [N, 0, 0, 0, 0, 0] & & \\ u(6) & n \equiv [n, 0, 0, 0, 0, 0] & n = N, N - 1, \dots, 0 & \\ so(6) & \gamma \equiv (\gamma, 0, 0) & \gamma = n, n - 2, \dots, 1 \text{ or } 0 \\ & & (n = \text{odd or even}) & \\ so_\rho(3) \oplus so_\lambda(3) & L_\rho, L_\lambda & \text{Algorithm 5} & \\ so(3) & L & |L_\rho - L_\lambda| \leq L \leq |L_\rho + L_\lambda| & \\ so(2) & M_L & -L \leq M_L \leq +L & \end{array} . \quad (7.184)$$

**Algorithm 5**

The values of  $L_\rho$  and  $L_\lambda$  are obtained by partitioning  $\gamma$  as

$$\gamma = 2\nu + L_\rho + L_\lambda \quad \text{with } \nu = 0, 1, \dots \quad (7.185)$$

*Subalgebra II.2*

This is the chain

$$u(7) \supset so(7) \supset so(6) \supset so_\rho(3) \oplus so_\lambda(3) \supset so(3) \supset so(2) . \quad (7.186)$$

States in this chain are characterized by

$$\left| \begin{array}{ccccccc} u(7) \supset so(7) \supset so(6) \supset so_\rho(3) \oplus so_\lambda(3) \supset so(3) \supset so(2) \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ N \quad \omega \quad \gamma \quad L_\rho \quad L_\lambda \quad L \quad M_L \end{array} \right\rangle , \quad (7.187)$$

with branching

$$\begin{array}{llll} u(7) & N \equiv [N, 0, 0, 0, 0, 0] & & \\ so(7) & \omega \equiv (\omega, 0, 0) & \omega = N, N - 2, \dots, 1 \text{ or } 0 & \\ so(6) & \gamma \equiv (\gamma, 0, 0) & \gamma = \omega, \omega - 1, \dots, 0 & \\ so_\rho(3) \oplus so_\lambda(3) & L_\rho, L_\lambda & \text{Algorithm 1} & \\ so(3) & L & |L_\rho - L_\lambda| \leq L \leq |L_\rho + L_\lambda| & \\ so(2) & M_L & -L \leq M_L \leq +L & \end{array} . \quad (7.188)$$

### Subalgebras III

Two other subalgebra chains have been used extensively in hadronic physics (three-quark system). [R. Bijker, F. Iachello and A. Leviatan, *Algebraic Models of Hadronic Structure. I. Nonstrange Baryons*, Ann. Phys. (NY) 236, 69 (1994).]

#### Subalgebra III.1

This is the chain

$$u(7) \supset u_\rho(3) \oplus u_\lambda(4) \supset u_\rho(3) \oplus u_\lambda(3) \supset so_\rho(3) \oplus so_\lambda(3) \supset so(3) \supset so(2) . \tag{7.189}$$

States in this chain are characterized by the quantum numbers

$$\left. \begin{array}{ccccccc} u(7) \supset u_\rho(3) \oplus u_\lambda(4) \supset u_\rho(3) \oplus u_\lambda(3) \supset so_\rho(3) \oplus so_\lambda(3) & & & & & & \\ \downarrow & & \downarrow & \downarrow & \downarrow & \downarrow & \\ N & & n_\rho & n_\lambda & L_\rho & L_\lambda & \\ & & & & & & \\ & & \supset so(3) \supset so(2) & & & & \\ & & \downarrow & \downarrow & & & \\ & & L & M_L & & & \end{array} \right\} , \tag{7.190}$$

with branching

$u(7)$	$N \equiv [N, 0, 0, 0, 0, 0, 0]$	
$u_\rho(3) \oplus u_\lambda(3)$	$n_\rho \equiv (n_\rho, 0, 0)$ $n_\lambda \equiv (n_\lambda, 0, 0)$	$n_\rho = 0, 1, \dots, N$ $n_\lambda = 0, 1, \dots, N - n_\rho$
$so_\rho(3) \oplus so_\lambda(3)$	$L_\rho$ $L_\lambda$	$L_\rho = n_\rho, n_\rho - 2, \dots, 1$ or $0$ . $L_\lambda = n_\lambda, n_\lambda - 2, \dots, 1$ or $0$
$so(3)$	$L$	$ L_\rho - L_\lambda  \leq L \leq  L_\rho + L_\lambda $
$so(2)$	$M_L$	$-L \leq M_L \leq +L$

(7.191)

#### Subalgebra III.2

This is the chain

$$u(7) \supset u_\rho(3) \oplus u_\lambda(4) \supset u_\rho(3) \oplus so_\lambda(4) \supset so_\rho(3) \oplus so_\lambda(3) \supset so(3) \supset so(2) . \tag{7.192}$$

States in this chain are characterized by the quantum numbers


$$\left. \begin{array}{ccccccc} u(7) \supset u_\rho(3) \oplus u_\lambda(4) \supset u_\rho(3) \oplus so_\lambda(4) \supset so_\rho(3) \oplus so_\lambda(3) & & & & & & \\ \downarrow & & \downarrow & \downarrow & \downarrow & \downarrow & \\ N & & n_\rho & \omega & L_\rho & L_\lambda & \\ & & & & & & \\ & & \supset so(3) \supset so(2) & & & & \\ & & \downarrow & \downarrow & & & \\ & & L & M_L & & & \end{array} \right\} , \tag{7.193}$$

with branching


$$\begin{array}{lll}
 u(7) & N \equiv [N, 0, 0, 0, 0, 0, 0] & \\
 u_\rho(3) \oplus so_\lambda(4) & n_\rho \equiv (n_\rho, 0, 0) & n_\rho = 0, 1, \dots, N \\
 & \omega \equiv (\omega, 0) & \omega = N - n_\rho, N - n_\rho - 2, \dots, 1 \text{ or } 0 \\
 so_\rho(3) \oplus so_\lambda(3) & L_\rho & L_\rho = n_\rho, n_\rho - 2, \dots, 1 \text{ or } 0 \\
 & L_\lambda & L_\lambda = 0, 1, \dots, \omega \\
 so(3) & L & |L_\rho - L_\lambda| \leq L \leq |L_\rho + L_\lambda| \\
 so(2) & M_L & -L \leq M_L \leq +L
 \end{array} \tag{7.194}$$

### Lattice of Algebras

The lattice of algebras for the chains discussed above is



and



Note that the two lattices merge at the level of  $so_\rho(3) \oplus so_\lambda(3)$ . [F. Iachello, *Algebraic Theory of the Three-body Problem*, in *Symmetries in Science VIII*, Edited by B. Gruber, Plenum Press, New York, 1995.]

# 8 Fermion Realizations

## 8.1 Fermion Operators

In Chap. 7 a realization of Lie algebras in terms of boson operators has been given. These operators have been used either in connection with coordinates and momenta, as in harmonic oscillator problems, see Chap. 7, Examples 1, 2 and 6, or as operators which create or annihilate particles with integer values of the angular momentum  $l = 0, 1, \dots$ . In the 1920's, it became evident that particles exist with half-integer values of the angular momentum,  $j = \frac{1}{2}, \frac{3}{2}, \dots$ . These particles are called fermions. For applications to physics, it is of interest to consider realizations of Lie algebras in terms of fermion operators. In view of the pervasive presence in physics of particles with half-integer spin, most notably electrons and nucleons, fermion realizations have become an important part of Lie algebraic methods. These will be discussed in this chapter.

We begin by introducing an operation called anticommutator, denoted by a curly bracket,  $\{, \}$ . The anticommutator of two quantities  $X, Y$  is

$$\{X, Y\} = XY + YX . \quad (8.1)$$

The anticommutator is sometimes denoted by  $[, ]_+$ . In this book, the curly bracket notation will be used. We then introduce fermion creation,  $a_i^\dagger$ , and annihilation,  $a_i$ , operators, satisfying anticommutation relations,

$$\{a_i, a_{i'}^\dagger\} = \delta_{ii'}; \quad \{a_i, a_{i'}\} = \{a_i^\dagger, a_{i'}^\dagger\} = 0 . \quad (8.2)$$

These operators are the key ingredient in the construction.

## 8.2 Lie Algebras Constructed with Fermion Operators

Lie algebras can be constructed from bilinear products of fermion creation and annihilation operators

$$g \doteq A_{ik} = a_i^\dagger a_k \quad (i, k = 1, \dots, n) , \quad (8.3)$$

as in the previous case of boson operators. The elements  $A_{ik}$  satisfy the commutation relations of the unitary algebra  $u(n)$  [and  $gl(n)$ ]

$$[A_{ik}, A_{st}] = A_{it} \delta_{ks} - A_{sk} \delta_{it} . \quad (8.4)$$

A basis can be constructed by acting with the creation operators on a vacuum state  $|0\rangle$ . This basis will be denoted by  $\mathcal{F}$ .

$$\mathcal{F} : \frac{1}{\mathcal{N}} a_i^\dagger a_{i'}^\dagger \dots |0\rangle . \quad (8.5)$$

In contrast with the case of boson operators which generate totally symmetric representations, the irreducible representations of  $u(n)$  generated by acting with fermion operators on  $|0\rangle$  are the totally antisymmetric representations, with Young tableau

$$\left. \begin{array}{c} \square \\ \square \\ \vdots \\ \square \end{array} \right\} N_F . \quad (8.6)$$

For  $u(n)$  the Young tableau has  $n$  entries. The totally antisymmetric representations have  $N_F$  entries 1 and  $n - N_F$  entries 0

$$\{N_F\} \equiv \underbrace{[1, 1, \dots, 1, 0, \dots, 0]}_N . \quad (8.7)$$

The short-hand notation  $\{N_F\}$  is often used to denote these representations. The zeros are usually not written, except for the identity representation  $[0, 0, \dots, 0] \equiv [0]$ . The basis  $\mathcal{F}$  is often called a Fermi-Dirac basis. Since any Lie algebra is a subalgebra of  $gl(n)$  it can be constructed with bilinear products of fermion operators.

### 8.3 Racah Form

While in the case of bosons both the uncoupled and coupled (Racah) form of the Lie algebra have been extensively used, in the case of fermions only the Racah form has been to a large extent used, and will be considered in this book. The Racah form of the Lie algebra can be obtained by introducing fermion operators that transform as representations  $|j, m\rangle$  of  $spin(3) \supset spin(2)$ , with  $j = \text{half-integer}$ . We use here  $spin(3)$  and  $spin(2)$  instead of  $so(3)$  and  $so(2)$ , since we need to consider explicitly spinor representations. The corresponding creation and annihilation operators will be denoted by



$$\begin{aligned} a_{j,m}^\dagger & \quad m = \pm\frac{1}{2}, \pm\frac{3}{2}, \dots, \pm j \\ a_{j,m} & \quad m = \pm\frac{1}{2}, \pm\frac{3}{2}, \dots, \pm j \end{aligned} \quad (8.8)$$

These operators satisfy anticommutation relations

$$\begin{aligned} \{a_{j,m}, a_{j',m'}^\dagger\} &= \delta_{jj'} \delta_{mm'} \\ \{a_{j,m}, a_{j',m'}\} &= \{a_{j,m}^\dagger, a_{j',m'}^\dagger\} = 0. \end{aligned} \quad (8.9)$$

In constructing the Lie algebra it is convenient to introduce the operators

$$\tilde{a}_{j,m} = (-1)^{j-m} a_{j,-m} \quad (8.10)$$

that transform as tensors under  $spin(3) \supset spin(2)$ . (An alternative definition is with a phase  $(-1)^{j+m}$ . This alternative definition introduces minus signs in some formulas, but it does not alter the algebraic structure. Operators with physical meaning constructed with creation and annihilation operators must be defined accordingly. This problem does not arise for Racah boson realizations, since  $l$  and  $m$  are integers.)

The Racah form of  $u(n)$  is

$$g \doteq A_\mu^{(\lambda)}(j, j') = [a_j^\dagger \times \tilde{a}_{j'}]_\mu^{(\lambda)}, \quad (8.11)$$

with  $|j + j'| \geq \lambda \geq |j - j'|$  and  $n = \sum_i (2j_i + 1)$ . The commutation relations are

$$\begin{aligned} [A_\mu^{(\lambda)}(j, j'), A_{\mu'}^{(\lambda')}(j'', j''')] &= - \sum_{\lambda'' \mu''} [(2\lambda + 1)(2\lambda' + 1)]^{1/2} \langle \lambda \mu \lambda' \mu' | \lambda'' \mu'' \rangle \\ &\quad \times \left[ (-)^{\lambda'' + j + j'''} \left\{ \begin{matrix} \lambda \lambda' \lambda'' \\ j''' j j' \end{matrix} \right\} \delta_{j' j''} A_{\mu''}^{(\lambda'')}(j, j''') \right. \\ &\quad \left. - (-)^{\lambda + \lambda' + j' + j''} \left\{ \begin{matrix} \lambda \lambda' \lambda'' \\ j'' j' j \end{matrix} \right\} \delta_{j j''} A_{\mu''}^{(\lambda'')}(j'', j') \right]. \end{aligned} \quad (8.12)$$

Note that, since  $j$  is half-integer, one can construct with fermion operators that transform as representations of  $spin(3) \supset spin(2)$  only unitary algebras in an even number of dimensions, a peculiarity of these realizations.

## 8.4 The Algebras $u(2j + 1)$

Consider a single value of  $j$ . There are  $(2j + 1)$  values of  $m$ . Thus  $n = 2j + 1$ . With this single value of  $j$ , we can construct the algebra  $u(2j + 1)$ . There are  $(2j + 1)^2$  elements in the algebra

$$A_\mu^{(\lambda)}(j, j) = [a_j^\dagger \times \tilde{a}_j]_\mu^{(\lambda)}. \quad (8.13)$$

To the algebra  $u(2j + 1)$  we can now apply the general procedure of constructing subalgebras.

### 8.4.1 Subalgebra Chain Containing $spin(3)$

A generic subalgebra chain (called a classification scheme) for  $u(2j + 1)$  constructed with fermion operators can be obtained as follows:

- (i) Exclude the element with  $\lambda = 0, \mu = 0$ ; this gives  $su(2j + 1)$ .
- (ii) Retain only terms with  $\lambda = \text{odd}$ ; this gives the Lie algebra  $sp(2j + 1, C) \equiv sp(2j + 1)$ . Note that this situation is different from that of Lie algebras constructed in terms of boson operators, since in that case, retaining terms with  $\lambda = \text{odd}$  generates the orthogonal Lie algebras,  $so(2l + 1)$ .
- (iii) Retain the term with  $\lambda = 1$ ; this gives the algebra  $spin(3)$ .
- (iv) Retain the term with  $\lambda = 1, \mu = 0$ ; this gives the algebra  $spin(2)$ . A generic subalgebra chain for single  $j$  is thus

$$u(2j + 1) \supset su(2j + 1) \supset sp(2j + 1) \supset \dots \supset spin(3) \supset spin(2). \quad (8.14)$$

Dots have been inserted between  $sp(2j + 1)$  and  $spin(3)$ , since, for large  $j$ , there may be intermediate steps. [R.H. Flowers, *Studies in  $j$ - $j$  Coupling. I. Classification of Nuclear and Atomic States*, Proc. Roy. Soc. A212, 248 (1952).]

### 8.4.2 The Algebras $u(2)$ and $su(2)$ . Spinors

Because of the pervasive presence in physics of particles with angular momentum  $\frac{1}{2}$ , it is of great interest to construct explicitly the Lie algebra  $u(2)$  in terms of fermion operators with  $j = \frac{1}{2}$ , called spinors. The fermion creation operators are written as

$$a_{\frac{1}{2}, +\frac{1}{2}}^\dagger, \quad a_{\frac{1}{2}, -\frac{1}{2}}^\dagger. \quad (8.15)$$

An alternative notation, often found in books in condensed matter physics, is  $a_\uparrow^\dagger, a_\downarrow^\dagger$ , called spin up, and spin down notation.

The 4 elements of  $u(2)$  are

$$\begin{aligned} A_\mu^{(1)}(\frac{1}{2}, \frac{1}{2}) &= \left[ a_{\frac{1}{2}}^\dagger \times \tilde{a}_{\frac{1}{2}} \right]_\mu^{(1)} \quad 3 \\ A_0^{(0)}(\frac{1}{2}, \frac{1}{2}) &= \left[ a_{\frac{1}{2}}^\dagger \times \tilde{a}_{\frac{1}{2}} \right]_0^{(0)} \quad 1 \end{aligned} \quad (8.16)$$

By deleting the element  $A_0^{(0)}$  one obtains the subalgebra  $su(2) \sim sp(2) \sim spin(3)$ . By considering only the element  $A_0^{(1)}$ , one obtains the subalgebra  $spin(2)$ . The basis is

$$\left| \begin{array}{ccc} u(2) \supset su(2) \supset spin(2) \\ \downarrow \qquad \qquad \downarrow \\ N_F \qquad \qquad M_J \end{array} \right\rangle. \quad (8.17)$$

Note that, for totally antisymmetric representations, no new quantum number is introduced when going from  $u(2)$  to  $su(2) \sim sp(2)$ . The elements of the Lie algebra  $u(2)$  constructed with spin  $\frac{1}{2}$  fermion operators have a straightforward physical meaning. They are the total spin operator,  $\hat{S}$ , and the number operator for fermions,  $\hat{N}_F$ ,

$$\begin{aligned}\hat{S}_\mu &= -\sqrt{\frac{1}{2}} \left[ a_{1/2}^\dagger \times \tilde{a}_{1/2} \right]_\mu^{(1)} \\ \hat{N}_F &= -\sqrt{2} \left[ a_{1/2}^\dagger \times \tilde{a}_{1/2} \right]_0^{(0)}\end{aligned}\tag{8.18}$$

(The minus sign arises from the choice of phases in (8.10).) The only states of  $u(2)$  that can be constructed with fermion operators are

$$\begin{aligned}\square &\equiv [1] \quad N_F = 1 \\ \square &\equiv [1, 1] \quad N_F = 2\end{aligned}\tag{8.19}$$

in addition to the vacuum  $|0\rangle$  that transforms as the identity representation  $[0]$ . This property is known to physicists as Pauli principle. The classification of antisymmetric states is rewritten, for comparison with the subsequent sections, as in Table 8.1. The algebra of  $u(2)$  constructed with fermion

**Table 8.1.** Classification of antisymmetric states of  $u(2)$

$j = 1/2$	$u(2)$	$su(2)$
	$[0]$	0
	$[1]$	1/2
	$[1, 1]$	0

operators has had many applications in physics, and hence many different notations have been used to label the states. Antisymmetric states of  $u(2)$  have been labeled either by  $\{N_F\}$  or by  $[\lambda_1, \lambda_2]$ . The number of fermions,  $N_F$ , is related to the labels  $[\lambda_1, \lambda_2]$  by

$$N_F = \lambda_1 + \lambda_2.\tag{8.20}$$

Also, in this case, in the general classification scheme of (8.14), all steps, except the last one, coincide, since  $su(2) \sim sp(2) \sim spin(3)$ . Finally, either  $u(2)$  or  $su(2)$  can be used to classify the states, and the use of both the labels  $[\lambda_1, \lambda_2]$  of  $u(2)$  and the label  $J$  of  $su(2)$  is redundant.

### 8.4.3 The Algebra $u(4)$

Another case of considerable interest is the case of  $j = \frac{3}{2}$ ,  $n = 4$ . The 16 elements of the algebra are

$$\begin{aligned}
 A_\mu^{(3)}\left(\frac{3}{2}, \frac{3}{2}\right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_\mu^{(3)} & 7 \\
 A_\mu^{(2)}\left(\frac{3}{2}, \frac{3}{2}\right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_\mu^{(2)} & 5 \\
 A_\mu^{(1)}\left(\frac{3}{2}, \frac{3}{2}\right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_\mu^{(1)} & 3 \\
 A_0^{(0)}\left(\frac{3}{2}, \frac{3}{2}\right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_0^{(0)} & 1
 \end{aligned} \tag{8.21}$$

By deleting the element  $A_0^{(0)}\left(\frac{3}{2}, \frac{3}{2}\right)$  one obtains the subalgebra  $su(4)$ . By retaining the 10 elements with  $\lambda = \text{odd}$

$$\begin{aligned}
 A_\mu^{(3)}\left(\frac{3}{2}, \frac{3}{2}\right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_\mu^{(3)} & 7 \\
 A_\mu^{(1)}\left(\frac{3}{2}, \frac{3}{2}\right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_\mu^{(1)} & 3
 \end{aligned} , \tag{8.22}$$

one obtains the algebra of  $sp(4)$ . By retaining only the elements with  $\lambda = 1$

$$A_\mu^{(1)}\left(\frac{3}{2}, \frac{3}{2}\right) = \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_\mu^{(1)} \quad 3 \tag{8.23}$$

one obtains the algebra of  $spin(3) \sim su(2)$ . Finally by retaining only

$$A_0^{(1)}\left(\frac{3}{2}, \frac{3}{2}\right) = \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_0^{(1)} \quad 1 \tag{8.24}$$

one obtains  $spin(2)$ . A basis for fermions with  $spin \frac{3}{2}$  is then

$$\left| \begin{array}{cccc}
 u(4) \supset & sp(4) & \supset & su(2) \supset spin(2) \\
 \downarrow & & \downarrow & \downarrow \\
 N_F & (n_1, n_2) & J & M_J
 \end{array} \right\rangle . \tag{8.25}$$

The values of  $n_1, n_2$  are restricted by the branching rules and only one quantum number is actually needed. The allowed states are

$\square \equiv [1]$	$N_F = 1$	
$\square$ $\square \equiv [1, 1]$	$N_F = 2$	
$\square$ $\square \equiv [1, 1, 1]$ $\square$	$N_F = 3$	(8.26)
$\square$ $\square$ $\square \equiv [1, 1, 1, 1]$ $\square$	$N_F = 4$	

The classification of antisymmetric states is given in Table 8.2. The number

**Table 8.2.** Classification of antisymmetric states of  $u(4)$

$j = 3/2$	$u(4)$	$sp(4)$	$su(2)$
	[0]	(0, 0)	0
	[1]	(1, 0)	3/2
	[1, 1]	(0, 0)	0
		(1, 1)	2
	[1, 1, 1]	(1, 0)	3/2
	[1, 1, 1, 1]	(0, 0)	0

$N_F$  is here

$$N_F = \sum_{i=1}^4 \lambda_i . \tag{8.27}$$

### 8.4.4 The Algebra $u(6)$

This algebra can be constructed with  $j = \frac{5}{2}$ . The 36 elements have the form

$$\begin{aligned}
 A_\mu^{(5)} \left( \frac{5}{2}, \frac{5}{2} \right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_\mu^{(5)} & 11 \\
 A_\mu^{(4)} \left( \frac{5}{2}, \frac{5}{2} \right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_\mu^{(4)} & 9 \\
 A_\mu^{(3)} \left( \frac{5}{2}, \frac{5}{2} \right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_\mu^{(3)} & 7 \\
 A_\mu^{(2)} \left( \frac{5}{2}, \frac{5}{2} \right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_\mu^{(2)} & 5 \\
 A_\mu^{(1)} \left( \frac{5}{2}, \frac{5}{2} \right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_\mu^{(1)} & 3 \\
 A_0^{(0)} \left( \frac{5}{2}, \frac{5}{2} \right) &= \left[ a_{3/2}^\dagger \times \tilde{a}_{3/2} \right]_0^{(0)} & 1
 \end{aligned} . \tag{8.28}$$

**Table 8.3.** Classification of antisymmetric states of  $u(6)$

$j = 5/2$	$u(6)$	$sp(6)$	$su(2)$
	[0]	(0, 0, 0)	0
	[1]	(1, 0, 0)	5/2
	[1, 1]	(0, 0, 0)	0
		(1, 1, 0)	2, 4
	[1, 1, 1]	(1, 0, 0)	5/2
		(1, 1, 1)	3/2, 9/2
	[1, 1, 1, 1]	(0, 0, 0)	0
		(1, 1, 0)	2, 4
	[1, 1, 1, 1, 1]	(1, 0, 0)	5/2
	[1, 1, 1, 1, 1, 1]	(0, 0, 0)	0

The basis  $\mathcal{F}$  can be labeled by

$$\left\langle \begin{array}{cccc} u(6) \supset & sp(6) & \supset & su(2) \supset spin(2) \\ \downarrow & \downarrow & \downarrow & \downarrow \\ N_F & (n_1, n_2, n_3) & J & M_J \end{array} \right\rangle. \tag{8.29}$$

The representations  $(n_1, n_2, n_3)$  of  $sp(6)$  are restricted and only one quantum number is needed to classify uniquely the states. The classification scheme is given in Table 8.3. The representations of  $u(6)$  are labelled here by their Young tableau  $[\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5, \lambda_6]$ . The number of fermions  $N_F$  is

$$N_F = \sum_{i=1}^6 \lambda_i. \tag{8.30}$$

Algebras with  $j > \frac{5}{2}$  and their classification scheme can be constructed in a similar way. [M. Hamermesh, *Group Theory and its Applications to Physical Problems*, Addison-Wesley, Reading, Massachusetts, 1962.]

### 8.5 The Algebra $u(\sum_i (2j_i + 1))$

A generalization of the construction given in the previous section, is the case when there are several values of  $j$ . These situations occur in atomic and nuclear physics where they are called mixed configurations. The construction is straightforward and it produces the Lie algebra  $u(n)$ , with  $n = \sum_i (2j_i + 1)$ . Note, however, once more, that with fermion operators that transform as representations of  $spin(3) \supset spin(2)$  it is possible to construct only unitary algebras in an even number of dimensions.

**Example 1.** *The algebras  $u(6)$  and  $u(12)$ .*

In the study of the spectroscopy of nuclei, one encounters situations in which the values of  $j$  are  $j = \frac{1}{2}, \frac{3}{2}$  or  $j = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}$ . The algebras constructed with these values are  $u(6)$  and  $u(12)$ . [F. Iachello and P. van Isacker, *The Interacting Boson-Fermion Model*, Cambridge University Press, 1991].

## 8.6 Internal Degrees of Freedom (Different Spaces)

### 8.6.1 The Algebras $u(4)$ and $su(4)$

One often encounters in physics particles with internal degrees of freedom. These objects transform as representation  $|j, m\rangle$  under  $spin(3) \supset spin(2)$ , and as representations of some internal symmetry group,  $G$ . A particularly interesting case is that of protons and neutrons. These particles transform as  $j = \frac{1}{2}$  under  $spin(3)$  and as  $t = \frac{1}{2}$  under another group called  $isospin(3)$ . The fermion creation operators are, in a double index notation:

$$\begin{aligned} a_{\frac{1}{2}, +\frac{1}{2}, \frac{1}{2}, +\frac{1}{2}}^\dagger & p\uparrow \\ a_{\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, +\frac{1}{2}}^\dagger & p\downarrow \\ a_{\frac{1}{2}, +\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}}^\dagger & n\uparrow \\ a_{\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}}^\dagger & n\downarrow \end{aligned} \quad (8.31)$$

generically denoted  $a_{s, m_s, t, m_t}^\dagger$ . The annihilation operators are  $a_{s, m_s, t, m_t}$ . The 16 bilinear products of creation and annihilation operators form an algebra, called Wigner  $u(4)$ . It is convenient to introduce the adjoint operators

$$\tilde{a}_{s, m_s, t, m_t} = (-)^{s-m_s+t-m_t} a_{s, -m_s, t, -m_t} \quad (8.32)$$

that transform as tensors under spin and isospin rotations. The Racah form of Wigner  $u(4)$  is obtained in terms of double tensors

$$\begin{aligned} \left[ a_{\frac{1}{2}, \frac{1}{2}}^\dagger \times \tilde{a}_{\frac{1}{2}, \frac{1}{2}} \right]_{(M_S, M_T)}^{(S, T)} &= \sum_{\substack{m, m' \\ m_t, m_t'}} \left\langle \frac{1}{2}, m_s, \frac{1}{2}, m'_s \mid S, M_S \right\rangle \\ &\times \left\langle \frac{1}{2}, m_t, \frac{1}{2}, m_{t'} \mid T, M_T \right\rangle a_{\frac{1}{2}, m_s, \frac{1}{2}, m_t}^\dagger \tilde{a}_{\frac{1}{2}, m'_s, \frac{1}{2}, m'_t} \end{aligned} \quad (8.33)$$

with  $S = 0, 1$  and  $T = 0, 1$ .

### Subalgebra Chain

Although a given algebra can be decomposed into several subalgebra chains, as shown in the previous chapter for bosonic realizations, in practice physical

considerations dictate what subalgebra chains are of interest. For algebras with internal degrees of freedom, the construction of subalgebra chains splits into two categories: different spaces and same spaces. For different spaces, the internal degrees of freedom are separated from the outset and not combined at any stage, while for same spaces, they are combined at some stage into a single algebra, whose elements are the sum of the elements of the two algebras. This procedure is elucidated in the paragraphs below.

In the case of Wigner  $u(4)$ , spin and isospin live on different spaces, the physical space and a fictitious isotopic spin space. It is assumed, on physical grounds, that they cannot be combined, and the appropriate chain is  $u(4) \supset su(4) \supset su_S(2) \oplus su_T(2) \supset spin_S(2) \oplus spin_T(2)$ . Here  $su_S(2) \sim spin(3)$ ,  $su_T(2) \sim isospin(3)$  and the subscript  $S, T$  has been added to distinguish spin from isospin. The basis  $\mathcal{F}$  for Wigner  $u(4)$  can then be written as

$$\left\langle \begin{array}{ccc} u(4) & \supset su_S(2) \oplus su_T(2) \supset spin_S(2) \oplus spin_T(2) \\ \downarrow & \downarrow & \downarrow \\ [\lambda_1, \lambda_2, \lambda_3, \lambda_4] & S, T & M_S, M_T \end{array} \right\rangle. \quad (8.34)$$

The full notation for the representations of  $u(4)$  has been restored and the intermediate step  $su(4)$  has been omitted, since no new quantum number appears. For a given representation of  $u(4)$ , the branching problem can be solved using the techniques of Chap. 4. The chain is non-canonical and requires a building-up process. It has become customary to introduce another notation, called Wigner notation. In this notation, one first goes from  $u(4)$  to  $su(4)$ , using the rules of Chap. 4. The representations of  $su(4)$  are labeled by  $[\lambda'_1, \lambda'_2, \lambda'_3]$  with

$$\lambda'_1 = \lambda_1 - \lambda_4, \lambda'_2 = \lambda_2 - \lambda_4, \lambda'_3 = \lambda_3 - \lambda_4. \quad (8.35)$$

The Wigner quantum numbers  $(P, P', P'')$  are defined as

$$(P, P', P'') = \left( \frac{\lambda'_1 + \lambda'_2 - \lambda'_3}{2}, \frac{\lambda'_1 - \lambda'_2 + \lambda'_3}{2}, \frac{\lambda'_1 - \lambda'_2 - \lambda'_3}{2} \right). \quad (8.36)$$

The branching of representations of  $su(4)$  into representations of  $su_S(2) \oplus su_T(2)$  can then be constructed. [M. Hamermesh, *Group Theory and its Applications to Physical Problems*, Addison-Wesley, Reading, Massachusetts,

**Table 8.4.** Branching  $su(4) \supset su_S(2) \oplus su_T(2)$

$su(4)$ $(P, P', P'')$	$\supset su_S(2) \oplus su_T(2)$ $(S, T)$	$\dim(P, P', P'')$
$(0, 0, 0)$	$(0, 0)$	1
$(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$	$(\frac{1}{2}, \frac{1}{2})$	4
$(1, 1, 1)$	$(0, 0), (1, 1)$	10
$(\frac{3}{2}, \frac{3}{2}, \frac{3}{2})$	$(\frac{1}{2}, \frac{1}{2}), (\frac{3}{2}, \frac{3}{2})$	20



1962).] A portion of this branching is given in Table 8.4. In the last column the dimension of the representations of  $su(4)$  is shown.

### 8.6.2 The Algebras $u(6)$ and $su(6)$

In this case, particles still transform as  $j = \frac{1}{2}$  under  $spin(3)$ , but they now transform as the fundamental representation [1] of an internal group  $su_F(3)$ , called the flavor group. The creation operators are denoted by  $a_{u\uparrow}^\dagger, a_{d\uparrow}^\dagger, a_{s\uparrow}^\dagger, a_{u\downarrow}^\dagger, a_{d\downarrow}^\dagger, a_{s\downarrow}^\dagger$ . The index  $u, d, s$  stands for  $u, d, s$  quarks. The bilinear product of creation and annihilation operators form a  $u(6)$  algebra, called the Gürsey-Radicati algebra.

#### Subalgebra Chain

The breaking of  $u(6)$  first proceeds to  $su_F(3) \oplus su_S(2)$ . The algebra  $su(3)$  can be broken in various ways. The breaking appropriate to the physical situation described by the Gürsey-Radicati algebra is  $su_F(3) \supset su_T(2) \oplus u_Y(1)$ , where  $T$  denotes the isospin as before, and  $u_Y(1)$  is an Abelian algebra called hypercharge (Y) algebra. The subalgebra chain is thus  $u(6) \supset su(6) \supset su_F(3) \oplus su_S(2) \supset su_T(2) \oplus u_Y(1) \oplus su_S(2) \supset spin_T(2) \oplus u_Y(1) \oplus spin_S(2)$ . The basis is labelled by

$$\left. \begin{array}{l} u(6) \supset su_F(3) \oplus su_S(2) \supset su_T(2) \oplus u_Y(1) \oplus su_S(2) \\ \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \\ [\lambda] \qquad [\mu_1, \mu_2] \qquad S \qquad T \qquad Y \\ \supset spin_T(2) \oplus u_Y(1) \oplus spin_S(2) \\ \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \\ \qquad \qquad \qquad T_3 \qquad \qquad \qquad S_3 \end{array} \right\}. \quad (8.37)$$

Here  $[\lambda] = [\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5, \lambda_6]$ . It has become customary to label the representations of  $su_F(3)$  not by their Young labels  $[\mu_1, \mu_2]$  but by the dimension of the representations,  $\dim [0, 0] = 1, \dim [2, 1] = 8$ , etc. and the representations of  $su_S(2)$  by their dimensions  $\dim S = 2S + 1$ , put as a superscript, that is  ${}^8$  denotes the representation  $[2, 1], S = \frac{1}{2}$  of  $su_F(3) \oplus su_S(2)$ . The branching of representations of  $u(6)$  into representations of  $su_F(3) \oplus su_S(2)$  can be constructed. [F. Gürsey and L.A. Radicati, *Spin and unitary spin independence of strong interactions*, Phys. Rev. Lett. 13, 173 (1964).] A portion of this branching is given in Table 8.5. The conversion between this notation and the standard notation for  $su(3)$  representations used in the preceding chapters is  $1 \equiv [0, 0], 3 \equiv [1, 0], \bar{3} \equiv [1, 1], 6 \equiv [2, 0], 8 \equiv [2, 1], 10 \equiv [3, 0]$ . The general formula for the dimension of the representations of  $u(n)$  is given in Chap. 4, Sect. 12. It is worth noting that with fermions with internal degrees of freedom, it is possible to construct algebras in both even and odd number of dimensions, in the present case  $su_F(3)$ .

**Table 8.5.** Branching  $su(6) \supset su_F(3) \oplus su_T(2)$

$su(6)$ $[\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5]$	$\supset$	$su_F(3) \oplus su_S(2)$ $(\dim [\mu_1, \mu_2], \dim S)$	$\dim [\lambda]$
$[0, 0, 0, 0, 0]$		$(1, 1)$	1
$[1, 0, 0, 0, 0]$		$(3, 2)$	6
$[1, 1, 1, 1, 1]$		$(\bar{3}, 2)$	6
$[1, 1, 0, 0, 0]$		$(\bar{3}, 3), (6, 1)$	15
$[1, 1, 1, 0, 0]$		$(8, 2), (1, 4)$	20
$[2, 0, 0, 0, 0]$		$(6, 3), (\bar{3}, 1)$	21
$[2, 1, 1, 1, 1]$		$(8, 3), (8, 1), (1, 3)$	35
$[3, 0, 0, 0, 0]$		$(10, 4), (8, 2)$	56
$[2, 1, 0, 0, 0]$		$(10, 2), (8, 4), (8, 2), (1, 2)$	70

## 8.7 Internal Degrees of Freedom (Same Space)

### 8.7.1 The Algebra $u((2l + 1)(2s + 1))$ : $L$ - $S$ Coupling

Another case of interest is that of particles with both orbital,  $l = \text{integer}$ , and spin,  $s = \text{half-integer}$ , angular momentum. (The spin angular momentum is the internal degree of freedom.) The corresponding creation and annihilation operators are

$$\begin{aligned} a_{l, m_l, s, m_s}^\dagger \\ \tilde{a}_{l, m_l, s, m_s} \end{aligned} = (-1)^{l-m_l+s-m_s} a_{l, -m_l, s, -m_s} . \tag{8.38}$$

The bilinear products

$$a_{l, m_l, s, m_s}^\dagger \tilde{a}_{l, m'_l, s, m'_s} \tag{8.39}$$

generate the Lie algebra  $u((2l + 1)(2s + 1))$ .

### Racah Form

The Racah form of the orbital-spin algebra is constructed with double tensors as in the preceding section,

$$\begin{aligned} [a_{l, s}^\dagger \times \tilde{a}_{l, s}]_{M_L, M_S}^{(L, S)} &= \sum_{\substack{m_s, m'_s \\ m_l, m'_l}} \langle l, m_l, l, m'_l | L, M_L \rangle \langle s, m_s, s, m'_s | S, M_S \rangle \\ &\times a_{l, m_l, s, m_s}^\dagger \tilde{a}_{l, m'_l, s, m'_s} . \end{aligned} \tag{8.40}$$

The classification scheme for this algebra is

$$\begin{aligned} u((2l + 1)(2s + 1)) &\supset u_L(2l + 1) \oplus su_S(2s + 1) \supset \dots \\ &\supset so_L(3) \oplus su_S(2) \supset spin_J(3) \supset spin_J(2) , \end{aligned} \tag{8.41}$$

called  $L$ - $S$  coupling. Since orbital and spin angular momentum are assumed to act on the same space, they can be combined. Here  $spin_J(3)$  denotes the algebra whose elements are the sum of the elements of  $so_L(3)$  and  $su_S(2)$ , sometimes called the diagonal algebra. For large  $l$ , some intermediate steps are needed between  $u_L(2l+1)$  and  $so_L(3)$  and the problem of missing labels becomes particularly acute since the representations of  $u_L(2l+1)$  that appear in (8.41) are not just one-row representations as in the case of bosons. For this reason, dots have been inserted in (8.41) between  $u_L(2l+1)$  and  $so_L(3)$ .

The representations of  $u((2l+1)(2s+1))$  that describe fermions are the totally antisymmetric representations  $\{N_F\}$  of (8.7) (one-column representations). Racah showed that, in order to obtain totally antisymmetric representations of  $u((2l+1)(2s+1))$ , the representations of  $u_L(2l+1)$  and  $su_S(2s+1)$  must be *dual* (sometimes called *conjugate*), that is obtained from each other by interchanging columns with rows. Particularly interesting is the case of  $s = \frac{1}{2}$  that describes electrons in atoms and nucleons in nuclei. In this case, the algebra is  $u(2(2l+1)) \supset u_L(2l+1) \oplus su_S(2)$ . The representations of  $su_S(2)$  are characterized by the quantum numbers  $S, M_S$ , and thus described by a Young tableau with two rows of length  $(N/2) + S$  and  $(N/2) - S$ . The representations of  $u_L(2l+1)$  which arise in the reduction of representations  $\{N_F\}$  of  $u(2(2l+1))$  must then be described by a Young tableau with two columns of these lengths.

**Example 2.** *p electrons in atoms*

In this case,  $l = 1$ , leading to the Lie algebra  $u(6)$ . The classification scheme is

$$u(6) \supset u_L(3) \oplus su_S(2) \supset so_L(3) \oplus su_S(2) \supset spin_J(3) \supset spin_J(2) . \quad (8.42)$$

The branching of the representations of  $u_L(3)$  into those of  $so_L(3)$  has been discussed in Chap. 4, Sect. 14. No additional subalgebra is needed.

**Example 3.** *f electrons in atoms*

In this case,  $l = 3$ , and the appropriate algebra is  $u(14)$ . This algebra has 196 elements. The classification scheme of (8.41) is

$$u(14) \supset u_L(7) \oplus su_S(2) \supset \dots \supset so_L(3) \oplus su_S(2) \supset spin_J(3) \supset spin_J(2) . \quad (8.43)$$

The classification scheme for the step  $u_L(7) \supset so_L(3)$  is incomplete. A complete classification scheme was found by Racah and it is

$$u_L(7) \supset so_L(7) \supset g_2 \supset so_L(3) \supset so_L(2) . \quad (8.44)$$

This classification scheme occupies a special role in Lie algebraic methods in physics since it was the first case in which an exceptional algebra,  $g_2$ , appeared. The elements of the algebras in (8.44) can be written easily in the double tensor notation of (8.40). They are

$$\begin{aligned}
 u(7) : & \quad [a^\dagger \times \tilde{a}]_{M_L,0}^{(L,0)} \quad L = 0, 1, 2, 3, 4, 5, 6 \quad 49 \\
 su(7) : & \quad [a^\dagger \times \tilde{a}]_{M_L,0}^{(L,0)} \quad L = 1, 2, 3, 4, 5, 6 \quad 48 \\
 so(7) : & \quad [a^\dagger \times \tilde{a}]_{M_L,0}^{(L,0)} \quad L = 1, 3, 5 \quad 21 \\
 g_2 : & \quad [a^\dagger \times \tilde{a}]_{M_L,0}^{(L,0)} \quad L = 1, 5 \quad 14 \\
 so(3) : & \quad [a^\dagger \times \tilde{a}]_{M_L,0}^{(L,0)} \quad L = 1 \quad 3 \\
 so(2) : & \quad [a^\dagger \times \tilde{a}]_{M_L,0}^{(1,0)} \quad 1
 \end{aligned} \tag{8.45}$$

where the indices  $l = 3, s = \frac{1}{2}$  have been omitted. States are characterized by

$$\left. \begin{array}{cccccc}
 u(7) & \supset & so(7) & \supset & g_2 & \supset & so(3) & \supset & so(2) \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 [\lambda_1, \dots, \lambda_7] & & [\mu_1, \mu_2, \mu_3] & & [\gamma_1, \gamma_2] & & L & & M_L
 \end{array} \right\} . \tag{8.46}$$

For the application described in this example, the representations of  $u(14)$  are characterized by the number of electrons  $N_F$ . The representations of  $su_S(2)$  are characterized by the total spin  $S$ . For each  $N_F$  and  $S$  the representations of  $u(7)$  are two-column representations with length  $(N_F/2)+S$  and  $(N_F/2)-S$ . Tables of reduction for  $u(7) \supset so(7)$ ,  $so(7) \supset g_2$ ,  $g_2 \supset so(3)$  can be found in Racah. Note that the reduction  $u(7) \supset so(7)$  for fermionic representations is different from that of bosonic representations given in Chap. 7. The same remark applies for the reduction  $so(7) \supset g_2$  and  $g_2 \supset so(3)$ . The reduction  $so(7) \supset g_2$  is given in Table 8.6. The reduction  $g_2 \supset so(3)$  is given in Table 8.7.

**Table 8.6.** Reduction  $so(7) \supset g_2$

$so(7)$ $[\mu_1, \mu_2, \mu_3]$	$\supset g_2$ $[\gamma_1, \gamma_2]$
[0, 0, 0]	[0, 0]
[1, 0, 0]	[1, 0]
[1, 1, 0]	[1, 0], [1, 1]
[2, 0, 0]	[2, 0]
[1, 1, 1]	[0, 0], [1, 0], [2, 0]
[2, 1, 0]	[1, 1], [2, 0], [2, 1]
[2, 1, 1]	[1, 0], [1, 1], [2, 0], [2, 1], [3, 0]
[2, 2, 0]	[2, 0], [2, 1], [2, 2]
[2, 2, 1]	[1, 0], [1, 1], [2, 0], [2, 1], [3, 0], [3, 1]
[2, 2, 2]	[0, 0], [1, 0], [2, 0], [3, 0], [4, 0]

**Table 8.7.** Reduction  $g_2 \supset so(3)$

$g_2$	$\supset so(3)$
$[\gamma_1, \gamma_2]$	$L$
$[0, 0]$	0
$[1, 0]$	3
$[1, 1]$	1, 5
$[2, 0]$	2, 4, 6
$[2, 1]$	2, 3, 4, 5, 7, 8
$[3, 0]$	1, 3, 4, 5, 6, 7, 9
$[2, 2]$	0, 2, 4, 5, 6, 8, 10
$[3, 1]$	1, 2, 3 <sup>2</sup> , 4, 5 <sup>2</sup> , 6 <sup>2</sup> , 7 <sup>2</sup> , 8, 9, 10, 11
$[4, 0]$	0, 2, 3, 4 <sup>2</sup> , 5, 6 <sup>2</sup> , 7, 8 <sup>2</sup> , 9, 10, 12

To complete the study of this chain, one needs also to construct the Casimir invariants and their eigenvalues. The construction is given in Racah. The general expression for the eigenvalues in the representation  $[\gamma_1, \gamma_2]$  are

$$\langle C_2(g_2) \rangle = (\gamma_1^2 + \gamma_1\gamma_2 + \gamma_2^2 + 5\gamma_1 + 4\gamma_2) / 12 . \tag{8.47}$$

(The factor of 12 comes from the definition of  $C_2(g_2)$ .) A detailed account is given in [G. Racah, *Theory of Complex Spectra. IV*, Phys. Rev. 76, 1352 (1949).] Also, note again that with fermions with internal degrees of freedom it is possible to construct unitary algebras in odd number of dimensions, here  $u(7)$ . Finally,  $L - S$  coupling plays a major role in the spectroscopy of atoms.

### 8.7.2 The Algebra $u\left(\sum_j (2j + 1)\right)$ : $j$ - $j$ Coupling

The orbital and spin angular momenta can be coupled from the outset. The corresponding creation and annihilation fermion operators are

$$a_{l,s,j,m_j}^\dagger; \quad \tilde{a}_{l,s,j,m_j} = (-1)^{j-m_j} a_{l,s,j,-m_j} , \tag{8.48}$$

with  $|l + s| \geq j \geq |l - s|$ . The bilinear products

$$a_{l,s,j,m_j}^\dagger \tilde{a}_{l,s,j',m'_j}$$

form the Lie algebra  $u(n)$  with  $n = \sum_j (2j + 1)$ , as in Chap. 5. The Racah form of the algebra and its branchings can be constructed. [F. Iachello and P. Van Isacker, *The Interacting Boson-Fermion Model*, Cambridge University Press, 1991.] Again here the case of interest is that in which  $s = \frac{1}{2}$ .

**Example 4.**  $p$  nucleons in nuclei

In this case,  $l = 1$ , and  $j = \frac{1}{2}, \frac{3}{2}$ , leading to the Lie algebra  $u(6)$ . The classification scheme is

$$u(6) \supset sp(6) \supset spin_J(3) \supset spin_J(2) . \tag{8.49}$$

$j$ - $j$  coupling plays a major role in the spectroscopy of nuclei.

### 8.7.3 The Algebra $u((\sum_l(2l+1))(2s+1))$ : Mixed $L$ - $S$ Configurations

In this case the orbital angular momentum  $l$  takes several values  $l_1, l_2, \dots$ . The creation and annihilation operators are still given by (8.48), but the elements of the Lie algebra are the bilinear products

$$a_{l,m_l,s,m_s}^\dagger \tilde{a}_{l',m'_l,s,m'_s} \quad (8.50)$$

where  $l, l'$  take the values  $l_1, l_2, \dots$ . A particularly important case described by this algebra is that of fermions with spin  $s = \frac{1}{2}$  moving in a harmonic oscillator potential in three dimensions. As discussed in Chap. 7, Sect. 4.1.2, for  $N \geq 2$  states of the three dimensional harmonic oscillator are degenerate with respect to the angular momentum. For  $N = 2$  the values of the orbital angular momentum are  $l = 0, 2$ , for  $N = 3$  they are  $l = 1, 3$ , etc. The algebra that describes fermions in a given degenerate shell (a given value of  $N$ ), called Elliott algebra, is  $u((\sum_l(2l+1))(2s+1))$ . [J.P. Elliott, *Collective Motion in the Nuclear Shell Model. I. Classification Schemes for States of Mixed Configurations*, Proc. Roy. Soc. A 245, 128 (1958).]

#### Example 5. $s$ - $d$ nucleons in nuclei

In this case  $l = 0, 2$  leading to the Lie algebra  $u(12)$ . The classification scheme is

$$\begin{aligned} u(12) &\supset u_L(6) \oplus su_S(2) \supset su_L(3) \oplus su_S(2) \supset so_L(3) \oplus su_S(2) \\ &\supset spin_J(3) \supset spin_J(2). \end{aligned} \quad (8.51)$$

Mixed  $L$ - $S$  configurations play a major role in the spectroscopy of light nuclei.

# 9 Differential Realizations

## 9.1 Differential Operators

Lie algebras can also be constructed with differential operators acting on a space of derivable functions  $f(x)$  of the coordinate  $x$ . The basic commutation relations are

$$\left[ x, \frac{d}{dx} \right] = x \frac{d}{dx} - \frac{d}{dx} x = -1 . \quad (9.1)$$

## 9.2 Unitary Algebras $u(n)$

The bilinear products of coordinates,  $x_i (i = 1, \dots, n)$ , and their derivatives,  $\frac{\partial}{\partial x_j} (j = 1, \dots, n)$  generate  $u(n)$

$$u(n) \doteq x_i \frac{\partial}{\partial x_j} . \quad (9.2)$$

Introducing the double index notation of the previous chapters

$$X_{ik} = x_i \frac{\partial}{\partial x_k} \quad i, k = 1, \dots, n \quad (9.3)$$

one obtains the commutation relations

$$[X_{ik}, X_{mn}] = \delta_{km} X_{in} - \delta_{in} X_{mk} . \quad (9.4)$$

For  $su(n)$ , one subtracts  $\sum_j X_{jj}$  from the diagonal elements

$$X'_{ii} = X_{ii} - \frac{1}{n} \sum_j X_{jj} . \quad (9.5)$$

and deletes  $X'_{nn}$ . (The  $n$  elements  $X'_{ii}$  are not linear independent since  $\sum_i X'_{ii} = 0$ .) The  $n^2 - 1$  elements  $X_{ik} (i \neq k = 1, \dots, n)$  and  $X'_{ii} (i = 1, \dots, n - 1)$  generate  $su(n)$ .

**Example 1.** *Differential realization of  $u(2)$*

Introduce two coordinates  $x, y$ . The differential realization is

$$X_{11} = x \frac{\partial}{\partial x}, X_{12} = x \frac{\partial}{\partial y}, X_{21} = y \frac{\partial}{\partial x}, X_{22} = y \frac{\partial}{\partial y}. \quad (9.6)$$

**Example 2.** *Differential realization of  $su(2)$*

From the preceding example, one obtains

$$X'_{11} = \frac{1}{2} \left( x \frac{\partial}{\partial x} - y \frac{\partial}{\partial y} \right), X_{12} = x \frac{\partial}{\partial y}, X_{21} = y \frac{\partial}{\partial x}. \quad (9.7)$$

### 9.3 Orthogonal Algebras $so(n)$

Differential realization are often used to construct orthogonal Lie algebras. Introduce  $n$  real coordinates,  $x_1, x_2, \dots, x_n$ . A construction of the Lie algebra  $so(n)$  is

$$so(n) \doteq x_i \frac{\partial}{\partial x_j} - x_j \frac{\partial}{\partial x_i}. \quad (9.8)$$

Introducing the notation

$$L_{ij} = x_i \frac{\partial}{\partial x_j} - x_j \frac{\partial}{\partial x_i}, \quad i < j = 1, \dots, n. \quad (9.9)$$

one obtains the commutation relations

$$[L_{ij}, L_{kl}] = \delta_{jk} L_{il} + \delta_{il} L_{jk} - \delta_{jl} L_{ik} - \delta_{ik} L_{jl}. \quad (9.10)$$

The  $\frac{n(n-1)}{2}$  elements  $L_{ij}$  are called angular momentum operators.

**Example 3.** *The Lie algebra  $so(2)$*


We introduce here two coordinates,  $x, y$ , as in part (a) of Fig. 9.1. The single element of  $so(2)$  is

$$X_1 = y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y}. \quad (9.11)$$

This element acts on functions  $f(x, y)$ .

**Example 4.** *The Lie algebra  $so(3)$*





**Fig. 9.1.** Coordinates for differential realizations of (a)  $so(2)$  and (b)  $so(3)$

We introduce here three coordinates,  $x, y, z$ , as in part (b) of Fig. 9.1. The three elements of  $so(3)$  are, in a single index notation,

$$\begin{aligned} X_1 &= z \frac{\partial}{\partial y} - y \frac{\partial}{\partial z} \\ X_2 &= x \frac{\partial}{\partial z} - z \frac{\partial}{\partial x} \\ X_3 &= y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} \end{aligned} \tag{9.12}$$

acting of  $f(x, y, z)$ . The elements satisfy the commutation relations

$$[X_1, X_2] = X_3 ; [X_2, X_3] = X_1 ; [X_3, X_1] = X_2 . \tag{9.13}$$

**Example 5.** *The Lie algebra  $so(4)$*

We introduce here four coordinates,  $x, y, z, t$ . The six elements of  $so(4)$  are

$$\begin{aligned} A_1 &= z \frac{\partial}{\partial y} - y \frac{\partial}{\partial z}, A_2 = x \frac{\partial}{\partial z} - z \frac{\partial}{\partial x}, A_3 = y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y}, \\ B_1 &= x \frac{\partial}{\partial t} - t \frac{\partial}{\partial x}, B_2 = y \frac{\partial}{\partial t} - t \frac{\partial}{\partial y}, B_3 = z \frac{\partial}{\partial t} - t \frac{\partial}{\partial z} . \end{aligned} \tag{9.14}$$

acting on  $f(x, y, z, t)$ .

The realization (9.8) is in  $n$  variables  $x_1, x_2, \dots, x_n$ . Another realization is in terms of  $n - 1$  angle variables  $\varphi_1, \varphi_2, \dots, \varphi_{n-1}$ . This realization is obtained by introducing hyperspherical coordinates,  $(x_1, x_2, \dots, x_n) \rightarrow (r, \varphi_{n-1}, \varphi_{n-2}, \varphi_{n-3}, \dots, \varphi_1)$

$$\begin{aligned}
x_1 &= r \sin \varphi_{n-1} \sin \varphi_{n-2} \dots \sin \varphi_2 \sin \varphi_1 \\
x_2 &= r \sin \varphi_{n-1} \sin \varphi_{n-2} \dots \sin \varphi_2 \cos \varphi_1 \\
&\dots \\
x_{n-1} &= r \sin \varphi_{n-1} \cos \varphi_{n-2} \\
x_n &= r \cos \varphi_{n-1}
\end{aligned} \tag{9.15}$$

and setting  $r = 1$  (realization on the unit  $n$  dimensional sphere.)

**Example 6.** *so(2) in polar coordinates*

Introducing

$$\begin{aligned}
x &= r \cos \varphi \\
y &= r \sin \varphi
\end{aligned} \tag{9.16}$$

one has

$$X_1 = \frac{\partial}{\partial \varphi} . \tag{9.17}$$

**Example 7.** *so(3) in spherical coordinates*

Introducing

$$\begin{aligned}
x &= r \sin \vartheta \cos \varphi \\
y &= r \sin \vartheta \sin \varphi , \\
z &= r \cos \vartheta
\end{aligned} \tag{9.18}$$

one has

$$\begin{aligned}
X_1 &= \cos \varphi \frac{\partial}{\partial \vartheta} + \sin \varphi \operatorname{ctg} \vartheta \frac{\partial}{\partial \varphi} \\
X_2 &= -\sin \varphi \frac{\partial}{\partial \vartheta} - \cos \varphi \operatorname{ctg} \vartheta \frac{\partial}{\partial \varphi} . \\
X_3 &= \frac{\partial}{\partial \varphi}
\end{aligned} \tag{9.19}$$

**Example 8.** *so(4) in hyperspherical coordinates*

One needs here three angles,  $\vartheta, \varphi, \psi$ . The corresponding form of the Lie algebra is called Pauli form.

### 9.3.1 Casimir Operators. Laplace-Beltrami Form

All operators are constructed in terms of differential operators. The differential form of the Casimir operators is often referred to as Laplace-Beltrami form.

**Example 9.** *The Laplace-Beltrami form of so(2)*

We have here trivially

$$C_1(so(2)) = \frac{\partial}{\partial\varphi}, \quad C_2(so(2)) = \frac{\partial^2}{\partial\varphi^2}. \quad (9.20)$$

**Example 10.** *The Laplace-Beltrami form of  $so(3)$*

We have here

$$C_2(so(3)) = \frac{1}{\sin^2\vartheta} \frac{\partial^2}{\partial\varphi^2} + \frac{1}{\sin\vartheta} \frac{\partial}{\partial\vartheta} \left( \sin\vartheta \frac{\partial}{\partial\vartheta} \right). \quad (9.21)$$

### 9.3.2 Basis for the Representations

One also needs to construct the basis for the representations. In the realization in terms of angles, the basis is constructed using the canonical chain

$$so(n) \supset so(n-1) \supset \dots \supset so(2). \quad (9.22)$$

**Example 11.** *Basis for  $so(3)$*

In this case one seeks simultaneous eigenfunctions of  $C_2(so(3))$  and  $C_2(so(2))$

$$\begin{aligned} C_2(so(2)) \Phi(\varphi) &= -m^2 \Phi(\varphi) \\ C_2(so(3)) \Theta(\vartheta) &= -l(l+1) \Theta(\vartheta). \end{aligned} \quad (9.23)$$

The eigenfunctions are the spherical harmonics

$$Y_l^m(\vartheta, \varphi) = \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} (-1)^m e^{im\varphi} P_l^m(\cos\vartheta). \quad (9.24)$$

It has become customary in physical applications to introduce the elements  $L_x, L_y, L_z$  obtained from those previously given by multiplication with  $(1/i)$

$$\begin{cases} L_x = \frac{1}{i} \left( \cos\varphi \frac{\partial}{\partial\vartheta} - \sin\varphi \operatorname{ctg}\vartheta \frac{\partial}{\partial\varphi} \right) \\ L_y = \frac{1}{i} \left( -\sin\varphi \frac{\partial}{\partial\vartheta} - \cos\varphi \operatorname{ctg}\vartheta \frac{\partial}{\partial\varphi} \right) \\ L_z = \frac{1}{i} \frac{\partial}{\partial\varphi} \end{cases} \quad (9.25)$$

From these one can also construct the elements in Cartan-Weyl form,  $L_z, L_{\pm} = L_x \pm iL_y$ . The action of those elements on the basis is

$$\begin{cases} L_z Y_l^m(\vartheta, \varphi) = m Y_l^m(\vartheta, \varphi) \\ L_+ Y_l^m(\vartheta, \varphi) = \sqrt{(l-m)(l+m+1)} Y_l^{m+1}(\vartheta, \varphi) \\ L_- Y_l^m(\vartheta, \varphi) = \sqrt{(l+m)(l-m+1)} Y_l^{m-1}(\vartheta, \varphi) \end{cases} . \quad (9.26)$$

The abstract notation is

$$\left. \begin{array}{c} so(3) \supset so(2) \\ \downarrow \quad \downarrow \\ l \quad m \end{array} \right\} \equiv Y_l^m(\vartheta, \varphi) . \quad (9.27)$$

Note that with differential realizations it is possible to construct only tensor representations of the orthogonal algebras. The construction of spinor representations requires the introduction of another mathematical framework, either that of spinors [E. Cartan, *The Theory of Spinors*, Hermann, Paris, 1966] or that of Grassmann variables [F.A. Berezin, *Introduction to Superanalysis*, D. Reidel, Dordrecht, 1987].

## 9.4 Orthogonal Algebras $so(n, m)$

Although the discussion in the preceding chapters has been devoted to compact Lie algebras, it is of interest to consider here non-compact Lie algebras,  $so(n, m)$ . A construction of the Lie algebra  $so(n, m)$  is

$$so(n, m) \doteq \delta_i x_i \frac{\partial}{\partial x_j} - \delta_j x_j \frac{\partial}{\partial x_i} , \quad (9.28)$$

where

$$\delta_i = \begin{cases} 1 & i = 1, \dots, n \\ -1 & i = n+1, \dots, n+m \end{cases} \quad (9.29)$$

**Example 12.** *The Lorentz algebra  $so(3, 1)$*

Introducing coordinates  $x, y, z, t$ , the elements are

$$\begin{aligned} A_1 &= x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x}, A_2 = y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y}, A_3 = z \frac{\partial}{\partial x} - x \frac{\partial}{\partial z}, \\ B_1 &= x \frac{\partial}{\partial t} + t \frac{\partial}{\partial x}, B_2 = y \frac{\partial}{\partial t} + t \frac{\partial}{\partial y}, B_3 = z \frac{\partial}{\partial t} + t \frac{\partial}{\partial z}. \end{aligned} \quad (9.30)$$

### 9.5 Symplectic Algebras $sp(2n)$

A differential realization of  $sp(2n)$  is obtained by introducing  $2n$  coordinates divided into  $x_1, \dots, x_n; x'_1, \dots, x'_n$ . A construction of  $sp(2n)$  is

$$\begin{array}{lll}
 x_i \frac{\partial}{\partial x'_i} & i = 1, \dots, n & n \\
 x'_i \frac{\partial}{\partial x_i} & i = 1, \dots, n & n \\
 x_i \frac{\partial}{\partial x_j} - x'_j \frac{\partial}{\partial x'_i} & i, j = 1, \dots, n & n^2 \\
 x_i \frac{\partial}{\partial x'_j} + x_j \frac{\partial}{\partial x'_i} & i < j = 1, \dots, n & \frac{n(n-1)}{2} \\
 x'_i \frac{\partial}{\partial x_j} + x'_j \frac{\partial}{\partial x_i} & i < j = 1, \dots, n & \frac{n(n-1)}{2}
 \end{array} \quad . \quad (9.31)$$

The number of elements is written to their right. There are in total  $n(2n + 1)$  elements.

**Example 13.** *The Lie algebra  $sp(2)$*

Introducing coordinates  $x_1, x'_1$  one has

$$X_1 = x_1 \frac{\partial}{\partial x'_1}, X_2 = x'_1 \frac{\partial}{\partial x_1}, X_3 = x_1 \frac{\partial}{\partial x_1} - x'_1 \frac{\partial}{\partial x'_1} . \quad (9.32)$$

All the differential realizations discussed above are real forms. [Jin-Qan Chen, *Group Representation Theory for Physicists*, World Scientific, Singapore, 1989.]

# 11 Spectrum Generating Algebras and Dynamic Symmetries

## 11.1 Spectrum Generating Algebras (SGA)

One of the most important applications of Lie algebras has been to the study of physical systems for which the Hamiltonian  $H$  and other operators of physical interest  $T$  can be written in terms of elements  $G_\alpha$  of a Lie algebra  $g$ ,

$$H = f(G_\alpha) \quad G_\alpha \in g \quad (11.1)$$

and

$$T = t(G_\alpha) \quad G_\alpha \in g. \quad (11.2)$$

The Lie algebra  $g$  is then called the Spectrum Generating Algebra (SGA) of the problem. The functionals  $f$  and  $t$  are usually polynomials in the elements  $G_\alpha$ , although cases have been studied in which  $1/H$  rather than  $H$  is a polynomial in the elements of the algebra (the so-called Coulomb problem discussed in Chap. 12). It turns out that most many-body problems, that is Hamiltonian problems for many interacting particles, can be cast in the form of a polynomial expansion in  $G_\alpha$ . The method is thus particularly important for this case. It is convenient to use the double index notation,  $G_{\alpha\beta}$ , of Chaps. 7 and 8. The expansion can then be written as

$$H = E_0 + \sum_{\alpha\beta} \varepsilon_{\alpha\beta} G_{\alpha\beta} + \frac{1}{2} \sum_{\alpha\beta\gamma\delta} u_{\alpha\beta\gamma\delta} G_{\alpha\beta} G_{\gamma\delta} + \dots \quad (11.3)$$

The term linear in the elements is called one-body term, the term quadratic in the elements is called two-body term, etc.. Most theories stop the expansion at two-body terms, although in some cases three- and higher-body terms have been considered. The term  $E_0$  is an overall constant, which sets the zero of the energy, and the coefficients  $\varepsilon_{\alpha\beta}, u_{\alpha\beta\gamma\delta}, \dots$  depend on the physical system under consideration.

## 11.2 Dynamic Symmetries (DS)

A particularly interesting situation occurs when the Hamiltonian  $H$  does not contain all elements of  $g$ , but only those combinations which form the Casimir operators of a chain of algebras originating from  $g \supset g' \supset g'' \supset \dots$

$$H = f(C_i) . \tag{11.4}$$

For these situations, called dynamic symmetries (DS), the eigenvalue problem for  $H$  can be solved in explicit analytic form, since the Casimir operators are diagonal in the basis provided by  $g \supset g' \supset g'' \supset \dots$ . The energy eigenvalues are given by a formula called *energy formula* in terms of the quantum numbers that characterize the representations of  $g \supset g' \supset \dots$ , while the matrix elements of the transition operators  $T$  are given in terms of isoscalar factor of the chain  $g \supset g' \supset g'' \supset \dots$ . When the Hamiltonian  $H$  is linear in the Casimir operators

$$H = E_0 + \alpha C(g) + \alpha' C(g') + \alpha'' C(g'') + \dots \tag{11.5}$$

the energy formula is simply

$$E = \langle H \rangle = E_0 + \alpha \langle C(g) \rangle + \alpha' \langle C(g') \rangle + \alpha'' \langle C(g'') \rangle + \dots , \tag{11.6}$$

where  $\langle \rangle$  denotes expectation value in the appropriate representation. Several examples will be discussed in the following sections. In these examples, taken from the fields of molecular physics, nuclear physics and hadronic physics, the Hamiltonian  $H$  is at most quadratic in the elements of the algebra  $g$ , and hence the Hamiltonian with dynamic symmetry contains at most Casimir operators of order 2. Systems for which the Hamiltonian has a dynamic symmetry are also called *exactly solvable problems*. Dynamic symmetries were introduced in [A.O. Barut and A. Böhm, *Dynamical Groups and Mass Formula*, Phys. Rev. B139, 1107 (1965)] and [Y. Dothan, M. Gell-Mann, and Y. Ne'eman, *Series of Hadron Levels as Representations of Non-compact Groups*, Phys. Lett. 17, 148 (1965)] and subsequently exploited in nuclear physics [F. Iachello, *Dynamical Symmetries in Nuclei*, in *Group Theoretical Methods in Physics*, Ed. A. Böhm, Lange Springer, Berlin, 1979, p.420] and other areas.

### 11.3 Bosonic Systems

Consider a system composed of  $N$  bosons  $b_\alpha (\alpha = 1, 2, \dots, n)$ . The (number conserving) Hamiltonian operator for this system can be written as

$$H = E_0 + \sum_{\alpha\beta} \tilde{\varepsilon}_{\alpha\beta} b_\alpha^\dagger b_\beta + \frac{1}{2} \sum_{\alpha\beta\gamma\delta} u_{\alpha\beta\gamma\delta} b_\alpha^\dagger b_\beta^\dagger b_\gamma b_\delta + \dots . \tag{11.7}$$

After rearrangement of some of the boson operators, this Hamiltonian can be written as above with  $G_{\alpha\beta} = b_\alpha^\dagger b_\beta$  and

$$\varepsilon_{\alpha\beta} = \tilde{\varepsilon}_{\alpha\beta} - \frac{1}{2} \sum_{\gamma} u_{\alpha\gamma\gamma\beta} . \tag{11.8}$$

The bilinear products  $b_\alpha^\dagger b_\beta$  are the elements of the Lie algebra  $u(n)$ . Hence  $u(n)$  is the spectrum generating algebra of this problem. The transition operators

$$T = \sum_{\alpha\beta} t_{\alpha\beta} b_\alpha^\dagger b_\beta + \dots \quad (11.9)$$

can also be written in terms of the elements of the Lie algebra  $u(n)$ . A physical system is characterized by a set of parameters  $\tilde{\varepsilon}_{\alpha\beta}$ ,  $u_{\alpha\beta\gamma\delta}$  and  $t_{\alpha\beta}$ . The methods of the previous chapters can then be used to solve the eigenvalue problem for  $H$ , that is to find the energy spectrum of the system under consideration, and to calculate matrix elements of operators.

Dynamic symmetries of this system can be studied by breaking  $u(n)$  in all possible ways. Often additional conditions are imposed on the breaking of the algebra  $g$  into its subalgebras  $g'$ . In most applications, the system under study is rotationally invariant. Hence the algebra  $so(3)$  must be included in the chain  $g \supset g' \supset g'' \supset \dots$ . The breakings of  $u(n)$  into its subalgebras containing  $so(3)$  were enumerated in Chap. 7. Knowing these breakings one can construct the corresponding dynamic symmetries. Two examples of boson dynamic symmetries will be given here.

### 11.3.1 Dynamic Symmetries of $u(4)$

Consider the algebra  $g \equiv u(4)$  discussed in Chap. 7. This algebra has two subalgebra chains containing  $so(3)$

$$u(4) \begin{cases} u(3) \supset so(3) \supset so(2) & (I) \\ so(4) \supset so(3) \supset so(2) & (II) \end{cases} \quad (11.10)$$

and correspondingly two possible dynamic symmetries.

#### *Dynamic symmetry I*

The Hamiltonian in this case can be written as

$$H^{(I)} = E_0 + \varepsilon C_1(u(3)) + \alpha C_2(u(3)) + \beta C_2(so(3)) . \quad (11.11)$$

The Casimir operator of  $so(2)$  could be added to this Hamiltonian. However, this corresponds physically to placing the system in an external field that splits the degeneracy of the angular momentum. In the absence of external fields, the Casimir operator of  $so(2)$  can be deleted. Also the invariant operators of  $u(4)$  could be included in the overall constant  $E_0$ ,


$$E_0 = E_{00} + E_{01} C_1(u(4)) + E_{02} C_2(u(4)) . \quad (11.12)$$

In the representation  $[N]$  of  $u(4)$  appropriate to bosons,

$$E_0 = E_{00} + E_{01} N + E_{02} N(N+3) \quad (11.13)$$

is a constant for all states in the basis.





**Fig. 11.1.** The energy level diagram of the dynamic symmetries of  $u(4)$ : (a)  $u(4) \supset u(3) \supset so(3) \supset so(2)$ ; (b)  $u(4) \supset so(4) \supset so(3) \supset so(2)$ . The representation [4] is shown

The eigenvalues of  $H^{(I)}$  in the basis  $|N, n_p, L, M_L\rangle$  can simply be found from the eigenvalues of the Casimir operators given in Chap. 5,

$$E^{(I)}(N, n_p, L, M_L) = E_0 + \varepsilon n_p + \alpha n_p(n_p + 2) + \beta L(L + 1) . \quad (11.14)$$

Usually, a hierarchy of couplings occurs, in such a way that successive splittings are smaller and smaller,  $|\varepsilon| \gg |\alpha| \gg |\beta|$ . It is customary to show the spectrum in an *energy level diagram*. The energy level diagram of the dynamic symmetry I of  $u(4)$ , with  $\varepsilon > 0$ , is shown in the top part of Fig. 11.1. This is the spectrum of a truncated three dimensional anharmonic oscillator. When  $\alpha = \beta = 0$ , the spectrum is called harmonic. The spectrum is truncated, since according to the branching rules of Chap. 7,  $n_p \leq N$ .


*Dynamic symmetry II*

The Hamiltonian for this case is

$$H^{(II)} = E_0 + AC_2(so(4)) + BC_2(so(3)) , \quad (11.15)$$

where again the Casimir operator of  $so(2)$  has been deleted. The eigenvalues of  $H^{(II)}$  in the basis  $|N, \omega, L, M_L\rangle$  are

$$E^{(II)}(N, \omega, L, M_L) = E_0 + A\omega(\omega + 2) + BL(L + 1) . \quad (11.16)$$



**Fig. 11.2.** A diatomic molecule is shown as an example of a truncated three-dimensional rotovibrator

Usually  $|A| \gg |B|$ . The energy level diagram of this dynamic symmetry, when  $A < 0$ , is shown in the bottom portion of Fig. 11.1. It represents the spectrum of the truncated three dimensional rotovibrator. In the figure, the usual vibrational quantum number is also shown. A three-dimensional rotovibrator is an object, such as a diatomic molecule, which can vibrate and rotate around an axis perpendicular to the line joining the two atoms, as shown in Fig. 11.2. This spectrum is characterized by a set of states belonging to the same representation of  $so(4)$ , called a rotational band. Different representations of  $so(4)$  correspond to different vibrational excitations. The spectrum is truncated since, according to the branching rules of Chap. 7,  $\omega \leq N$ .

The two dynamic symmetries represent special situations. In general, the Hamiltonian  $H$  will contain Casimir operators of both chains

$$H = E_0 + \varepsilon C_1(u(3)) + \alpha C_2(u(3)) + AC_2(so(4)) + BC_2(so(3)) . \quad (11.17)$$

The eigenvalues must be obtained numerically. For this most general case, Lie algebraic methods provide a basis upon which the numerical diagonalization is done. In the case of  $u(4)$ , there are two such bases, corresponding to the two dynamic symmetries (I) and (II).

Models based on the algebra of  $u(4)$  and its dynamic symmetries have found many applications in molecular physics. [F. Iachello and R.D. Levine, *Algebraic Theory of Molecules*, Oxford University Press, Oxford, 1995.]

### 11.3.2 Dynamic Symmetries of $u(6)$

For  $g \equiv u(6)$ , the subalgebra chains containing  $so(3)$  are

$$\begin{array}{l}
 u(6) \begin{cases} \nearrow \\ - \\ \searrow \end{cases} \begin{array}{l}
 u(5) \supset so(5) \supset so(3) \supset so(2) \quad (I) \\
 su(3) \supset so(3) \supset so(2) \quad (II) \\
 so(6) \supset so(5) \supset so(3) \supset so(2) \quad (III)
 \end{array}
 \end{array} \quad (11.18)$$

and correspondingly there are three possible dynamic symmetries.

*Dynamic symmetry I*

The Hamiltonian is

$$H^{(I)} = E_0 + \varepsilon C_1(u(5)) + \alpha C_2(u(5)) + \beta C_2(so(5)) + \gamma C_2(so(3)) , \quad (11.19)$$

where again the Casimir operator of  $so(2)$  has been omitted. The eigenvalues in the basis  $|N, n_d, v, n_\Delta, L, M_L\rangle$  are

$$E^{(I)}(N, n_d, v, n_\Delta, L, M_L) = E_0 + \varepsilon n_d + \alpha n_d(n_d + 4) + \beta v(v + 3) + \gamma L(L + 1) . \quad (11.20)$$


Usually  $|\varepsilon| \gg |\alpha| \gg |\beta| \gg |\gamma|$ . The energy level diagram, when  $\varepsilon > 0$ , is shown in Fig. 11.3 (top panel). It represents the spectrum of the truncated five-dimensional anharmonic oscillator. If  $\alpha = \beta = \gamma = 0$ , the spectrum is called harmonic.

*Dynamic symmetry II*


The Hamiltonian is

$$H^{(II)} = E_0 + \kappa C_2(su(3)) + \kappa' C_2(so(3)) . \quad (11.21)$$

Its eigenvalues in the basis  $|N, \lambda, \mu, K, L, M_L\rangle$  are



**Fig. 11.3.** The energy level diagram of the dynamic symmetries of  $u(6)$ : (a)  $u(6) \supset u(5) \supset so(5) \supset so(3) \supset so(2)$ ; (b)  $u(6) \supset su(3) \supset so(3) \supset so(2)$ ; (c)  $u(6) \supset so(6) \supset so(5) \supset so(3) \supset so(2)$ . The representation [3] is shown



**Fig. 11.4.** A nucleus deformed in the shape of an ellipsoid is shown as an example of a truncated five-dimensional rotovibrator

$$E^{(II)}(N, \lambda, \mu, K, L, M_L) = E_0 + \kappa(\lambda^2 + \mu^2 + \lambda\mu + 3\lambda + 3\mu) + \kappa' L(L+1). \quad (11.22)$$

Usually  $|\kappa| \gg |\kappa'|$ . The corresponding energy level diagram, when  $\kappa < 0$ , is shown in Fig. 11.3 (middle panel). This is the spectrum of the truncated five-dimensional rotovibrator. A five-dimensional rotovibrator is an object, such as an atomic nucleus deformed in the shape of a quadrupole, which can vibrate and rotate around an axis perpendicular to the symmetry axis of the quadrupole, as shown in Fig. 11.4. Each representation  $(\lambda, \mu)$  represents a rotational band. Different representations  $(\lambda, \mu)$  label vibrational excitations.

*Dynamic symmetry III*

The Hamiltonian is

$$H^{(III)} = E_0 + AC_2(so(6)) + BC_2(so(5)) + CC_2(so(3)). \quad (11.23)$$

Its eigenvalues in the basis  $|N, \sigma, \tau, \nu_\Delta, L, M_L\rangle$  are

$$E^{(III)}(N, \sigma, \tau, \nu_\Delta, L, M_L) = E_0 + A\sigma(\sigma+4) + B\tau(\tau+3) + CL(L+1). \quad (11.24)$$

Usually  $|A| \gg |B| \gg |C|$ . The energy level diagram when  $A < 0$  is shown in Fig. 11.3 (bottom panel). It is called the five dimensional  $\gamma$ - unstable rotovibrator.

The three dynamic symmetries represent special situations. In general, the Hamiltonian contains Casimir operators of all three chains

$$H = E_0 + \varepsilon C_1(u(5)) + \alpha C_2(u(5)) + \beta C_2(so(5)) + \gamma C_2(so(3)) \\ + \kappa C_2(su(3)) + AC_2(so(6)) \quad (11.25)$$

and must be diagonalized numerically. Models based on the algebra  $u(6)$  and its dynamic symmetries have had many applications in nuclear physics. [F. Iachello and A. Arima, *The Interacting Boson Model*, Cambridge University Press, Cambridge, 1987.]

## 11.4 Fermionic Systems

Spectrum generating algebras and dynamic symmetries can also be used for fermions. Consider a system composed of  $N_F$  fermions  $a_i$  ( $i = 1, 2, \dots, n$ ). The number conserving Hamiltonian operator for this system can be written as

$$H = E_0 + \sum_{ii'} \tilde{\eta}_{ii'} a_i^\dagger a_{i'} + \frac{1}{2} \sum_{ii'kk'} v_{ii'kk'} a_i^\dagger a_{i'}^\dagger a_k a_{k'} + \dots \quad (11.26)$$

After rearrangement of some of the fermion operators, the Hamiltonian can be rewritten as

$$H = E_0 + \sum_{ii'} \eta_{ii'} G_{ii'} + \frac{1}{2} \sum_{iki'k'} v_{ii'kk'} G_{ik} G_{i'k'} + \dots \quad (11.27)$$

The elements  $G_{ii'}$  span the Lie algebra  $u(n)$ . Hence  $u(n)$  is the spectrum generating algebra of this problem. All operators can be expanded into elements of this algebra. The transition operators

$$T = \sum_{ii'} t_{ii'} a_i^\dagger a_{i'} + \dots \quad (11.28)$$

are written as

$$T = \sum_{ii'} t_{ii'} G_{ii'} + \dots \quad (11.29)$$

A physical system is characterized by a set of parameters  $\eta_{ii'}$ ,  $v_{ii'kk'}$  and  $t_{ii'}$ . One can use the methods of the previous chaps. to solve the eigenvalue problem for  $H$  and to calculate the matrix elements of operators.

Dynamic symmetries of these systems can be studied by breaking  $u(n)$  in all possible ways. There are two types of problems, the atomic and nuclear many body problem in which the Hamiltonian is rotationally invariant and thus  $so(3)$  must be included in the chain  $g \supset g' \supset g'' \supset \dots$ , and other problems in which one treats internal degrees of freedom and thus rotational invariance need not to be imposed. The former problem is identically to that discussed in Chap. 3. In the following subsections two examples will be given of the latter problem.

### 11.4.1 Dynamic Symmetry of $u(4)$

The Wigner algebra  $u(4)$  is described by the chain

$$u(4) \supset su(4) \supset su_T(2) \oplus su_S(2) \supset spin_T(2) \oplus spin_S(2) \quad (11.30)$$

The basis states can be denoted by  $|(P, P', P''); T, S; M_T, M_S\rangle$ . The Hamiltonian with dynamic symmetry is

$$H = E_0 + aC_2(su(4)) + bC_2(su_T(2)) + cC_2(su_S(2)) + dC_1(spin_T(2)) . \quad (11.31)$$

In this Hamiltonian, the Casimir operator of  $spin_S(2)$  is not included, unless the system is placed in an external field that splits the  $su_S(2)$  degeneracy, but the Casimir operator of  $spin_T(2)$  is included. This group acts on an abstract isotopic spin space. In this space, there are interactions (for example the electromagnetic interaction) that split the  $su_T(2)$  degeneracy. The eigenvalues of the Hamiltonian are

$$E(P, P', P''; T, S; M_T, M_S) = E_0 + a(P^2 + 4P + P'^2 + P''^2) \\ + bT(T + 1) + cS(S + 1) + dM_T . \quad (11.32)$$

A simple case is when  $b = c = d = 0$ . [P. Franzini and L.A. Radicati, *On the Validity of the Supermultiplet Model*, Phys. Lett. 6, 32 (1963)].

### 11.4.2 Dynamic Symmetry of $u(6)$

The Gürsey-Radicati  $u(6)$  is described by the chain

$$u(6) \supset su(6) \supset su_F(3) \oplus su_S(2) \supset su_T(2) \oplus u_Y(1) \oplus su_S(2) \\ \supset spin_T(2) \oplus u_Y(1) \oplus spin_S(2) . \quad (11.33)$$


The basis states are denoted by  $|\dim[\lambda];^{\dim S} \dim[\mu_1, \mu_2]; T, Y, S; M_T, M_S\rangle$ . For applications to elementary particle physics, the mass operator  $M$  rather than the Hamiltonian  $H$  is expanded into the elements of a Lie algebra. The mass operator with dynamic symmetry is written in terms of Casimir invariants as

$$M = M_0 + aC_1(u_Y(1)) + b \left[ C_2(su_T(2)) - \frac{1}{4}C_1(u_Y(1)) \right] \\ + cC_2(su_S(2)) + dC_1(spin_T(2)) , \quad (11.34)$$

with eigenvalues

$$M([\lambda], [\mu_1, \mu_2]; T, Y, S; M_T, M_S) = M_0 + aY + b \left[ T(T + 1) - \frac{1}{4}Y^2 \right] \\ + cS(S + 1) + dM_T . \quad (11.35)$$

This formula is called a *mass formula* as it gives the masses of the particles. The mass level diagram for the representation  $[\lambda] = [3, 0, 0, 0, 0]$  with  $\dim[\lambda] = 56$  is shown in Fig. 11.5. [F. Gürsey and L.A. Radicati, *Spin and Unitary Spin Dependence of Strong Interactions*, Phys. Rev. Lett. 13, 173 (1964).] The  $su_F(3)$  flavor part of this formula, called the Gell-Mann-Okubo *mass*



**Fig. 11.5.** The mass level diagram of the representation [3] of  $u(6)$  with dimension 56. States are labeled by quantum numbers and by names of particles to which they correspond

*formula* was the first explicit example of dynamic symmetry in physics [M. Gell-Mann, *Symmetries of Baryons and Mesons*, Phys. Rev. 125, 1067 (1962); S. Okubo, *Note on Unitary Symmetry in Strong Interactions*, Progr. Theor. Phys. 27, 222 (1962)].

# 10 Matrix Realizations

## 10.1 Matrices

Lie algebras can be constructed with  $n \times n$  square matrices

$$A = \begin{pmatrix} \dots \\ \dots \\ \dots \end{pmatrix} \quad (10.1)$$

acting on the right on column vectors with  $n$  rows

$$\chi = \begin{pmatrix} \cdot \\ \cdot \\ \cdot \end{pmatrix}, \quad (10.2)$$

and on the left on row vectors with  $n$  columns

$$\tilde{\chi} = (\cdot \cdot \cdot). \quad (10.3)$$

Properties of matrices have been given in Chap. 3.

## 10.2 Unitary Algebras $u(n)$

The matrices

$$E_{km} = \begin{pmatrix} 0 & & & & & \\ & 0 & & & & \\ & & 0 & & & \\ 0 & 0 & 0 & 1 & 0 & 0 \\ & & & 0 & & \\ & & & 0 & & \\ & & & 0 & & \end{pmatrix} \quad (10.4)$$

with 1 in the  $k$ -th row and  $m$ -th column and zero otherwise, satisfy the commutation relations of  $u(n)$  [and  $gl(n)$ ]

$$[E_{km}, E_{st}] = \delta_{sm} E_{kt} - \delta_{kt} E_{sm}. \quad (10.5)$$

For  $su(n)$ , one subtracts the unit matrix,  $I$ , divided by  $n$ ,



$$I = \begin{pmatrix} 1 & & & 0 \\ & 1 & & \\ & & \cdots & \\ & & & 1 \\ 0 & & & & 1 \end{pmatrix} \quad (10.6)$$

from  $E_{kk}$

$$E'_{kk} = E_{kk} - \frac{1}{n}I \quad (10.7)$$

and deletes  $E'_{nn}$ . The remaining  $n^2 - 1$  matrices are the elements of  $su(n)$ . This procedure provides real forms of  $u(n)$  and  $su(n)$ .

**Example 1.** *Matrix realization of  $u(2)$*

A real matrix realization of  $u(2)$  is

$$\begin{aligned} E_{11} &= \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, & E_{12} &= \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \\ E_{21} &= \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, & E_{22} &= \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}. \end{aligned} \quad (10.8)$$

**Example 2.** *Matrix realization of  $su(2)$*

A real matrix realization of  $su(2)$  is

$$\begin{aligned} E'_{11} &= \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, & E_{12} &= \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \\ E_{21} &= \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}. \end{aligned} \quad (10.9)$$

It is of interest in physics to introduce also complex forms of the algebras  $u(n)$  and  $su(n)$ . A common construction is: (i) Retain  $E_{kk}$  for  $u(n)$  and  $E'_{kk}$  for  $su(n)$  for the elements with  $k = m$ . (ii) Take the combinations  $(E_{km} + E_{mk})$ ,  $-i(E_{km} - E_{mk})$  for the elements with  $k \neq m$ . This gives the complex form of  $u(n)$

$$\begin{aligned} E_{kk} & & k &= 1, \dots, n \\ E_{+,km} &= E_{km} + E_{mk} & k < m &= 1, \dots, n & \frac{n(n-1)}{2} \\ E_{-,km} &= -i(E_{km} - E_{mk}) & k < m &= 1, \dots, n & \frac{n(n-1)}{2} \end{aligned} \quad (10.10)$$

The number of elements of each type is written to the far right.

**Example 3.** *Complex form of  $u(2)$*

The complex form of  $u(2)$  is

$$\begin{aligned} E_{11} &= \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, & E_{22} &= \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \\ E_{+,12} &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, & E_{-,12} &= \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}. \end{aligned} \quad (10.11)$$

**Example 4.** *Complex form of  $su(2)$* 

This algebra is composed of three elements

$$\begin{aligned} E_{+,12} &= E_{12} + E_{21} \\ E_{-,12} &= -i(E_{12} - E_{21}) \\ E'_{11} &= E_{11} - \frac{1}{2}I. \end{aligned} \quad (10.12)$$

The corresponding matrices, denoted by  $\sigma_x, \sigma_y, \sigma_z$ , are

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (10.13)$$

The last matrix,  $E'_{11}$ , has been multiplied by 2 to conform with the usual normalization of  $\sigma_z$ . They are called Pauli matrices and are widely used in quantum mechanics. The Pauli matrices, together with the unit matrix,  $I$ ,

$$I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (10.14)$$

that is the matrices of  $u(2)$ , form also another algebra, not discussed here, called a Clifford algebra.

**Example 5.** *Complex form of  $u(3)$* 

The complex form of  $u(3)$  is

$$\begin{aligned} E_{+,12} &= \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & E_{-,12} &= \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & E_{+,13} &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\ E_{-,13} &= \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, & E_{+,23} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, & E_{-,23} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \\ E_{11} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & E_{22} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & E_{33} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \end{aligned} \quad (10.15)$$

**Example 6.** *Complex form of  $su(3)$*

In the complex form of  $su(3)$  the first six matrices remain the same

$$\begin{aligned} X_1 &= \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, X_2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, X_3 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\ X_4 &= \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, X_5 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, X_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}. \end{aligned} \quad (10.16)$$

The last three matrices are replaced by the two matrices

$$X_7 = \frac{1}{3} \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, X_8 = \frac{1}{3} \begin{pmatrix} -1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{pmatrix}. \quad (10.17)$$

Note that the choice of the traceless matrices is not unique. In applications, often different choices are made. A commonly used choice is

$$X_7 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, X_8 = \sqrt{\frac{1}{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}. \quad (10.18)$$

The corresponding matrices are called Gell-Mann matrices.

### 10.3 Orthogonal Algebras $so(n)$

A real matrix realization of the algebras  $so(n)$  with square  $n \times n$  matrices is

$$L_{km} = E_{km} - E_{mk} \quad k < m, \quad (10.19)$$

where the matrices  $E_{km}$  are given above.

**Example 7.** *Matrix realization of  $so(2)$*

A real matrix realization of  $so(2)$  is

$$L_{12} = E_{12} - E_{21} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (10.20)$$

This algebra is Abelian, and can be written in many ways.

**Example 8.** *Matrix realization of  $so(3)$*

A real matrix realization of  $so(3)$  is

$$L_{12} = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, L_{13} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, L_{23} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}. \quad (10.21)$$

In applications in quantum mechanics, the complex form  $iL_{km}$  is used. However, matrix realizations of  $so(n)$  are rarely used.

### 10.4 Symplectic Algebras $sp(2n)$

The construction of symplectic algebras is more involved. It is necessary here to introduce  $2n \times 2n$  matrices

$$E_{k,m} = \begin{pmatrix} & & & 0 \\ & & & 0 \\ 0 & 0 & \cdots & 0 & 1 & 0 & 0 \\ & & & & 0 \\ & & & & \cdots \\ & & & & 0 \\ & & & & 0 \end{pmatrix}. \tag{10.22}$$

The rows and columns are labelled by the indices  $k, m = 1, \dots, n; n + 1, \dots, 2n$ . A real matrix realization of  $sp(2n)$  is

$$\begin{array}{lll} E_{k,n+k} & k = 1, \dots, n & n \\ E_{n+k,k} & k = 1, \dots, n & n \\ E_{k,m} - E_{n+m,n+k} & k, m = 1, \dots, n & n^2 \\ E_{k,n+m} + E_{m,n+k} & k < m = 1, \dots, n & \frac{n(n-1)}{2} \\ E_{n+k,m} + E_{n+m,k} & k < m = 1, \dots, n & \frac{n(n-1)}{2} \end{array}. \tag{10.23}$$

The number of elements of each type is shown to the right.

**Example 9.** *The algebra  $sp(2)$*

In this case  $n = 1$ . The construction gives

$$E_{12} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}; E_{21} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}; E_{11} - E_{22} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \tag{10.24}$$

Note that  $sp(2) \sim su(2)$ . The matrix realization of  $sp(2)$  is identical to that of  $su(2)$  given in (10.9), apart from a normalization of the element  $E'_{11} = \frac{1}{2}(E_{11} - E_{22})$ .

**Example 10.** *The algebra  $sp(4)$*

In this case  $n = 2$ . The construction gives the ten elements

$$\begin{array}{ll} E_{13}; E_{24}; E_{31}; E_{42} & 4 \\ (E_{11} - E_{33}); (E_{12} - E_{43}); (E_{21} - E_{34}); (E_{22} - E_{44}) & 4 \\ (E_{14} + E_{23}); (E_{32} + E_{41}) & 2 \end{array}. \tag{10.25}$$

where

$$E_{13} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \text{ etc.} \tag{10.26}$$

This provides a real form of  $sp(2n)$ .

## 10.5 Basis for the Representation

A basis for the fundamental  $n$  dimensional representation of  $u(n)$  and  $su(n)$  is provided by  $n$ -columns vectors

$$\mathcal{B} : \begin{pmatrix} 1 \\ 0 \\ \cdots \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ \cdots \\ 0 \\ 0 \end{pmatrix}, \dots, \begin{pmatrix} 0 \\ 0 \\ \cdots \\ 0 \\ 1 \end{pmatrix}. \quad (10.27)$$

**Example 11.** *The basis for  $su(2)$*

The basis for the 2-dimensional representation of  $su(2)$  is formed by two columns vectors, often denoted by  $\alpha, \beta$

$$\alpha = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \beta = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (10.28)$$

The matrix realization of the algebra can be re-written in Cartan-Weyl form, introducing the matrices  $\sigma_{\pm} = \frac{1}{2}(\sigma_x \pm i\sigma_y)$ ,

$$\sigma_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \sigma_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (10.29)$$

The action of the elements on the basis is

$$\begin{aligned} \sigma_+ \alpha &= 0 & \sigma_- \alpha &= \beta & \sigma_z \alpha &= \alpha \\ \sigma_+ \beta &= \alpha & \sigma_- \beta &= 0 & \sigma_z \beta &= -\beta \end{aligned} \quad (10.30)$$

One of the most important applications of  $su(2)$  in physics is in the description of particles with spin  $S = \frac{1}{2}$ . The three spin operators,  $S_x, S_y, S_z$  can be simply be written as  $S_k = \frac{1}{2}\sigma_k$ . The “spin” basis was discussed in Chap. 8 where it was denoted by

$$\left| \begin{array}{cc} su(2) \supset & spin(2) \\ \downarrow & \downarrow \\ S = \frac{1}{2} & M_S = \pm \frac{1}{2} \end{array} \right\rangle. \quad (10.31)$$

The relation between the basis here and that discussed in Chap. 8 is

$$\alpha \equiv \left| \frac{1}{2}, +\frac{1}{2} \right\rangle, \beta \equiv \left| \frac{1}{2}, -\frac{1}{2} \right\rangle. \quad (10.32)$$

**Example 12.** *The basis for  $su(3)$*

The basis for the fundamental representation of  $su(3)$  is formed by three column vectors

$$\alpha_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \alpha_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \alpha_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}. \quad (10.33)$$

The matrix realization of  $su(3)$  was given in Example 6. In applications to particle physics the Gell-Mann matrices were used. This basis was discussed in Chap. 8, where it was denoted by

$$\left\langle \begin{array}{ccc} su(3) & \supset & su(2) & \supset & spin(2) \\ \downarrow & & \downarrow & & \downarrow \\ [\lambda_1, \lambda_2] \equiv [1] & & T = 0, \frac{1}{2} & & M_T = 0, \pm \frac{1}{2} \end{array} \right\rangle. \quad (10.34)$$

The relation between the basis (10.33) and that discussed in Chap. 8 is

$$\alpha_1 = \left| [1], \frac{1}{2}, +\frac{1}{2} \right\rangle \equiv u, \alpha_2 = \left| [1], \frac{1}{2}, -\frac{1}{2} \right\rangle \equiv d, \alpha_3 = |[1], 0, 0] \equiv s, \quad (10.35)$$

where the notation  $u, d, s$  used in particle physics has also been included.

For representations of  $u(n)$  and  $su(n)$  other than the fundamental representation, matrix realizations are seldom used.

## 10.6 Casimir Operators

The Casimir operators can be constructed from matrices. For the algebras  $u(n)$  and  $su(n)$ , the unit matrix  $I$  commutes with all elements and thus is a Casimir operator.

**Example 13.** *The Casimir operator of  $su(2)$*

For applications to physics it is of particular interest to construct the Casimir operator of  $su(2)$ . This algebra has only one invariant,  $C_2(su(2))$ . It can be easily shown that the invariant is

$$C_2(su(2)) = \sigma_x^2 + \sigma_y^2 + \sigma_z^2 = 3I. \quad (10.36)$$

(The condition  $\sigma_i^2 = I$  is one of the axioms defining a Clifford algebra.) The eigenvalues of the Casimir operator in the basis are

$$\langle \alpha | \sum_k \sigma_k^2 | \alpha \rangle = 3, \quad \langle \beta | \sum_k \sigma_k^2 | \beta \rangle = 3. \quad (10.37)$$

In application in physics, the algebra  $su(2)$  often describes spin  $\frac{1}{2}$  particles. The eigenvalues of the Casimir invariant in the 2-dimensional representation of  $su(2)$  is

$$\langle C_2(su(2)) \rangle = \frac{3}{4}. \quad (10.38)$$

This value agrees with the eigenvalues of the Casimir operator in the generic representation  $|J, M\rangle$  of  $su(2)$  given in Chap. 8,

$$\langle C_2(su(2)) \rangle = J(J+1), \quad (10.39)$$

since here  $J = S = \frac{1}{2}$ .

# 12 Degeneracy Algebras and Dynamical Algebras

## 12.1 Degeneracy Algebras

Another important application of algebraic methods in physics is to the study of exactly solvable problems in quantum mechanics. Consider quantum mechanics in  $\nu$  dimensions described by the Hamiltonian

$$H = -\frac{\hbar^2}{2m}\nabla^2 + V(\mathbf{r}) \quad (12.1)$$

where  $\nabla^2$  is the Laplace operator and  $\mathbf{r} \equiv (x_1, x_2, \dots, x_\nu)$  denotes a vector in  $\nu$  dimensions with components  $x_1, x_2, \dots, x_\nu$ . This Hamiltonian is obtained from the classical Hamiltonian

$$H = \frac{\mathbf{p}^2}{2m} + V(\mathbf{r}) \quad (12.2)$$

by the usual quantization procedure  $\mathbf{p} \rightarrow \frac{\hbar}{i}\nabla$ . If the Hamiltonian (12.1) can be written in terms of the Casimir operator  $C$  of an algebra  $g$ ,

$$H = f(C) \quad (12.3)$$

the eigenvalue problem for  $H$  can be solved in explicit analytic form,

$$E = \langle f(C) \rangle . \quad (12.4)$$

This situation is a dynamic symmetry, Sect. 11.2, except that only the Casimir operator of  $g$  and not those of the subalgebra chain  $g \supset g' \supset g'' \supset \dots$  appears in (12.3). The representations  $[\lambda]$  of  $g$  still label the eigenstates of the Hamiltonian and the symbol  $\langle \rangle$  denotes expectation value in the representation  $[\lambda]$ . If  $\dim[\lambda] \neq 1$ , more than one state has energy  $E$ . The state is said to be degenerate and the algebra  $g$  is called the *degeneracy algebra*,  $g_c$ , of the problem.

## 12.2 Degeneracy Algebras in $\nu \geq 2$ Dimensions

A particularly interesting class of problems is that of quantum mechanics in  $\nu \geq 2$  dimensions with rotationally invariant potentials,  $V = V(r)$ . Here



$$r = (x_1^2 + x_2^2 + \dots + x_\nu^2)^{1/2}. \quad (12.5)$$

This problem admits two and only two *exactly solvable* cases, the isotropic harmonic oscillator with  $V(r) = \frac{1}{2}kr^2$ , and the Coulomb (or Kepler) problem with  $V(r) = \frac{k}{r}$ .

### 12.2.1 The Isotropic Harmonic Oscillator

The Hamiltonian operator, in units where  $\hbar = m = 1$  and  $k = 1$ , is

$$H = \frac{1}{2}(p^2 + r^2) = \frac{1}{2}(-\nabla^2 + r^2). \quad (12.6)$$

Introducing the bosonic realization of Chap. 7 written in differential form and generalizing the results of Example 3 one can write the linear Casimir operator of  $u(\nu)$  as

$$C_1(u(\nu)) = \frac{1}{2} \sum_{j=1}^{\nu} \left( x_j - \frac{\partial}{\partial x_j} \right) \left( x_j + \frac{\partial}{\partial x_j} \right). \quad (12.7)$$

The basic commutation relations

$$\left[ x_i, \frac{\partial}{\partial x_j} \right] = -\delta_{ij} \quad (12.8)$$

give

$$H = C_1(u(\nu)) + \frac{\nu}{2}. \quad (12.9)$$

The degeneracy algebra of the  $\nu$ -dimensional harmonic oscillator is thus  $u(\nu)$ . [J.M. Jauch and E.H. Hill, *On the Problem of Degeneracy in Quantum Mechanics*, Phys. Rev. 57, 641 (1940).] States are characterized by the totally symmetric irreducible representations  $[n, 0, \dots, 0] \equiv [n]$  of  $u(\nu)$ , with eigenvalues

$$E(n) = n + \frac{\nu}{2} \quad n = 0, 1, \dots, \infty. \quad (12.10)$$

Although harmonic oscillator problems are best attacked by bosonic realizations of Lie algebras, it is still of interest to consider differential realizations in terms of coordinates  $\mathbf{r} \equiv (x_1, x_2, \dots, x_\nu)$  and momenta  $\mathbf{p} = \frac{1}{i} \nabla \equiv \frac{1}{i} \left( \frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_\nu} \right)$ .

#### Example 1. Isotropic harmonic oscillator in three dimensions

As discussed in Chap. 7, this problem is best solved in spherical coordinates  $r, \vartheta, \varphi$ . The Hamiltonian (12.2) is

$$H = \frac{\mathbf{p}^2}{2} + \frac{\mathbf{r}^2}{2}, \quad (12.11)$$

where  $\mathbf{r}$  and  $\mathbf{p}$  are here three-dimensional vectors. The nine operators

$$\begin{aligned}
 H &= \frac{1}{2}(\mathbf{r}^2 + \mathbf{p}^2) \\
 \mathbf{L} &= -i\sqrt{2}[\mathbf{r} \times \mathbf{p}]^{(1)} \\
 \mathbf{Q} &= [\mathbf{r} \times \mathbf{r} + \mathbf{p} \times \mathbf{p}]^{(2)} \quad , \quad (12.12)
 \end{aligned}$$

where  $\mathbf{L}$  and  $\mathbf{Q}$  are the angular momentum vector and quadrupole (rank-2) tensor in three dimensions, satisfy commutation relations isomorphic to those of  $u(3)$ . (The notation used here is that of [A. deShalit and I. Talmi, *Nuclear Shell Theory*, Academic Press, 1963], where  $\mathbf{r}$  and  $\mathbf{p}$  are rank-1 tensors, and the superscript denotes tensor couplings. The operator  $\mathbf{Q}$  has an additional factor, often  $\sqrt{8}$ , in other definitions.) The algebra  $u(3)$  has three invariant Casimir operators,  $C_1, C_2$  and  $C_3$ . The Hamiltonian  $H$  of (12.11) can be rewritten as

$$H = C_1(u(3)) + \frac{3}{2}. \quad (12.13)$$


(Note that  $H$  commutes with the eight elements of the algebra  $su(3)$ ,  $\mathbf{L}$  and  $\mathbf{Q}$ .) The eigenstates are labelled by the irreducible representations  $[n, 0, 0] \equiv [n]$  of  $u(3)$ . A complete labelling is  $|n, l, m\rangle$  with

$$\left| \begin{array}{ccc}
 u(3) \supset so(3) \supset so(2) \\
 \downarrow \quad \downarrow \quad \downarrow \\
 n \quad \quad l \quad \quad m
 \end{array} \right\rangle . \quad (12.14)$$

The branching of the representations  $[n]$  of  $u(3)$  into representations of  $so(3) \supset so(2)$  has been discussed in Chap. 7. The values of  $l$  for each  $n$  are given by  $l = n, n - 2, \dots, 1$  or  $0$  ( $n = \text{odd or even}$ ). The values of  $m$  are  $m = -l, \dots, +l$ . The quantum number  $n = 0, \dots, \infty$ . The energy level diagram

$$E(n) = n + \frac{3}{2} \quad (12.15)$$

for this case is shown in Fig. 12.1. The three dimensional harmonic oscillator finds useful applications in a variety of problems in physics.



**Fig. 12.1.** The spectrum of the three-dimensional harmonic oscillator. States up to  $n = 4$  are shown

**Example 2.** *Isotropic harmonic oscillator in five dimensions*

This problem is of interest in nuclear physics, in the study of quadrupole oscillations of a liquid drop. The problem is best solved by introducing coordinates  $\alpha_\mu$  ( $\mu = 0, \pm 1, \pm 2$ ) and momenta  $\pi_\mu$  ( $\mu = 0, \pm 1, \pm 2$ ), called Bohr variables. The coordinates  $\alpha_\mu$  are the five components of a quadrupole (rank-2) tensor with respect to rotations. The Hamiltonian in dimensionless units is

$$H = \frac{1}{2} \left( \sum_{\mu} \pi_{\mu}^2 + \sum_{\mu} \alpha_{\mu}^2 \right). \tag{12.16}$$

This Hamiltonian can be rewritten as

$$H = C_1(u(5)) + \frac{5}{2}. \tag{12.17}$$


The basis states are labelled by the totally symmetric representations  $[n, 0, 0, 0, 0] \equiv [n]$  of  $u(5)$ . The complete labelling is  $|n, \tau, \nu_{\Delta}, l, m\rangle$  with

$$\left| \begin{array}{cccc} u(5) \supset so(5) \supset so(3) \supset so(2) \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ n \quad \tau, \nu_{\Delta} \quad l \quad m \end{array} \right\rangle. \tag{12.18}$$

The branching of the representations  $[n]$  of  $u(5)$  is given in Chap. 7, Sect. 6. The energy level diagram for this case

$$E(n) = n + \frac{5}{2} \tag{12.19}$$

is given in Fig. 12.2.



**Fig. 12.2.** The spectrum of the five-dimensional harmonic oscillator. States up to  $n = 4$  are shown

### 12.2.2 The Coulomb Problem

The Hamiltonian operator for this problem, when  $\hbar = m = 1$  and  $k = 1$ , is

$$H = \frac{p^2}{2} - \frac{1}{r} = -\frac{\nabla^2}{2} - \frac{1}{r}. \tag{12.20}$$

While the derivation of (12.13) is straightforward, the rewriting of  $H$  in terms of Casimir operators is more involved (see Examples 3 and 4). For bound states, where  $E = \langle H \rangle < 0$ , the Hamiltonian  $H$  can be rewritten as

$$H = -\frac{1}{2 \left[ C_2(so(\nu + 1)) + \left(\frac{\nu-1}{2}\right)^2 \right]}. \tag{12.21}$$

The degeneracy algebra of the  $\nu$ -dimensional Coulomb problem is thus  $so(\nu + 1)$ . (The Coulomb problem is a case in which the Hamiltonian  $H$  is not linear in the Casimir operators, but rather its inverse,  $1/H$ , is.) The eigenstates are characterized by the totally symmetric irreducible representations  $[\omega, 0, 0, \dots, 0] \equiv [\omega]$  of  $so(\nu + 1)$  with eigenvalues

$$E(\omega) = -\frac{1}{2 \left[ \omega(\omega + \nu - 1) + \left(\frac{\nu-1}{2}\right)^2 \right]} \quad \omega = 0, 1, \dots, \infty. \tag{12.22}$$

**Example 3.** *Coulomb problem in three dimensions*

The Coulomb problem in three dimensions (also called the Kepler problem) was the first problem for which degeneracy algebras were introduced, initially by Pauli [W. Pauli, *Über das Wasserstoffspektrum von Standpunkt der neuen Quantenmechanik*, Z. Phys. 36, 336 (1926)] following the classical treatment of Runge and Lenz, and subsequently by Fock [V.A. Fock, *Zur Theorie des Wasserstoffatoms*, Z. Phys. 98, 145 (1935)] and Bargmann [V. Bargmann, *Zur Theorie des Wasserstoffatoms*, Z. Phys. 99, 576 (1936)].

The problem is best solved in the familiar spherical coordinates,  $r, \vartheta, \varphi$ . In order to derive (12.21), one begins by introducing the angular momentum operator  $\mathbf{L}$  and the so-called normalized Runge-Lenz vector  $\mathbf{A}$

$$\begin{aligned} \mathbf{L} &= \mathbf{r} \times \mathbf{p} \\ \mathbf{A} &= \frac{1}{\sqrt{-2H}} \left[ \frac{1}{2} (\mathbf{p} \times \mathbf{L} - \mathbf{L} \times \mathbf{p}) - \frac{\mathbf{r}}{r} \right]. \end{aligned} \tag{12.23}$$

Using the commutation relations of  $\mathbf{r}$  and  $\mathbf{p} = \frac{1}{i}\nabla$ , one can show that the six components  $L_i$  ( $i = 1, 2, 3$ ) and  $A_i$  ( $i = 1, 2, 3$ ) satisfy, when  $E < 0$ , the commutation relations of the Lie algebra  $so(4)$

$$\begin{aligned} [L_i, L_j] &= i\varepsilon_{ijk}L_k \\ [L_i, A_j] &= i\varepsilon_{ijk}A_k \\ [A_i, A_j] &= i\varepsilon_{ijk}L_k, \end{aligned} \tag{12.24}$$

while when  $E > 0$  they satisfy the commutation relations of the non-compact algebra  $so(3, 1)$

$$\begin{aligned} [L_i, L_i] &= i\varepsilon_{ijk}L_k \\ [L_i, A_k] &= i\varepsilon_{ijk}A_k \\ [A_i, A_j] &= -i\varepsilon_{ijk}L_k . \end{aligned} \tag{12.25}$$

The components of the angular momentum and Runge-Lenz vector are thus, when  $E < 0$ , elements of the Lie algebra  $so(4)$ . This algebra has two quadratic Casimir operators,  $C_2$  and  $C'_2$  that can be written as

$$\begin{aligned} C_2(so(4)) &= (\mathbf{L}^2 + \mathbf{A}^2) \\ C'_2(so(4)) &= \mathbf{L} \cdot \mathbf{A} . \end{aligned} \tag{12.26}$$

After lengthy manipulations, the Hamiltonian operator can be written as

$$H = -\frac{1}{2(C'_2(so(4)) + 1)} . \tag{12.27}$$

As one can see from the definition (12.23) of the Runge-Lenz vector, the scalar product of  $\mathbf{L}$  and  $\mathbf{A}$  vanishes,  $\mathbf{L} \cdot \mathbf{A} = 0$ . The second invariant operator does not appear therefore in  $H$ . The basis states of  $so(4)$  are labelled by two quantum numbers  $[\omega_1, \omega_2]$ . However, due to the vanishing of  $C'_2(so(4))$ , only the totally symmetric representations  $[\omega] \equiv [\omega, 0]$  appear. The energy eigenvalues can be obtained from the eigenvalues of the Casimir operators in Chap. 5. They are given by


$$E(\omega) = -\frac{1}{2(\omega(\omega + 2) + 1)} \quad \omega = 0, 1, \dots, \infty . \tag{12.28}$$

In order to label completely the states one needs to consider the branching  $so(4) \supset so(3) \supset so(2)$ . This branching was considered in Chap. 7. The complete labelling of states is  $|\omega, l, m\rangle$ , with branching  $l = \omega, \omega - 1, \dots, 0$  and  $m = -l, \dots, +l$ . It is customary to introduce the “principal quantum number”  $n = \omega + 1$ . In terms of this quantum number the energy levels are

$$E(n, l, m) = -\frac{1}{2n^2} . \tag{12.29}$$

This is the celebrated Bohr formula that gives the energy levels of the non-relativistic hydrogen atom. The corresponding energy level diagram is shown in Fig. 12.3. The states  $|n, l, m\rangle$  are degenerate with total degeneracy  $n^2$ . Therefore  $so(4)$  is the degeneracy algebra of the Coulomb problem in  $\nu = 3$  dimensions. A detailed account is given in [B.G. Wybourne, *Classical Groups for Physicists*, J. Wiley and Sons, New York (1974), Chap. 21.]

**Example 4.** *Coulomb problem in six dimensions*



**Fig. 12.3.** The spectrum of the three-dimensional Coulomb problem. States up to  $\omega = 3$  are shown. The principal quantum number is  $n = \omega + 1$

This problem and its application to the three-body problem is best solved in hyperspherical Jacobi coordinates. One first introduces two vectors  $\boldsymbol{\rho}$  and  $\boldsymbol{\lambda}$  or equivalently  $\rho, \vartheta_\rho, \varphi_\rho$  and  $\lambda, \vartheta_\lambda, \varphi_\lambda$ . The Jacobi hyperspherical coordinates are  $r, \xi, \vartheta_\rho, \varphi_\rho, \vartheta_\lambda, \varphi_\lambda$  with

$$r = (\rho^2 + \lambda^2)^{1/2} \quad \xi = \arctan(\rho/\lambda) . \tag{12.30}$$

Introducing the notation

$$q_i \equiv (\boldsymbol{\rho}, \boldsymbol{\lambda}) \quad p_i \equiv (\mathbf{p}_\rho, \mathbf{p}_\lambda) \quad i = 1, \dots, 6 , \tag{12.31}$$

one can construct the 15 elements of the Lie algebra  $so(6)$  by

$$L_{ij} = q_i p_j - q_j p_i, \quad i, j = 1, \dots, 6 \quad i < j . \tag{12.32}$$

To these, one can add the six components of the normalized Runge-Lenz vector

$$A_i = \frac{1}{\sqrt{-2H}} \left[ \frac{1}{2} (L_{ij} p_j - p_j L_{ji}) - \frac{q_i}{r} \right] \quad i = 1, \dots, 6 . \tag{12.33}$$

The 21 elements  $L_{ij}, A_i$  satisfy the commutation relations

$$\begin{aligned} [L_{ij}, L_{kl}] &= i (\delta_{ik} L_{jl} + \delta_{jl} L_{ik} - \delta_{il} L_{jk} - \delta_{jk} L_{il}) \\ [A_i, A_j] &= i L_{ij} \\ [L_{ij}, A_k] &= i (\delta_{ik} A_j - \delta_{jk} A_i) . \end{aligned} \tag{12.34}$$

These commutation relations are isomorphic to those of  $so(7)$ . The 21 elements also commute with the Hamiltonian  $H$

$$[L_{ij}, H] = [A_i, H] = 0 \tag{12.35}$$

where

$$H = \frac{1}{2} (\mathbf{p}_\rho^2 + \mathbf{p}_\lambda^2) - \frac{1}{r}. \tag{12.36}$$

The algebra  $so(7)$  possesses three Casimir operators  $C_2, C_4, C_6$ . The quadratic operator is

$$C_2(so(7)) = \sum_{i < j}^6 L_{ij}^2 + \sum_i^6 A_i^2. \tag{12.37}$$

After some lengthy manipulations the Hamiltonian can be written as

$$H = -\frac{1}{2 [C_2(so(7)) + \frac{25}{4}]} . \tag{12.38}$$


The expectation value of this Hamiltonian in the representation  $[\omega] \equiv [\omega, 0, 0]$  is

$$E(\omega) = -\frac{1}{2 [\omega(\omega + 5) + \frac{25}{4}]}, \quad \omega = 0, 1, \dots, \infty. \tag{12.39}$$

The representations of  $so(7)$  that appear are the totally symmetric representations  $[\omega]$ , in view of conditions analogous to  $\mathbf{L} \cdot \mathbf{A} = 0$  in the three dimensional case. In order to label completely the states, one needs to study the branching of representations of  $so(7)$ . This can be done using the techniques of Chap. 4. The labelling of states appropriate for the choice of coordinates in this example is  $so(7) \supset so(6) \supset so_\rho(3) \oplus so_\lambda(3) \supset so(3) \supset so(2)$  with quantum numbers

$$\left. \begin{array}{cccccc} so(7) \supset so(6) \supset so(3) \oplus so(3) \supset so(3) \supset so(2) \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ \omega \quad \quad \gamma \quad \quad l_\rho, l_\lambda \quad \quad L \quad \quad M \end{array} \right\}. \tag{12.40}$$

Here  $\gamma = \omega, \omega - 1, \dots, 1, 0$ , and the values of  $l_\rho, l_\lambda$  are obtained by partitioning  $\gamma$  as  $\gamma = 2n_\gamma + l_\rho + l_\lambda, n_\gamma = 0, 1, \dots$ . The values of  $L$  are obtained from  $|l_\rho + l_\lambda| \geq L \geq |l_\rho - l_\lambda|$  and  $M = -L, \dots, +L$  as usual. The energy level diagram of the six dimensional Coulomb problem is shown in Fig. 12.4. The



**Fig. 12.4.** The spectrum of the six-dimensional Coulomb problem. States up to  $\omega = 2$  are shown

spectrum is degenerate and  $so(7)$  is the degeneracy algebra. The Coulomb problem in six dimensions is of interest in the three-body problem, in particular the three quark system in hadronic physics. [E. Santopinto, M. Giannini, and F. Iachello, *Algebraic Approach to the HyperCoulomb Problem*, in *Symmetry in Science VII*, Edited by B. Gruber, Plenum Press, New York, 1995.]

### 12.3 Degeneracy Algebra in $\nu = 1$ Dimension

The case  $\nu = 1$  is a special case, since the degeneracy algebra of the one-dimensional Hamiltonian, in units  $\hbar = m = 1$ ,

$$H = -\frac{1}{2} \frac{d^2}{dx^2} + V(x) \tag{12.41}$$

is always the trivial algebra  $u(1) \sim so(2)$ . Because in one dimension there is no rotational invariance to impose, the class of exactly solvable problems in  $\nu = 1$  dimension, that is of problems that can be written in terms of Casimir operators of an algebra  $g$ , is much wider than that in  $\nu \geq 2$  dimensions. A partial list is given in Table 12.1. The Hamiltonian for these problems is either linear, case IV, or quadratic, cases I–III, in the Casimir operators and hence the eigenvalues,  $E(n)$ , are either linear or quadratic in the quantum number  $n$ . The strength of the potential is given in the table either by  $\kappa(\kappa - 1)$  or by  $\kappa^2$ .

**Example 5.** *The one dimensional harmonic oscillator*

The one dimensional harmonic oscillator,  $V(x) = \frac{1}{2}kx^2$ , in units  $k = 1$ , has been treated in Chap. 7. Its Hamiltonian can be written as

$$H = C_1(u(1)) + \frac{1}{2} \tag{12.42}$$

with eigenvalues

$$E(n) = n + \frac{1}{2} \quad n = 0, 1, \dots, \infty . \tag{12.43}$$

**Table 12.1.** Exactly solvable quantum mechanical problems in one dimension

Case	$V(x)$	$E(n)$
(I)	$\frac{1}{2} \frac{\kappa(\kappa - 1)}{\sin^2 x}$	$\frac{1}{2} (\kappa + 2n)^2$
(II)	$-\frac{1}{2} \frac{\kappa(\kappa - 1)}{\cosh^2 x}$	$-\frac{1}{2} (\kappa - 1 - n)^2$
(III)	$\kappa^2 (e^{-2x} - 2e^{-x})$	$-\kappa^2 + \frac{1}{2} \left[ 2\sqrt{2}\kappa \left( n + \frac{1}{2} \right) - \left( n + \frac{1}{2} \right)^2 \right]$
(IV)	$\frac{1}{2} \kappa^2 x^2$	$\kappa \left( n + \frac{1}{2} \right)$



## 12.4 Dynamical Algebras

The degeneracy algebra  $g_d$  allows one to solve the eigenvalue problem and thus classify the degenerate multiplets. However, in the cases discussed in the previous Chap. 11 all states of the system were assigned to an irreducible representation of an algebra  $g$ , and all operators  $H, T, \dots$  were written in terms of elements of  $g$ . It is of interest to do the same for quantum mechanical problems. This problem is often called the *embedding* problem. The algebra  $g$  that contains the degeneracy algebra  $g \supset g_d$  is called the dynamical algebra. Included in the dynamical algebra  $g$  there are now raising and lowering operators that relate the different representations of  $g$ . The embedding problem does not have a unique solution. An additional complication is that often the quantum mechanical problem has an infinite number of bound ( $E < 0$ ) eigenstates, as in the two cases of Chap. 2. The dynamical algebra must therefore be either a non-compact algebra or obtained by a limiting process (called contraction) from a compact algebra. Both of these problems are outside the scope of this book and will therefore be only mentioned here.

## 12.5 Dynamical Algebras in $\nu \geq 2$ Dimensions

### 12.5.1 Harmonic Oscillator

The degeneracy algebra is  $u(\nu)$ . A commonly used dynamical algebra is the symplectic algebra  $sp(2\nu, R) \supset u(\nu)$ . [R.C. Hwa and T. Nuyts, *Group Embedding for the Harmonic Oscillator*, Phys. Rev. 145, 1188 (1966).] However, this embedding has the disadvantage that two irreducible representations are needed to accommodate all the states of the harmonic oscillator. A simpler embedding is obtained by drawing on the results of Chap. 7. Introducing a fictitious coordinate,  $s$ , and momentum,  $p_s = \frac{1}{i} \frac{d}{ds}$ , one can construct the algebra  $u(\nu + 1)$ . The algebra  $u(\nu + 1) \supset u(\nu)$  can be used as a dynamical algebra of the harmonic oscillator. All states are assigned to the representation  $[N]$  of  $u(\nu + 1)$  with  $N \rightarrow \infty$ . The degenerate multiplets are labelled by  $n = 0, 1, \dots, N \rightarrow \infty$ .

### 12.5.2 Coulomb Problem

The degeneracy algebra is  $so(\nu + 1)$ . The non-compact algebras  $so(\nu + 1, 2) \supset so(\nu + 1, 1) \supset so(\nu + 1)$  have been used as dynamical algebras [H. Bacry, *The de Sitter Group  $L_{4,1}$  and the Bound States of the Hydrogen Atom*, Nuovo Cimento 41A, 222 (1966)], [E.C.G. Sudarshan, N. Mukunda, and L. O’Raifeartaigh, *Group Theory of the Kepler problem*, Phys. Lett. 19, 322 (1965)], [A.O. Barut, *Dynamical Groups and their Currents*, Springer Tracts Mod. Phys. 50, 1 (1969).] However, the representation theory of non-compact algebras is rather complicated and requires special attention. Again, a simpler

embedding is obtained by introducing a fictitious coordinate  $s$  and momentum  $\frac{1}{i} \frac{d}{ds}$ , and constructing the algebra  $so(\nu + 2)$ . This algebra, instead of  $so(\nu + 1, 1)$ , can be used as dynamical algebra of the Coulomb problem in  $\nu \geq 2$  dimensions. All states are assigned to a representation  $[I]$  of  $so(\nu + 2)$  with  $I \rightarrow \infty$ . The degenerate multiplets are labelled by  $\omega = 0, 1, \dots, I \rightarrow \infty$ .

## 12.6 Dynamical Algebra in $\nu = 1$ Dimension

The algebra  $u(2)$  has been used extensively as a dynamical algebra in  $\nu = 1$  dimension. In particular, this algebra has been used in the study of two quantum mechanical problems of practical interest, already listed in Chap. 3,  $V(x) = -\frac{k}{\cosh^2 x}$  (called the Pöschl-Teller potential) and  $V(x) = k(e^{-2x} - 2e^{-x})$  (called the Morse potential).

### 12.6.1 Pöschl-Teller Potential

The Hamiltonian operator for this problem in units  $\hbar = m = 1$  is

$$H = -\frac{1}{2} \frac{d^2}{dx^2} - \frac{k}{\cosh^2 x}, \quad (12.44)$$

where  $x$  is a dimensionless coordinate. In order to construct the dynamical algebra  $u(2) \supset su(2)$ , introduce two variables  $\vartheta$  and  $\varphi$  and consider the differential realization of  $su(2) \sim so(3)$  on the sphere, given in Chap. 9,

$$\begin{aligned} I_z &= -i \frac{\partial}{\partial \varphi} \\ I_{\pm} &= -ie^{\mp i\varphi} \left( \frac{\partial}{\partial \vartheta} \mp i \cot \vartheta \frac{\partial}{\partial \varphi} \right) \end{aligned} \quad (12.45)$$

and

$$I^2 = - \left[ \frac{1}{\sin \vartheta} \frac{\partial}{\partial \vartheta} \left( \sin \vartheta \frac{\partial}{\partial \vartheta} \right) + \frac{1}{\sin^2 \vartheta} \frac{\partial^2}{\partial \varphi^2} \right]. \quad (12.46)$$

The simultaneous eigenfunctions of  $I^2$  and  $I_z$

$$\begin{aligned} I^2 \chi_j^m &= j(j+1) \chi_j^m \\ I_z \chi_j^m &= m \chi_j^m \end{aligned} \quad (12.47)$$

with  $j$  and  $m$  integer, are

$$\chi_j^m(\vartheta, \varphi) = u_j^m(\vartheta) e^{im\varphi} \quad (12.48)$$

where  $u_j^m(\vartheta)$  satisfies the equation

$$\left[ -\frac{1}{\sin \vartheta} \frac{\partial}{\partial \vartheta} \left( \sin \vartheta \frac{\partial}{\partial \vartheta} \right) + \frac{m^2}{\sin^2 \vartheta} \right] u_j^m(\vartheta) = j(j+1) u_j^m(\vartheta). \quad (12.49)$$

The solutions of (12.50) are the associated Legendre functions  $P_j^m(\cos \vartheta)$  and the functions  $\chi_j^m(\vartheta, \varphi)$  are just the spherical harmonics  $Y_{jm}(\vartheta, \varphi)$ . The substitution

$$\cos \vartheta = \tanh x \quad -\infty < x < +\infty \quad (12.50)$$

brings (12.49) to the form

$$\left[ -\frac{d^2}{dx^2} - \frac{j(j+1)}{\cosh^2 x} \right] u_j^m(x) = -m^2 u_j^m(x). \quad (12.51)$$

This is, apart from a factor 2, the Schrödinger equation with Pöschl-Teller potential. The strength of the potential  $k$  is related to the eigenvalue of the Casimir operator of  $su(2) \sim so(3)$ ,  $j(j+1)$ , by  $k = \frac{1}{2}j(j+1)$ . The Hamiltonian operator can be written as

$$H = -\frac{1}{2}I_z^2 \quad (12.52)$$


with eigenvalues

$$E(m) = -\frac{1}{2}m^2. \quad (12.53)$$

In the bra-ket notation, the eigenfunctions can be written as

$$\left| \begin{array}{cc} u(2) \supset so(2) \\ \downarrow \quad \quad \downarrow \\ j \quad \quad m \end{array} \right\rangle. \quad (12.54)$$

All eigenstates are assigned to the representation  $j$  of  $u(2)$ . This algebra is the dynamical algebra of the problem. The Hamiltonian is written in terms of elements of this algebra, in fact in terms of the Casimir operator  $I_z$  of the subalgebra  $so(2)$ , which is the degeneracy algebra of the problem. There is a peculiarity in this case due to the fact that the algebra  $so(2)$  is an orthogonal algebra in an even number of dimensions and thus  $m = -j, \dots, +j$ . Since the eigenvalues depend only on  $m^2$ , they are double degenerate. They correspond to the eigenvalues of the potential and its reflection. The algebra  $u(2)$  should therefore be used as dynamical algebra with the proviso that only states with  $m \geq 0$  should be considered. The spectrum of the Pöschl-Teller potential with this proviso is shown in Fig. 12.5. The Pöschl-Teller potential is of interest in molecular physics. [F. Iachello and S. Oss, *Algebraic Model of Bending Vibrations of Complex Molecules*, Chem. Phys. Lett. 205, 285 (1993).] In applications in molecular physics, often the quantum number  $m = j, j-1, \dots, 1, 0$  is replaced by the so-called vibrational quantum number  $n = j-m, n = 0, 1, \dots, j$ . In contrast with the harmonic oscillator of Example 6, the Pöschl-Teller potential has a finite number of bound states. Thus  $j = \text{finite}$ , and the representations of  $u(2)$  that appear in this problem are the usual finite dimensional representations.



**Fig. 12.5.** The spectrum of the one-dimensional Pöschl-Teller potential superimposed to the potential. The strength of the potential is characterized by  $j = 4$

### 12.6.2 Morse Potential

The Hamiltonian for this problem in units  $\hbar = m = 1$  is

$$H = -\frac{1}{2} \frac{d^2}{dx^2} + k [e^{-2x} - 2e^{-x}] . \tag{12.55}$$

In order to construct the spectrum generating algebra, introduce again two variables  $s$  and  $t$ . A realization of  $u(2)$  in terms of differential operators  $s \frac{\partial}{\partial s}, s \frac{\partial}{\partial t}, t \frac{\partial}{\partial s}, t \frac{\partial}{\partial t}$  is

$$\begin{aligned} \hat{F}_x &= \frac{1}{2} \left( st - \frac{\partial^2}{\partial s \partial t} \right) \\ \hat{F}_y &= \frac{1}{2i} \left( s \frac{\partial}{\partial t} - t \frac{\partial}{\partial s} \right) \\ \hat{F}_z &= \frac{1}{4} \left( s^2 - t^2 - \frac{\partial^2}{\partial s^2} + \frac{\partial^2}{\partial t^2} \right) \\ \hat{N} &= \frac{1}{2} \left( s^2 + t^2 - \frac{\partial^2}{\partial s^2} - \frac{\partial^2}{\partial t^2} - 2 \right) . \end{aligned} \tag{12.56}$$

This realization is the oscillator realization of Sect. 7.3 written in terms of differential operators. A hat is put in (12.56) to distinguish an operator from its eigenvalue. Consider now the simultaneous eigenstates of  $\hat{N}$  and  $\hat{F}_y$ . Introducing polar coordinates

$$\begin{aligned} s &= r \cos \varphi & 0 \leq r < \infty \\ t &= r \sin \varphi & 0 \leq \varphi < 2\pi , \end{aligned} \tag{12.57}$$

the two operators can be rewritten as

$$\begin{aligned}\hat{F}_y &= -\frac{i}{2} \frac{\partial}{\partial \varphi} \\ \hat{N} &= \frac{1}{2} \left( r^2 - \frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial}{\partial r} - \frac{1}{r^2} \frac{\partial^2}{\partial \varphi^2} \right) - 1.\end{aligned}\quad (12.58)$$

The simultaneous eigenstates of  $\hat{N}$  and  $\hat{F}_y$  can be written as

$$\psi_{N,m_y}(r, \varphi) = R_{N,m_y}(r) e^{2im_y \varphi} \quad (12.59)$$

where  $2m_y = \text{integer}$  since  $\psi$  should be periodic in  $\varphi$  with period  $2\pi$ , and  $R_{N,m_y}(r)$  satisfies

$$\frac{1}{2} \left( -\frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial}{\partial r} + \frac{4m_y^2}{r^2} + r^2 \right) R_{N,m_y}(r) = (N+1) R_{N,m_y}(r). \quad (12.60)$$

By the change of variable

$$r = \sqrt{N+1} e^{-x/2} \quad (12.61)$$

this equation can be brought to the form

$$\left[ -\frac{d^2}{dx^2} + \left( \frac{N+1}{2} \right)^2 (e^{-2x} - 2e^{-x}) \right] R_{N,m_y}(x) = -m_y^2 R_{N,m_y}(x). \quad (12.62)$$

This is, apart from a factor 2, the Schrödinger equation with Morse potential. The strength of the potential  $k$  is related to the eigenvalue of the Casimir operator of  $su(2)$  by

$$k = \frac{1}{2} \left( \frac{N+1}{2} \right)^2. \quad (12.63)$$

The Hamiltonian operator can be written as

$$H = -\frac{1}{2} \hat{F}_y^2 \quad (12.64)$$


with eigenvalues

$$E(m_y) = -\frac{1}{2} m_y^2. \quad (12.65)$$

In bra-ket notation, the eigenfunctions can be written as

$$\left| \begin{array}{cc} u(2) \supset u(1) \\ \downarrow \quad \downarrow \\ F \quad m_y \end{array} \right\rangle, \quad (12.66)$$

where, as discussed in Chap. 7, Sect. 3,  $F = \frac{N}{2}$ . The values of  $m_y$  are  $m_y = -\frac{N}{2}, \dots, +\frac{N}{2} = -F, \dots, +F$ , but, for reasons given above, only  $m_y \geq 0$  need be considered. The spectrum of the Morse potential is shown in Fig. 12.6.



**Fig. 12.6.** The spectrum of the one-dimensional Morse potential. The strength of the potential is characterized by  $F = \frac{N}{2} = 4$

The Morse potential is of great interest in molecular physics. [Y. Alhassid, F. Iachello and F. Gürsey, *Group Theory of the Morse Oscillator*, Chem. Phys. Lett. 99, 27 (1983).] Here also it is customary to introduce a vibrational quantum number  $v = \frac{N}{2} - m_y, v = 0, 1, \dots, \frac{N}{2}$  or  $\frac{N-1}{2}$  ( $N = \text{even or odd}$ ), and the number of bound states is finite. The Morse potential problem in one dimension is soluble not only for the values such that  $N$  integer, but for any strength  $k$ . The general solution requires however the use of *projective* representations of  $u(2)$  rather than tensor representations. Projective representations of  $u(2)$  [V. Bargmann, *Irreducible Unitary Representations of the Lorentz Group*, Ann. Math. 48, 568 (1947)] are outside the scope of these lectures note and will not be discussed.

The Morse potential has also been associated with representations of the non-compact algebra  $su(1,1)$  [P. Cordero and S. Hojman, *Algebraic Solution of a Short-Range Potential Problem*, Lett. Nuovo Cimento 4, 1123 (1970)].

### 12.6.3 Lattice of Algebras

The concept of lattice of algebras introduced in Chap. 7, can be used here as well. One dimensional exactly solvable problems are characterized by the lattice of algebras

$$\begin{array}{c} u(2) \\ | \\ u(1) \sim so(2) \end{array} \tag{12.67}$$

Here  $u(2)$  is the dynamical algebra and  $u(1) \sim so(2)$  is the degeneracy algebra. Because of the isomorphism  $u(1) \sim so(2)$ , all exactly solvable problems in  $\nu = 1$  dimension have the same structure. In particular, the two problems discussed in Sects. 6.1 and 6.2 have the same bound state spectrum

$$E(m) = -\frac{1}{2}m^2 . \tag{12.68}$$

Problems with the same bound ( $E < 0$ ) spectrum are called *isospectral*. (They differ, however, in the scattering ( $E > 0$ ) spectrum, not discussed here.)

The algebra  $u(2)$  can also be used as a dynamical algebra of the harmonic oscillator of Chap. 3. The Hamiltonian is now linear in the Casimir operator  $C_1$  of  $u(1)$  with eigenvalue

$$E(n) = n + \frac{1}{2}. \quad (12.69)$$

The harmonic oscillator has an infinite number of bound states and thus  $N \rightarrow \infty$  and  $n = 0, 1, \dots, \infty$ .

## References

- Alhassid, Y., Iachello, F., and Gürsey, F., *Group Theory of the Morse Oscillator*, Chem. Phys. Lett. 99, 27 (1983).
- Bacry, H., *The de Sitter Group  $L_{4,1}$  and the Bound States of the Hydrogen Atom*, Nuovo Cimento 41A, 222 (1966).
- Baird, G.E., and Biedenharn, L.C., *On the Representations of Semisimple Lie Groups-II*, J. Math. Phys. 4, 1449 (1963).
- Bargmann, V., *Zur Theorie des Wasserstoffatoms*, Z. Phys. 99, 576 (1936).
- Bargmann, V., *Irreducible Unitary Representations of the Lorentz Group*, Ann. Math. 48, 568 (1947).
- Bröcker, T., and tom Dieck, T., *Representations of Compact Lie Groups*, Springer-Verlag, New York, 1985.
- Barut, A.O., *Dynamical Groups and their Currents*, Springer Tracts in Mod. Phys. 50,1 (1969).
- Barut, A.O., and Raçzka, R., *Theory of Group Representations and Applications*, World Scientific, Singapore, 1986.
- Barut, A.O., and Böhm, A., *Dynamical Groups and Mass Formula*, Phys. Rev. B139, 1107 (1965).
- Berezin, F.A., *An Introduction to Superanalysis*, D. Riedel, Dordrecht, 1987.
- Bijker, R., Iachello, F., and Leviatan, A., *Algebraic Models of Hadronic Structure. I. Non Strange Baryons*, Ann. Phys. (NY) 236, 69 (1994).
- Bijker, R., and Leviatan, A., *Algebraic Treatment of Collective Excitations in Baryon Spectroscopy*, in *Symmetries in Science VII: Spectrum Generating Algebras and Dynamic Symmetries in Physics*, Eds. B. Gruber and T. Otsuka, Plenum Press, New York, 1994.
- Cartan, E., *The Theory of Spinors*, Hermann, Paris, 1966.
- Cartan, E., *Sur la Structure des Groupes de Transformation Finis et Continus*, Thèse, Paris, 1894.
- Casimir, H., *Über die Konstruktion einer zu den irreduzibelen Darstellung halbeinfacher kontinuierlichen Gruppen gehörigen Differential-gleichung*, Proc. Roy. Akad. Amst. 34, 844 (1931).
- Chaichian, M., and Hagedorn, R., *Symmetry in Quantum Mechanics: from Angular Momentum to Supersymmetry*, Institute of Phys. Publ., Philadelphia, PA, 1998.



- Chen, J.Q., *Group Representation Theory for Physicists*, World Scientific, Singapore, 1989.
- Chen, J.-Q., Chen, B.-Q., and Klein, A., *Factorization of Commutators: The Wick Theorem for Coupled Operators*, Nucl. Phys. A554, 61 (1993).
- Cordero, P., and Hojman, S., *Algebraic Solution of a Short-Range Potential Problem*, Lett. Nuovo Cimento 4,1123 (1970).
- De Shalit, A. and Talmi, I., *Nuclear Shell Theory*, Academic Press, New York, 1963.
- de Swart, J.J., *The Octet Model and its Clebsch-Gordan coefficients*, Rev. Mod. Phys. 35, 916 (1963).
- Dothan, Y., Gell-Mann, M., and Ne'eman, Y., *Series of Hadron Levels as Representations of Non-compact Groups*, Phys. Lett. 17, 148 (1965).
- Dynkin, E.B., *The Structure of Semisimple Lie Algebras*, Usp. Mat. Nauk (N.S.), 2, 59 (1947). Translated in Am. Math. Soc. Transl. (1), 9, 308 (1962).
- Elliott, J.P., *Collective Motion in the Nuclear Shell Model. I. Classification Scheme for States of Mixed Configurations*, Proc. Roy. Soc. A 245, 128 (1958).
- Flowers, R.H., *Studies in  $j$ - $j$  Coupling. I. Classification of Nuclear and Atomic States*, Proc. Roy. Soc. A212, 248 (1952).
- Fock, V.A., *Zur theorie des Wasserstoffatoms*, Z. Phys. 98, 145 (1935).
- Frank, A., and van Isacker, P., *Algebraic Methods in Molecular and Nuclear Structure Physics*, J. Wiley and Sons, New York, 1994.
- Franzini, P., and Radicati, L.A., *On the Validity of the Supermultiplet Model*, Phys. Lett. 6, 32 (1963).
- French, J.B., *Multipole and Sum-rule Methods in Spectroscopy*, in *Proceedings of the International School of Physics "Enrico Fermi"*, Course XXXVI, Ed. C. Bloch, Academic Press, New York, 1966, p. 278.
- Fulton, W., and Harris, J., *Representation Theory: A First Course*, Springer, New York, 1991.
- Gel'fand, I.M., and Cetlin, M.L., *Finite-Dimensional Representations of a Group of Unimodular Matrices*, Dokl. Akad. Nauk SSSR 71, 8, 825 (1950); *Finite-Dimensional Representations of Groups of Orthogonal Matrices*, ibid. 71, 1017 (1950).
- Gell-Mann, M., *Symmetries of Baryons and Mesons*, Phys. Rev. 125, 1067 (1962).
- Georgi, H., *Lie Algebras in Particle Physics*, Perseus Books, Reading, MA., 1999.
- Gilmore, R., *Lie Groups, Lie Algebras and Some of Their Applications*, J. Wiley and Sons, New York, 1974.
- Gürsey, F., and Radicati, L.A., *Spin and Unitary Spin Independence of Strong Interactions*, Phys. Rev. Lett. 13, 173 (1964).
- Hamermesh, M., *Group Theory and its Applications to Physical Problems*, Addison-Wesley, Reading, Massachusetts, 1962.

- Hermann, R., *Lie Groups for Physicists*, W.A. Benjamin, New York, 1966.
- Hladik, J., *Spinors in Physics*, translated from the French edition *Spineuren Physique*, Springer, New York, 1999.
- Humphreys, J.E., *Introduction to Lie Algebras and Representation Theory*, Graduate Texts in Mathematics, Springer-Verlag, New York, 1972.
- Hwa, R.C., and Nuyts, T., *Group Embedding for the Harmonic Oscillator*, Phys. Rev. 145, 1188 (1966).
- Iachello, F. and Arima, A., *The Interacting Boson Model*, Cambridge University Press, Cambridge, England, 1987.
- Iachello, F. and Levine, R.D., *Algebraic Theory of Molecules*, Oxford University Press, Oxford, England, 1995.
- Iachello, F. and van Isacker, P., *The Interacting Boson Fermion Model*, Cambridge University Press, Cambridge, England, 1991.
- Iachello, F., *Algebraic Theory of the Three-Body Problem*, in *Symmetries in Science VIII*, Ed. B. Gruber, Plenum Press, New York, 1995.
- Iachello, F., and Oss, S., *Algebraic Approach to Molecular Spectra: Two-dimensional Problems*, J. Chem. Phys. 104, 6956 (1996).
- Iachello, F., and Oss, S., *Algebraic Model of Bending Vibrations of Complex Molecules*, Chem. Phys. Lett. 205, 285 (1993).
- Jauch, J.M., and Hill, E.H., *On the Problem of Degeneracy in Quantum Mechanics*, Phys. Rev. 57, 641 (1940).
- Kirillov, A.A., *Elements of the Theory of Representation*, Nauka, Moscow, 1978.
- Killing, W., *Die Zusammensetzung der Stetigen Endlichen Transformationsgruppen, I-IV*, Math. Ann. 31, 252 (1888); 33,1 (1889); 34,57 (1889); 36, 161 (1890).
- Lie, S. and Scheffers, G., *Vorlesungen über Kontinuerliche Gruppen*, Leipzig, 1893.
- Messiah, A., *Quantum Mechanics*, J. Wiley and Sons, New York, translated from the French edition, Dunod, Paris, 1958.
- Okubo, S., *Note on Unitary Symmetry in Strong Interactions*, Progr. Theor. Phys. 27, 949 (1962).
- Pauli, W., *Über das Wasserstoffspektrum von Standpunkt der neuen Quantenmechanik*, Z. Phys. 36, 336 (1926).
- Perelomov, A.M., and Popov, V.S., *Casimir Operators for  $u(n)$  and  $su(n)$* , Sov. J. of Nucl. Phys. 3, 676 (1966); *Casimir Operators of Orthogonal and Symplectic Groups*, *ibid.* 3, 819 (1966).
- Racah, G., *Group Theory and Spectroscopy*, Mimeographed Lecture Notes, Princeton, New Jersey, 1951, reprinted in Springer Tracts in Modern Physics 37, 28 (1965).
- Racah, G., *Theory of Complex Spectra-IV*, Phys. Rev. 76, 1352 (1949).
- Santopinto, E., Giannini, M., and Iachello, F., *Algebraic Approach to the HyperCoulomb Problem*, in *Symmetry in Science VII*, B. Gruber, ed., Plenum Press, New York, 1995.

- Schwinger, J., in *Quantum Theory of Angular Momentum*, L.C. Biedernharn and H. Van Dam, eds., Academic Press, New York, 1965, p. 229.
- Sattinger, D.H., and Weaver, O.L., *Lie Groups, Lie Algebras with Applications to Physics, Geometry and Mechanics*, Springer-Verlag, New York, 1986.
- van der Waerden, B.L., *Die Gruppentheoretische Methode in der Quantenmechanik*, Math. Zeitschr. 37, 446 (1933).
- Van Roosmalen, O.S., *Algebraic Description of Nuclear and Molecular Rotation-Vibration Spectra*, Ph.D. Thesis, University of Groningen, The Netherlands, 1982.
- Varadarajan, V.S., *Lie Groups, Lie Algebras, and Their Representations*, Springer-Verlag, New York, 1984.
- Weyl, H., *Theorie der Darstellung kontinuierlicher halbeinfacher Gruppen durch lineare Transformationen, I-IV*, Math. Zeitschr. 23, 271 (1925); 24, 328, 377 (1926).
- Wybourne, B.G., *Classical Groups for Physicists*, J. Wiley and Sons, New York, 1974.

# Index

- Ado's theorem 24
- angular momentum operator 148
- anticommutator 131
- associativity 27
  
- basis
  - Bose-Einstein 92
  - Fermi-Dirac 132
  - irreducible 39, 75
- boson calculus 103
- bosonic operator 91
- branching problem 93
  
- Campbell-Hausdorff formula 38
- Cartan criterion 8
- Cartan-Weyl form 15, 17
  - of  $so(3)$  18
  - of  $su(2)$  95
  - of  $su(3)$  18
- Casimir operator
  - complete set 65
  - definition 10, 63
  - eigenvalue of 66–74
  - Laplace-Beltrami form 150
  - matrix form 161
  - of exceptional algebra 65
  - of  $so(n)$  64
  - of  $sp(n)$  65
  - of  $su(n)$  64
  - of  $u(n)$  63
- change of basis 3
- Clebsch-Gordan coefficient *see*
  - coupling coefficient
- Clifford algebra 157, 161
- closure 27
- commutation relations
  - Bose 24, 91
  - Fermi 25, 131
- commutator 1
  - tensor coupled 98
- Coulomb problem 177
  - three-dimensional 177
  - six-dimensional 178
  - embedding 182
- coupling
  - $j$ - $j$  145
  - $L$ - $S$  142
- coupling coefficient
  - computation 85
  - definition 76
  - $so(3)$  77, 84
  - $so(4)$  80
  - $su(3)$  81
  - symmetry properties 84
- degeneracy algebra 173
  - in one dimension 181
- derivation 9
- direct sum 5
- dynamic symmetry 163
  - bosonic system 164
  - fermionic system 170
- dynamical algebra 182
- Dynkin diagram 20
  
- embedding problem 182
- energy formula 164
- energy level diagram
  - $u(4)$  166
  - $u(6)$  168
- enveloping algebra 23
- equivalence relations 42
- exactly solvable problem 173, 174, 181
- exponential map 37
  
- fermionic operator 131

- Gel'fand pattern 47  
 Gell-Mann matrix 158  
 Gürsey-Radicati  $u(6)$  141, 171  
 harmonic oscillator  
   one-dimensional 92, 188  
   two-dimensional 94  
   three-dimensional 101, 103, 174  
   five-dimensional 176  
   embedding 182  
 homomorphism 37  
 hyperspherical coordinates 149  
  
 Iachello-Arima  $u(6)$  169  
 Iachello-Levine  $u(4)$  167  
 ideal *see* subalgebra, invariant  
 identity 27  
 internal degrees of freedom 139, 142  
 invariant operator *see* Casimir  
   operator  
   for non-semisimple algebra 12  
 inverse 27  
 isomorphism 3, 21, 49  
 isoscalar factor 80  
 isospin 139  
  
 Jacobi identity 2  
 Jordan-Schwinger construction 96  
  
 Kepler problem *see* Coulomb problem  
 Killing form 8  
 Kirillov classification 12  
 Kronecker product 40  
  
 label 39  
   missing 61  
 lattice of algebras  
    $u(2)$  97, 187  
    $u(3)$  103  
    $u(4)$  109, 114  
    $u(6)$  123  
    $u(7)$  130  
 Legendre function 184  
 Lie algebra  
   Abelian 5  
   action of elements 53  
    $A_l$  *see* Lie algebra, special unitary  
   angular momentum *see* Lie algebra,  
     special orthogonal  
    $B_l$  *see* Lie algebra, special  
     orthogonal  
    $C_l$  *see* Lie algebra, symplectic  
   classification 21  
   compact 9  
   complex extension 4  
   definition 1  
    $D_l$  *see* Lie algebra, special  
     orthogonal  
    $e(2)$  7, 9, 10, 13  
    $E_6$  21  
    $E_7$  21  
    $E_8$  21  
   Euclidean 7  
   exceptional 21  
    $F_4$  21  
    $G_2$  16, 21, 124, 143  
   Lorentz 3, 152  
   nilpotent 10  
   orthogonal 21  
   Racah form 97, 132, 142  
   real form 21  
   semisimple 7, 8, 21  
    $so(2)$  4, 7, 12, 158  
    $so(2)$  148  
    $so(2, 1)$  3, 8, 9, 11, 13  
    $so(3)$  3, 7–9, 11, 13, 16, 18, 57, 99,  
     148, 158  
    $so(3, 1)$  6, 152, 178  
    $so(4)$  5, 16, 48, 149, 177  
    $so(5)$  16, 20  
    $so(6)$  58  
    $so(7)$  181  
   solvable 9  
    $sp(2)$  153, 159  
    $sp(4)$  16, 159  
   special orthogonal 21, 99, 148, 158  
   special unitary 21, 99, 134  
    $spin(3)$  134  
   spinor 44  
    $su(2)$  16, 55, 95, 134, 156, 160, 161  
    $su(2)$  148  
    $su(3)$  16, 18, 59, 99, 157, 161  
    $su_F(3)$  141, 171  
    $su(4)$  139  
    $su(6)$  141  
   symplectic 21, 134, 153, 159  
    $t(2)$  5, 7, 12

- $t(3)$  13
- tensor coupled form 98
- translation 5
- $u(1)$  12, 92
- $u(2)$  54, 93, 134, 156
- $u(2)$  148
- $u(3)$  59, 99, 157
- $u(4)$  108, 136, 139, 165, 170
- $u(5)$  116
- $u(6)$  115, 137, 141, 167
- $u(7)$  123, 143
- $u(12)$  146
- $u(14)$  143
- unitary 21, 99, 134, 147, 155
- Lie group
  - compact 33
  - definition 27
  - general linear 29
  - Lorentz 33
  - matrix 27
  - non-compact 33
  - orthogonal 30
  - rotation *see* Lie group, special
    - orthogonal
    - $SO(1, 1)$  33
    - $SO(2)$  32
    - $SO(3)$  34, 35
  - special linear 29
  - special orthogonal 31
  - special unitary 30
  - $SU(2)$  34, 35
  - symplectic 31
  - unitary 29
- mass formula 171
- matrix exponential 38
- matrix properties 28
- metric tensor 8
- Morse potential 185
- operator
  - lowering 18
  - raising 18
- Pauli matrix 157
- Pöschl-Teller potential 183
- quantum number 39
- Racah form *see* Lie algebra, Racah form
- Racah's factorization lemma 79
- rank 15
- realization
  - bosonic 24, 91
  - differential 23, 147
  - fermionic 25, 131
  - Grassman 24, 152
  - matrix 23, 155
- recoupling coefficient
  - computation 86
  - definition 83
  - double 87
  - $so(3)$  83
  - symmetry properties 86
- reduction formula 88, 89
- representation
  - decomposition of 59, 62
  - dimension of 51
  - fundamental 45
  - irreducible 39, 42
  - mixed symmetry 50
  - of  $so(n)$  43, 44
  - of  $sp(n)$  43
  - of  $su(n)$  42
  - of  $u(n)$  42
  - spinor 44
  - tensor 42
  - totally antisymmetric 50
  - totally symmetric 50
- root 15
  - long 20
  - short 20
- root diagram 16
- rotovibrator
  - three-dimensional 167
  - five-dimensional 169
- Runge-Lenz vector 177
- semidirect sum 7
- spectrum generating algebra 163
- spherical harmonic 151
- spinor 152
- structure constants 2
- subalgebra
  - Cartan 15
  - definition 4
  - invariant 7

- maximal Abelian *see* subalgebra, Cartan
- subalgebra chain 46
  - canonical 46
  - containing  $so(3)$  99
  - containing  $spin(3)$  134
  - non-canonical 58
- tensor
  - antisymmetric 40
  - irreducible 40
  - symmetric 40
- tensor calculus 106
- tensor operator
  - adjoint 81
  - coupling of 88
  - definition 75
  - double 142
  - $so(3)$  75
- tensor product 56
- vacuum 24
- vector
  - column 155
  - row 155
- vector space 2, 39
- Wigner  $u(4)$  139, 170
- Wigner-Eckart theorem 77
- Young tableau 40

# Lecture Notes in Physics

For information about earlier volumes  
please contact your bookseller or Springer  
LNP Online archive: [springerlink.com](http://springerlink.com)

- Vol.663: A. Kalloniatis, D. Leinweber, A. Williams (Eds.), Lattice Hadron Physics
- Vol.664: R. Wielebinski, R. Beck (Eds.), Cosmic Magnetic Fields
- Vol.665: V. Martinez (Ed.), Data Analysis in Cosmology
- Vol.666: D. Britz, Digital Simulation in Electrochemistry
- Vol.667: W. D. Heiss (Ed.), Quantum Dots: a Doorway to Nanoscale Physics
- Vol.668: H. Ocampo, S. Paycha, A. Vargas (Eds.), Geometric and Topological Methods for Quantum Field Theory
- Vol.669: G. Amelino-Camelia, J. Kowalski-Glikman (Eds.), Planck Scale Effects in Astrophysics and Cosmology
- Vol.670: A. Dinklage, G. Marx, T. Klinger, L. Schweikhard (Eds.), Plasma Physics
- Vol.671: J.-R. Chazottes, B. Fernandez (Eds.), Dynamics of Coupled Map Lattices and of Related Spatially Extended Systems
- Vol.672: R. Kh. Zeytounian, Topics in Hypersonic Flow Theory
- Vol.673: C. Bona, C. Palenzuela-Luque, Elements of Numerical Relativity
- Vol.674: A. G. Hunt, Percolation Theory for Flow in Porous Media
- Vol.675: M. Kröger, Models for Polymeric and Anisotropic Liquids
- Vol.676: I. Galanakis, P. H. Dederichs (Eds.), Half-metallic Alloys
- Vol.677: A. Loiseau, P. Launois, P. Petit, S. Roche, J.-P. Salvetat (Eds.), Understanding Carbon Nanotubes
- Vol.678: M. Donath, W. Nolting (Eds.), Local-Moment Ferromagnets
- Vol.679: A. Das, B. K. Chakrabarti (Eds.), Quantum Annealing and Related Optimization Methods
- Vol.680: G. Cuniberti, G. Fagas, K. Richter (Eds.), Introducing Molecular Electronics
- Vol.681: A. Llor, Statistical Hydrodynamic Models for Developed Mixing Instability Flows
- Vol.682: J. Souchay (Ed.), Dynamics of Extended Celestial Bodies and Rings
- Vol.683: R. Dvorak, F. Freistetter, J. Kurths (Eds.), Chaos and Stability in Planetary Systems
- Vol.684: J. Dolinšek, M. Vilfan, S. Žumer (Eds.), Novel NMR and EPR Techniques
- Vol.685: C. Klein, O. Richter, Ernst Equation and Riemann Surfaces
- Vol.686: A. D. Yaghjian, Relativistic Dynamics of a Charged Sphere
- Vol.687: J. W. LaBelle, R. A. Treumann (Eds.), Geospace Electromagnetic Waves and Radiation
- Vol.688: M. C. Miguel, J. M. Rubi (Eds.), Jamming, Yielding, and Irreversible Deformation in Condensed Matter
- Vol.689: W. Pötz, J. Fabian, U. Hohenester (Eds.), Quantum Coherence
- Vol.690: J. Asch, A. Joye (Eds.), Mathematical Physics of Quantum Mechanics
- Vol.691: S. S. Abdullaev, Construction of Mappings for Hamiltonian Systems and Their Applications
- Vol.692: J. Frauendiener, D. J. W. Giulini, V. Perlick (Eds.), Analytical and Numerical Approaches to Mathematical Relativity
- Vol.693: D. Alloin, R. Johnson, P. Lira (Eds.), Physics of Active Galactic Nuclei at all Scales
- Vol.694: H. Schwoerer, J. Magill, B. Beleites (Eds.), Lasers and Nuclei
- Vol.695: J. Dereziński, H. Siedentop (Eds.), Large Coulomb Systems
- Vol.696: K.-S. Choi, J. E. Kim, Quarks and Leptons From Orbifolded Superstring
- Vol.697: E. Beaupaire, H. Bulou, F. Scheurer, J.-P. Kappler (Eds.), Magnetism: A Synchrotron Radiation Approach
- Vol.698: S. Bellucci (Ed.), Supersymmetric Mechanics – Vol. 1
- Vol.699: J.-P. Rozelot (Ed.), Solar and Heliospheric Origins of Space Weather Phenomena
- Vol.700: J. Al-Khalili, E. Roeckl (Eds.), The Euroschool Lectures on Physics with Exotic Beams, Vol. II
- Vol.701: S. Bellucci, S. Ferrara, A. Marrani, Supersymmetric Mechanics – Vol. 2
- Vol.702: J. Ehlers, C. Lämmerzahl, Special Relativity
- Vol.703: M. Ferrario, G. Ciccotti, K. Binder (Eds.), Computer Simulations in Condensed Matter Systems Volume 1
- Vol.704: M. Ferrario, G. Ciccotti, K. Binder (Eds.), Computer Simulations in Condensed Matter Systems Volume 2
- Vol.705: P. Bhattacharyya, B.K. Chakrabarti (Eds.), Modelling Critical and Catastrophic Phenomena in Geoscience
- Vol.706: M.A.L. Marques, C.A. Ullrich, F. Nogueira, A. Rubio, K. Burke, E.K.U. Gross (Eds.), Time-Dependent Density Functional Theory
- Vol.707: A.V. Shepetilov, Calculus and Mechanics on Two-Point Homogenous Riemannian Spaces
- Vol.708: F. Iachello, Lie Algebras and Applications